

isib/ms/2005/10  
February 21st, 2005  
<http://www.isibang.ac.in/~statmath/eprints>

# Degree-one maps, surgery and four-manifolds

SIDDHARTHA GADGIL

Indian Statistical Institute, Bangalore Centre  
8th Mile Mysore Road–560 059, India



# DEGREE-ONE MAPS, SURGERY AND FOUR-MANIFOLDS

SIDDHARTHA GADGIL

ABSTRACT. We give a description of degree-one maps between closed, oriented 3-manifolds in terms of surgery. Namely, we show that there is a degree-one map from a closed, oriented 3-manifold  $M$  to a closed, oriented 3-manifold  $N$  if and only if  $M$  can be obtained from  $N$  by surgery about a link in  $N$  each of whose components is an unknot.

We use this to interpret the existence of degree-one maps between closed 3-manifolds in terms of smooth 4-manifolds. More precisely, we show that there is a degree-one map from  $M$  to  $N$  if and only if there is a smooth embedding of  $M$  in  $W = (N \times I) \#_n \overline{\mathbb{C}P^2} \#_m \mathbb{C}P^2$ , for some  $m \geq 0$ ,  $n \geq 0$  which separates the boundary components of  $W$ . This is motivated by the relation to topological field theories, in particular the invariants of Ozsvath and Szabo.

## 1. INTRODUCTION

We assume that all manifolds are connected and that all 3-manifolds are smooth. For closed, oriented 3-manifolds  $M$  and  $N$ , we say that  $M$  *dominates*  $N$  (or  $M$  1-dominates  $N$ ) if there is a degree-one map from  $M$  to  $N$ . This gives a transitive relation on closed, oriented 3-manifolds which has been extensively studied by several authors (for instance, see [1], [5], [10], [11], [12], [13], [15]). Note that every manifold dominates  $S^3$  and that if  $M$  dominates  $N$  then there is a surjection from  $\pi_1(M)$  to  $\pi_1(N)$ .

In this paper, we characterise dominance in terms of Dehn surgery. We use this to interpret dominance in terms of smooth 4-manifolds. The latter is motivated by the relation to topological field theories, in particular the invariants of Ozsvath and Szabo [8][9].

Suppose  $N$  is a closed 3-manifold and  $M$  is obtained from  $N$  by surgery about a link in  $N$  all of whose components are homotopically trivial, then it is easy to see that there is a degree-one map from  $M$  to  $N$ . Our first result is the converse, namely that if there is a degree-one map from  $M$  to  $N$ , then  $M$  can be obtained from  $N$  by surgery about a link  $L \subset N$  all of whose components are homotopically trivial. In fact we can find  $L$  each of whose components is an unknots.

**Theorem 1.1.** *For closed oriented 3-manifolds  $M$  and  $N$ , there is a degree-one map from  $M$  to  $N$  if and only if  $M$  can be obtained from  $N$  by surgery about a link in  $N$  each of whose components is an unknot in  $N$ .*

We next interpret dominance of 3-manifolds in terms of 4-manifolds. Observe that a partial ordering on closed orientable 3-manifolds can be defined by saying that  $M$  *strongly dominates*  $N$  if there is a smooth embedding  $i: M \rightarrow N \times (0, 1) \subset N \times [0, 1]$  so that  $i(M)$  separates the two boundary components  $N \times \{0\}$  and  $N \times \{1\}$ , with the *appropriate orientation*. Observe that if  $M$  strongly dominates  $N$ , the the composition  $\pi \circ i$  of the embedding  $i$  with the projection  $\pi: N \times [0, 1] \rightarrow N$  has degree  $\pm 1$ . We say that the embedding has the *appropriate orientation* if the degree of this map is one.

This definition is motivated by the relation to ((3+1)-dimensional) *topological field theories*, in particular the invariants of Ozsvath and Szabo. Recall that a degree-one map  $f: M \rightarrow N$  induces a surjection

---

*Date:* February 25, 2005.

*1991 Mathematics Subject Classification.* Primary 57N10 ; Secondary 57N13, 57M27.

*Key words and phrases.* Degree-one maps, Dehn Surgery, Topological Field theories.

$f_*$  on the fundamental group. Hence if  $\pi_1(N)$  is non-trivial so is  $\pi_1(M)$ . Further  $f^*: H^*(N) \rightarrow H^*(M)$  is an injection, which shows that if  $H^k(N) \neq 0$  then  $H^k(M) \neq 0$ .

We see that an analogous result holds for any topological field theory, with dominance replaced by strong dominance. Recall that a  $(3+1)$ -dimensional topological field theory associates to each closed, oriented 3-manifold  $M$  a vector space  $V(M)$  and to each cobordism  $W$  from  $M$  to another closed, oriented 3-manifold  $N$  a linear transformation  $T(W): V(M) \rightarrow V(N)$ . Further this satisfies functorial properties, namely a product cobordism induces the identity map and if  $W_1$  is a cobordism from  $M_1$  to  $M_2$  and  $W_2$  is a cobordism from  $M_2$  to  $M_3$  then for the cobordism  $W_1 \amalg_{M_2} W_2$  from  $M_1$  to  $M_3$ ,  $T(W_1 \amalg_{M_2} W_2) = T(W_2) \circ T(W_1)$ .

Suppose  $M$  strongly dominates  $N$ , then splitting  $N \times [0, 1]$  along the given embedding of  $M$  gives two cobordisms,  $W_1$  from  $N$  to  $M$  and  $W_2$  from  $M$  to  $N$ . The composition of these is the product cobordism  $N \times [0, 1]$ , which induced the identity map on  $T(N)$ . It follows that the identity map on  $V(N)$  factors through  $V(M)$ , and in particular  $V(N) \neq 0$  implies that  $V(M) \neq 0$ . This is the analogue of the corresponding results for  $\pi_1$  and  $H^*$  with respect to degree-one maps. Thus *strong dominance* plays the same role in the *bordism category* as dominance in the homotopy category.

We shall see that relation of strong dominance is stronger than dominance. In particular we shall see in Theorem 4.1 that for lens spaces  $L(p, q)$  and  $L(p, q')$ , if  $L(p, q)$  strongly dominates  $L(p, q')$  then  $L(p, q)$  and  $L(p, q')$  are homeomorphic. This holds even if the embedding is only assumed to be topologically locally flat (rather than smooth). However, there are degree-one maps, in fact homotopy equivalences, between lens spaces which are not homeomorphic (for example  $L(7, 1)$  and  $L(7, 2)$ ). We get more subtle examples using Gauge theory, namely the Poincaré homology sphere does not (smoothly) strongly dominate  $S^3$ . However there is a topologically locally flat embedding of the Poincaré homology sphere in  $S^3 \times [0, 1]$ .

We show, however, that dominance is equivalent to a relation obtained using 4-manifolds similar to the above one except that we allow ‘positive and negative blow-ups’.

**Theorem 1.2.** *For closed orientable 3-manifolds  $M$  and  $N$ , there is a degree-one map from  $M$  to  $N$  if and only if there is a smooth embedding of  $M$  in  $\text{int}(W)$ ,  $W = (N \times I) \#_n \overline{\mathbb{C}P^2} \#_m \mathbb{C}P^2$  for some  $m > 0$ ,  $n > 0$  which separates the boundary components of  $W$ , with the embedding having the appropriate orientation.*

There is a relation in between dominance and strong dominance which is of interest. Namely, we say that  $M$  *negatively dominates*  $N$  if there is an embedding of  $M$  into  $W = (N \times I) \#_n \overline{\mathbb{C}P^2}$ , for some  $n \geq 0$ , which separates the two boundary components. This is of interest because the Ozsvath-Szabo invariants (as also the Seiberg-Witten invariants) behave well under blowing up.

We shall see (in Theorem 4.2) that the Poincaré homology sphere does not even negatively dominate  $S^3$ . We shall study negative dominance elsewhere.

## 2. DEGREE-ONE MAPS AND SURGERY

In this section, we give a proof of Theorem 1.1. We first recall the (well known) easy part of the result (see, for example, [1]). Namely, let  $M$  be obtained from  $N$  by surgery about a link  $L \subset N$  each of whose components  $K_i$  is homotopically trivial. Then we have a corresponding link  $L' \subset M$  with components  $K'_i$ , and an identification  $M - \text{int}(Nbd(L')) = N - \text{int}(Nbd(L))$ . We construct a degree-one map  $f: M \rightarrow N$  as follows. Define  $f$  on  $M - \text{int}(Nbd(L')) = N - \text{int}(Nbd(L))$  to be the identity map. We extend this to each solid torus  $X_i = Nbd(K'_i) = D^2 \times S^1$ .

Fix an identification  $X_i = D^2 \times S^1$ ,  $D^2 = \{z \in \mathbb{C} : |z| \leq 1\}$  and let  $X'_i \subset X_i$  be the solid torus  $\{z \in \mathbb{C} : |z| \leq 1/2\} \times S^1$ . Let  $T_i = \partial X'_i$ . We identify  $A_i = T_i \times [1/2, 1] = X_i - \text{int}(X'_i)$  with  $S^1 \times S^1 \times [1/2, 1]$  so that the curves of the form  $S^1 \times \{p\} \times \{1\}$  are mapped by  $f$  to meridians of  $Nbd(K_i)$ . Then  $f$  extends to a map on  $A_i$ , with image contained in  $Nbd(K_i)$ , so that curves of the form  $S^1 \times \{p\} \times \{1/2\}$  are mapped to points on  $K_i$ .

Now identify  $X'_i$  with  $D^2 \times S^1$ , which gives a new identification of  $T_i$  with  $S^1 \times S^1$ . Under this identification, we can assume that curves of the form  $S^1 \times \{p\}$  are mapped to  $K_i = S^1$  by a smooth map  $h$ , which we identify with a map  $h: S^1 \rightarrow K$ . As  $K$  is homotopic to the trivial curve, this extends to a smooth map  $h: D^2 \rightarrow K$ . Define  $f$  on  $X'_i = D^2 \times S^1$  by  $f(x, y) = h(x)$ . Thus the map  $f$  is defined on all of  $M$ . We see that this has degree one as any point in  $N$  disjoint from  $Nbd(L) \cup \text{image}(h)$  is a regular value for  $f$  and has inverse image a single point  $P$ , with  $f$  the identity on a neighbourhood of  $P$ .

The converse is based on the following theorem of Haken [4] and Waldhausen [14] (see also [12]).

**Theorem** (Haken-Waldhausen). *Let  $f: M \rightarrow N$  be a degree-one map and let  $N = H_1 \cup H_2$  be a Heegaard decomposition of  $N$  with  $H_1$  and  $H_2$  handlebodies. Then  $f$  is homotopic to a map  $g$  such that  $g|_{g^{-1}(H_1)}: g^{-1}(H_1) \rightarrow H_1$  is a homeomorphism.*

Such a map is called a 1-pinch. Thus if  $M$  dominates  $N$ , there is a 1-pinch  $g: M \rightarrow N$ .

*Proof of Theorem 1.1.* Assume  $M$  dominates  $N$  and let  $g, H_1$  and  $H_2$  be as above. We shall first show that  $M$  can be obtained from  $N$  by surgery about a link each of whose components is homotopically trivial.

Consider a collection of properly embedded discs  $D_i$ ,  $1 \leq i \leq n$  in  $H_2$  such that on splitting  $H_2$  along  $D_i$ , we get a 3-ball  $B$ . We can assume that  $g$  is transversal to  $D_i$  for all  $i$ ,  $1 \leq i \leq n$ . Let  $F_i = g^{-1}(D_i)$  and let  $K = g^{-1}(H_2)$ . Note that  $g|_{\partial H_2}$  is a homeomorphism and hence  $F_i$  consists of a compact surface with a single boundary component and a (possibly empty) collection of closed surfaces. First, note that by performing a homotopy of  $g$  we can assume that each  $F_i$  is connected. This follows (as the induced map on  $\pi_1$  is a surjection) by using standard techniques using binding ties as in Stallings's proof of the Kneser conjecture (see for example the proof of Kneser's conjecture in [6]). Hence  $F_i$ ,  $1 \leq i \leq n$ , is a compact surface with a single boundary component.

We first consider the special case when each  $F_i$  is a disc. After a homotopy of  $g$ , we can assume that  $F_i$  maps homeomorphically onto  $D_i$  for  $1 \leq i \leq n$ . On splitting  $K$  along the properly embedded discs  $F_i$ ,  $1 \leq i \leq n$ , we get a manifold  $K'$  with boundary a 2-sphere. By the theorem of Lickorish and Wallace, this can be obtained from  $B$  by surgery about a link  $L$  in  $B$ . Note that each component of  $L$  is contained in the 3-ball  $B$  and hence is homotopically trivial. Indeed the Lickorish-Wallace theorem says that we can take each component to be an unknot.

We now consider the general case. For each  $i$ ,  $1 \leq i \leq n$ , consider a collection  $L_i$  of disjoint, embedded simple closed curves on  $F_i$  which do not separate  $F_i$  and are maximal with respect to this property. Then  $L' = \cup_i L_i$  is a link in  $M$ . We consider a corresponding framed link (also denoted  $L'$ ), with the framing of a component of  $L_i$  given by the normal to  $F_i$ . Let  $M'$  be the manifold obtained from  $M$  by surgery about the framed link  $L'$ .

We shall see that the map  $g$  induces a degree-one map  $g'$  from  $M'$  to  $N$  with  $g'^{-1}(D_i)$  obtained from  $F_i$  by compressing along the components of  $L_i$ . By the choice of  $L_i$  it follows that  $g'^{-1}(D_i)$  is a disc for all  $i$ ,  $1 \leq i \leq n$ .

Let the components of  $L_i$  be  $C_j^i$  and let  $T_j^i$  denote a regular neighbourhood of  $C_j^i$ . On the complement of  $\bigcup_{i,j} \text{int}(T_j^i)$ , we let  $g' = g$ . After surgery, each  $T_j^i$  is replaced by a solid torus  $X = X_j^i$  with the

same boundary as  $T_j^i$ . Now,  $F_i \cap \partial X$  consists of two meridional curves (on  $\partial X$ ) which bound properly embedded discs  $E_1$  and  $E_2$  in  $X$ . As  $g(\partial E_i) \subset D_i$ , the map  $g'$  extends to  $E_i$  with  $g'(E_i) \subset D_i$ . Using transversality of  $g$  to  $D_i$ , we see that we can extend  $g'$  to a regular neighbourhood  $E_i \times [-1, 1]$  of  $E_i$  with  $g'(E_i \times [-1, 1] - E_i) \cap D_i = \emptyset$ . Finally,  $X - (E_1 \times (-1, 1)) - (E_2 \times (-1, 1))$  is the union of two balls, each of whose boundaries is mapped by  $g'$  into  $B$ . Thus  $g'$  extends to a map on  $X$  with  $(g'|_T)^{-1}(D_i) = E_1 \cup E_2$ . Making this construction for each  $X_j^i$ , we get a map  $g'$  as claimed.

Now  $g'$  is a pinch which is as in the special case, i.e.  $g'^{-1}(D_i)$  is a disc for all  $i$ ,  $1 \leq i \leq n$ . Thus  $M'$  can be obtained from  $N$  by surgery about a link  $L_0 \subset N$ , each of whose components is homotopically trivial in  $N$ . We now perform surgeries about knots  $\gamma_j^i \subset M'$  so that the surgery about  $\gamma_j^i$  cancels the surgery about  $C_j^i \subset L'$ . Thus on performing such surgeries we obtain  $M$ . The knots  $\gamma_j^i$  can be regarded as knots in  $H_2 \subset N$  and their union, with framing corresponding to the cancelling surgeries, is a framed link  $L_1 \subset N$ .

Thus  $M$  is obtained from  $N$  by surgery about the framed link  $L = L_0 \cup L_1$ , with each component of  $L_0$  homotopically trivial. We next show that the knots  $\gamma_j^i$ , regarded as curves in  $N$ , are trivial. Recall that, as  $\gamma_j^i$  is the knot corresponding to the surgery cancelling the surgery about  $C_j^i$ , it is given by an unknot bounding a disc that intersects  $C_j^i$  transversally in a single point and is contained in a ball around a point in  $C_j^i$ . Hence we can assume that  $\gamma_j^i$  intersects  $F_i$  transversally in two points, with opposite signs of intersection, and  $\gamma_j^i$  is disjoint from  $F_k$  for  $k \neq i$ .

Note that  $\pi_1(H_2)$  is a free group with generators  $\alpha_i$ ,  $1 \leq i \leq n$ , corresponding to the discs  $D_i$ . Further, if  $\gamma$  is a curve transversal to the discs  $D_i$ ,  $1 \leq i \leq n$ , then (up to conjugacy) the word represented by  $\gamma$  is determined by the intersection points with the discs  $D_i$ . Namely, if the points of  $\gamma \cap (\cup_i D_i)$ , in cyclic order around  $\gamma$ , are contained in  $D_{i_1}, \dots, D_{i_k}$  with signs of intersection  $\epsilon_j = \pm 1$ , then  $\gamma = \alpha_{i_1}^{\epsilon_1} \dots \alpha_{i_k}^{\epsilon_k}$  up to conjugacy.

As  $\gamma_j^i$  intersects  $F_i$  (hence  $D_i$ ) transversally in two points, with opposite signs of intersection, and  $\gamma_j^i$  is disjoint from  $F_k$  (hence  $D_k$ ) for  $k \neq i$ , it follows that  $\gamma_j^i$  represents the trivial word in  $\pi_1(H_2)$ , and hence is homotopically trivial in  $N$ .

Now as each component of  $L$  is homotopically trivial, there is a sequence of crossing changes so that on performing these crossing changes we obtain a link all of whose components are unknots. Observe that each crossing change of a knot  $\kappa$  is locally of a standard form. Namely, there is a ball  $B \subset M$  which intersects  $\kappa$  in a pair of arcs  $c_1$  and  $c_2$ , and the crossing change corresponds to a crossing of these arcs to give new arcs  $c'_1$  and  $c'_2$  with the same endpoints as  $c_1$  and  $c_2$ .

Further, if  $K_i$  is an unknot in  $B$  unlinked from the arcs  $c_i$  with framing  $\pm 1$ , then on performing the Kirby moves of sliding  $c_1$  and  $c_2$  over  $K_i$ , with opposite orientations, we get the knot obtained by crossing  $c_1$  and  $c_2$ . In this manner we can obtain both positive and negative crossing changes.

Note that replacing  $L$  by its union with unknots and performing the Kirby moves as above does not change the resulting manifold. Thus, we can replace  $L$  by a framed link in  $N$ , each of whose components is an unknot, so that the result of surgery about the link is  $M$ .

□

### 3. SURGERY AND 4-MANIFOLDS

We now characterise dominance in terms of 4-manifolds.

*Proof of Theorem 1.2.* Suppose  $M$  embeds in  $W$  as in the hypothesis. Then  $W - M$  has two components with closures  $K_1$  and  $K_2$  so that  $\partial[K_1] = [M] - [N \times \{0\}]$ . Hence  $[M]$  is homologous to  $[N \times \{0\}]$ . Now by identifying all the points in each  $\mathbb{C}P^2$  and  $\overline{\mathbb{C}P^2}$  in  $W = (M \times I) \#_n \overline{\mathbb{C}P^2} \#_m \mathbb{C}P^2$  to a single point, we get a blow-down map  $\pi: W \rightarrow N \times [0, 1]$ . By composing with the projection, we get a map

$p: W \rightarrow N$  with  $p: N \times \{0\} \rightarrow N$  being the identity map. This restricts to a map  $p: M \rightarrow N$ . As  $[M]$  is homologous to  $[N \times \{0\}]$ ,  $p_*([M]) = [N]$ , i.e.,  $M$  has degree one.

Conversely, assume  $M$  and  $N$  are as in the hypothesis with  $\pi_1(M) = 1$ . By Theorem 1.1,  $M$  can be obtained from  $N$  by surgery about a framed link  $L$ , all of whose components are unknots in  $N$ . Hence  $L$  can be obtained from an unlink  $L_0 \subset N$  by a sequence of (say  $p$ ) crossings.

Let  $K_1, \dots, K_n$  a collection of unknots in  $M$ , with  $n \geq p$  to be specified later, so that  $L_0 \cup \{K_1, \dots, K_n\}$  forms an unlink. Let  $W$  be obtained by attaching a 2-handle with framing  $\pm 1$  (with signs to be chosen later) to  $M \times [0, 1/2]$  along each of  $K_0, K_1, \dots, K_n$ . Note that  $W = (M \times [0, 1]) \#_k \overline{\mathbb{C}P^2} \#_l \mathbb{C}P^2$  for some  $k$  and  $l$ .

We shall construct a different Kirby diagram for  $W$ . Corresponding to the  $p$  crossings of  $L_0$  required to make it isotopic to  $L$  we can find disjoint balls  $B_i$ ,  $1 \leq i \leq p$ , in which the crossing is made. By an isotopy, we can assume that for  $1 \leq i \leq p$ ,  $K_i$  is contained in  $B_i$ . Performing the Kirby moves corresponding to the crossing changes in each of these  $B_i$ , we get a Kirby diagram for  $W$  with a sublink isotopic to  $L$ . Furthermore, by performing the Kirby move of sliding over the unknots  $K_{p+1}, \dots, K_n$  (with framing  $\pm 1$ ) we can ensure this sublink is isotopic to  $L$  as a framed link (as such a Kirby move changes the framing by  $\pm 1$  without changing the link). Consider the corresponding Morse function for  $W$  with the 2-handles corresponding to components of  $L$  attached first. The level set on attaching  $L$  is the result of surgery about  $L$ . But this is  $M$ , and hence we get an embedding of  $M$  separating the boundary components of  $W$ .  $\square$

#### 4. EXAMPLES

We now consider the relation of strong dominance where we allow no blowing-up. We see that we have restrictions in this case even for topologically locally flat embeddings.

**Theorem 4.1.** *If the lens space  $L(p, q)$  (topologically) strongly dominates the lens space  $L(p, q')$  for a prime  $p$  then  $L(p, q)$  and  $L(p, q')$  are homeomorphic.*

*Proof.* By hypothesis we have an embedding of  $L(p, q)$  in  $L(p, q') \times [0, 1]$  separating the two boundary components. Splitting along  $L(p, q)$ , we get a cobordism from  $L(p, q')$  to  $L(p, q)$ . By the Seifert-Van Kampen theorem and a Mayer-Vietoris argument we see that this is an h-cobordism. By a result of Milnor [7], it follows that  $L(p, q)$  and  $L(p, q')$  are homeomorphic.  $\square$

More subtle obstructions can be obtained using Gauge theory. Any 3-manifold dominates the 3-sphere. Furthermore, by a theorem of Freedman [3], we have a topological locally flat embedding of any homology sphere into  $S^3 \times [0, 1]$ . However in the smooth case, we see that  $S^3$  is not a minimal element with respect to strong dominance or even negative dominance.

**Theorem 4.2.** *For  $n \geq 0$ , there is no embedding of the Poincaré homology sphere in  $(S^3 \times I) \#_n \overline{\mathbb{C}P^2}$  which separates the boundary components.*

*Proof.* Note that the Poincaré homology sphere can be obtained from  $S^3$  by surgery on the  $E_8$  link, and hence, with one of its orientations, bounds a 4-manifold  $W$  with positive definite intersection pairing. Denote the Poincaré homology sphere with this orientation as  $M$ .

Suppose, for some  $n \geq 0$ , there is an embedding of the Poincaré homology sphere in  $(S^3 \times I) \#_n \overline{\mathbb{C}P^2}$  which separates the boundary components. Then by capping off the boundary components  $S^3 \times \{0\}$  and  $S^3 \times \{1\}$ , we get an embedding of the Poincaré homology sphere in  $\#_n \overline{\mathbb{C}P^2}$ . Splitting along the embedding and using the Mayer-Vietoris sequence, we get 4-manifolds  $W_1$  and  $W_2$  bounding  $M$  and  $-M$  with positive definite intersection forms. If  $\partial W_1 = -M$ , then  $Y = W \amalg_M W_1$  is a smooth 4-manifold

with  $H_1(Y, \mathbb{Z}) = 0$  and the intersection form on  $H_2(Y, \mathbb{Z})$  being positive definite but not diagonal, contradicting Donaldson's theorem [2].

□

## REFERENCES

1. Boileau, Michel; Wang, Shicheng *Non-zero degree maps and surface bundles over  $S^1$* . J. Differential Geom. **43** (1996), 789–806.
2. Donaldson, S. K. *An application of gauge theory to four-dimensional topology*. J. Differential Geom. 18 (1983), no. 2, 279–315.
3. Freedman, Michael Hartley *The topology of four-dimensional manifolds*. J. Differential Geom. **17** (1982), 357–453.
4. Haken, Wolfgang *On homotopy 3-spheres*. Illinois J. Math. **10** (1966) 159–178.
5. Hayat-Legend, Claude; Wang, Shicheng; Zieschang, Heiner *Any 3-manifold 1-dominates at most finitely many 3-manifolds of  $S^3$ -geometry*. Proc. Amer. Math. Soc. **130** (2002), 3117–3123.
6. Hempel, John *3-Manifolds*. Ann. of Math. Studies, No. **86**. Princeton University Press, Princeton, 1976.
7. Milnor, J. *Whitehead torsion* Bull. Amer. Math. Soc. **72** (1966) 358–426.
8. Ozsvath, Peter; Szabo, Zoltan *Holomorphic discs and topological invariants for closed three-manifolds*, to appear in the Annals of Mathematics.
9. Ozsvath, Peter; Szabo, Zoltan *Holomorphic triangles and invariants of smooth four-manifolds*, preprint
10. Reid, Alan W.; Wang, Shicheng *Non-Haken 3-manifolds are not large with respect to mappings of non-zero degree*. Comm. Anal. Geom. **7** (1999), 105–132.
11. Rong, Yong Wu *Degree one maps between geometric 3-manifolds*. Trans. Amer. Math. Soc. **332** (1992), 411–436.
12. Rong, Yong Wu; Wang, Shi Cheng *The preimages of submanifolds* Math. Proc. Cambridge Philos. Soc. **112** (1992), 271–279.
13. Soma, Teruhiko *Non-zero degree maps to hyperbolic 3-manifolds*. J. Differential Geom. **49** (1998), 517–546.
14. Waldhausen, Friedhelm *On mappings of handlebodies and of Heegaard splittings* Topology of Manifolds (Proc. Inst., Univ. of Georgia, Athens, Ga., 1969) 205–211
15. Wang, Shicheng; Zhou, Qing *Any 3-manifold 1-dominates at most finitely many geometric 3-manifolds*. Math. Ann. **322** (2002), 525–535.

STAT MATH UNIT,, INDIAN STATISTICAL INSTITUTE,, BANGALORE 560059, INDIA

*E-mail address:* [gadgil@isibang.ac.in](mailto:gadgil@isibang.ac.in)