

isibang/ms/2006/10

April 4th, 2006

<http://www.isibang.ac.in/~statmath/eprints>

# New constructions of homogeneous operators

GADHADHAR MISRA AND ADAM KORÁNYI

Indian Statistical Institute, Bangalore Centre  
8th Mile Mysore Road, Bangalore, 560059 India



# NEW CONSTRUCTIONS OF HOMOGENEOUS OPERATORS

ADAM KORÁNYI AND GADADHAR MISRA

ABSTRACT. New examples of homogeneous operators involving infinitely many parameters are constructed. They are realized on Hilbert spaces of holomorphic functions with reproducing kernels which are computed explicitly. All the examples are irreducible and belong to the Cowen - Douglas class. Even though the construction is completely explicit, it is based on certain facts about Hermitian holomorphic homogeneous vector bundles. These facts also make possible a description of all homogeneous Cowen - Douglas operators, in a somewhat less explicit way.

An operator  $T$  on a Hilbert space is said to be *homogeneous* if its spectrum is contained in the closure of the unit disc  $\mathbb{D}$  in  $\mathbb{C}$  and if  $g(T)$  is unitarily equivalent to  $T$  for every element  $g$  of the holomorphic automorphism group  $G$  of  $\mathbb{D}$ . There are general results about such operators, but relatively few examples are known (cf. [7, 2, 3, 1]). In this note a large family of examples is constructed and a step is made towards the description of all such operators in the Cowen - Douglas class of  $\mathbb{D}$  (see [4]).

## 1. CONSTRUCTION OF THE EXAMPLES

Let  $\tilde{G}$  denote the universal covering group of  $G$ . For  $\lambda > 0$ , let  $A^{(\lambda)}$  denote the Hilbert space of holomorphic functions on  $\mathbb{D}$  with reproducing kernel  $(1 - z\bar{w})^{-2\lambda}$ . The well-known discrete series  $D_\lambda^+$  of unitary representation of  $\tilde{G}$  acts on  $A^{(\lambda)}$  by

$$(D_\lambda^+(g)f)(z) = \left(\frac{\partial(g^{-1}z)}{\partial z}\right)^\lambda f(g^{-1}(z))$$

with the power defined to ensure continuity on  $\tilde{G} \times \mathbb{D}$ .

Let  $m \geq 1$  be an integer and let  $\lambda > \frac{m}{2}$ . For  $0 \leq j \leq m$  we write  $\lambda_j = \lambda - \frac{m}{2} + j$  and define the operator  $\Gamma_j : A^{(\lambda_j)} \rightarrow \text{Hol}(\mathbb{D}, \mathbb{C}^{m+1})$  by

$$(\Gamma_j f)_\ell = \begin{cases} 0 & \text{if } \ell < j \\ \binom{\ell}{j} \frac{1}{(2\lambda_j)_{\ell-j}} f^{(\ell-j)} & \text{if } \ell \geq j \end{cases}.$$

Denote by  $\mathbf{A}^{(\lambda_j)}$  the image of  $\Gamma_j$ . The algebraic sum of the spaces  $\mathbf{A}^{(\lambda_j)}$  can be shown to be direct. Hence, for any  $\mu_1, \dots, \mu_m$  and  $\mu_0 = 1$  we can define a norm on the direct sum by

$$\left\langle \sum_{j=0}^m \Gamma_j f_j, \sum_{j=0}^m \Gamma_j g_j \right\rangle = \sum_{j=0}^m \mu_j^2 \langle f_j, g_j \rangle, \quad f_j, g_j \in A^{(\lambda_j)}.$$

We denote the Hilbert space obtained in this way by  $\mathbf{A}^{(\lambda, \boldsymbol{\mu})}$  (we write  $\boldsymbol{\mu} = (\mu_0, \mu_1, \dots, \mu_m)$ ).

---

This research was supported in part by DST - NSF S&T Cooperation Programme.

The direct sum of maps  $\Gamma := \oplus \mu_j \Gamma_j$  is then a Hilbert space isomorphism of  $\oplus A^{(\lambda_j)}$  onto  $\mathbf{A}^{(\lambda, \mu)}$ . This isomorphism transfers the representation  $\oplus D_{\lambda_j}^+$  to  $\mathbf{A}^{(\lambda, \mu)}$ ; we denote its image by  $U$ .

For  $g \in \tilde{G}$ , we have  $g''(z) = -2c g'(z)^{3/2}$  with a constant  $c$  depending on  $g$ . We use  $c$  with this meaning in the following theorem.

**Theorem 1.1.**  *$U$  is a multiplier representation of  $\tilde{G}$ , that is,  $(U(g)f)(z) = J(g^{-1}, z)f(g^{-1}(z))$  for  $g \in \tilde{G}$ ,  $z \in \mathbb{D}$ . The matrix entries of the multiplier are given by*

$$J(g, z)_{p, \ell} = \begin{cases} 0 & \text{if } p < \ell \\ \binom{p}{\ell} (-c)^{p-\ell} g'(z)^{\lambda - \frac{m}{2} + \frac{p+\ell}{2}} & \text{if } p \geq \ell. \end{cases}$$

The theorem is proved by direct computation.

It is well known that in each  $A^{(\lambda_j)}$  the monomials, appropriately normalized, form an orthonormal basis. Using the isomorphism  $\Gamma$  we obtain an orthonormal basis of  $\mathbf{A}^{(\lambda, \mu)}$ .

Explicitly, the basis is  $\{\mu_j e_n^j\}$ ,  $0 \leq j \leq m$ ,  $n \geq 0$ , where the  $\ell$ 'th component of  $e_n^j$  is

$$e_n^j(z)_\ell = \begin{cases} 0 & \text{if } \ell < j \text{ or } \ell > n + j \\ \binom{\ell}{j} \frac{\sqrt{n!}}{(n-\ell+j)!} \frac{\sqrt{(2\lambda_j)_n}}{(2\lambda_j)_{\ell-j}} z^{(n-\ell+j)} & \text{if } j \leq \ell \leq n + j. \end{cases}$$

Here  $(x)_p := x(x+1) \cdots (x+p-1)$  is the Pochhammer symbol. It follows that the polynomials form a dense set in  $\mathbf{A}^{(\lambda, \mu)}$ . Therefore, the linear operator  $M = M^{(\lambda, \mu)}$  on  $\mathbf{A}^{(\lambda, \mu)}$  defined by

$$(Mf)(z) = zf(z)$$

is densely defined. To find the expression of  $M$  in terms of our orthonormal basis, we define the subspace  $H(n)$  as the linear span of the vectors  $\{e_{n-j}^j : 0 \leq j \leq \min(n, m)\}$  for each  $n \geq 0$ . Clearly  $M$  maps  $H(n)$  to  $H(n+1)$ . We have

$$M \mu_j e_{n-j}^j = \sum_{k=0}^m M(n)_{k,j} \mu_k e_{n+1-k}^k.$$

We define the number  $E(n)_{\ell j}$  by  $(e_{n-j}^j(z))_\ell = E(n)_{\ell j} z^{n-\ell}$  and write  $E(n)$  for the matrix  $E(n)_{\ell j}$ . We define the diagonal matrix  $D(\mu)$  by  $D(\mu)_{\ell j} = \mu_\ell \delta_{\ell j}$ . Now we have for  $n \geq m$

$$M(n) = D(\mu)^{-1} E(n+1)^{-1} E(n) D(\mu),$$

(with a small modification for  $0 \leq n < m$ ). Using Stirling's formula one verifies that, as  $n \rightarrow \infty$ ,

$$(1) \quad M(n) \sim I_{m+1} + O\left(\frac{1}{n}\right),$$

where  $O\left(\frac{1}{n}\right)$  stands for an  $(m+1) \times (m+1)$  matrix whose entries are  $O\left(\frac{1}{n}\right)$ .

**Theorem 1.2.** *The operator  $M = M^{(\lambda, \mu)}$  defined on  $\mathbf{A}^{(\lambda, \mu)}$  by  $(Mf)(z) = zf(z)$  belongs to the Cowen - Douglas class and is homogeneous.*

For the proof one remarks that by (1), the operator  $M$  splits into the sum of two operators. The first one is the direct sum of  $(m+1)$  copies of the standard isometric shift operator and

the second one belongs to the Hilbert-Schmidt class. This implies that it is bounded and is in the Cowen - Douglas class.

Using Theorem 1.1 one can verify, in a standard way, that  $g(M) = U_g^{-1}MU_g$  proving that  $M$  is homogeneous.

## 2. INEQUIVALENCE

**Theorem 2.1.** *For every  $m \geq 1$ , the operators  $M^{(\lambda, \boldsymbol{\mu})}$ ,  $\lambda > \frac{m}{2}; \mu_1, \dots, \mu_m > 0$  are mutually unitarily inequivalent.*

One essential ingredient of the proof is the following.

**Theorem 2.2.** *Suppose  $E$  is a Hermitian holomorphic vector bundle over  $\mathbb{D}$  and for every  $g \in G$  there exists an automorphism  $\hat{g}$  of  $E$  whose action on  $\mathbb{D}$  coincides with  $g$ . Then  $\hat{G}$  acts on  $E$  by automorphisms in a unique way.*

In the proof one considers the connected component  $\hat{G}$  of the full automorphism group of  $E$ . There is a natural homomorphism  $\hat{G} \rightarrow G$  which has a compact kernel  $N$ . One shows that  $\hat{G}$  is reductive so it contains a normal subgroup which is a covering of  $G$ .

For the proof of Theorem 2.1 we consider the representation of the Lie algebra  $\mathfrak{g}$  induced by  $U$  and extend it to the complexification  $\mathfrak{g}^{\mathbb{C}} = \mathfrak{sl}(2, \mathbb{C})$ , still denoting it by  $U$ .

Restricting  $U$  to the triangular subalgebra  $\mathfrak{t}$  of  $\mathfrak{g}^{\mathbb{C}}$  (the Lie algebra of the stabilizer of 0 in  $SL(2, \mathbb{C})$  acting on the extended complex plane), direct computation using Theorem 1.1 shows that  $(U(X)f)(0) = \rho_{\lambda, \boldsymbol{\mu}}(X)f(0)$  for all  $X \in \mathfrak{t}$  and  $f \in \mathbf{A}^{(\lambda, \boldsymbol{\mu})}$ , with  $\rho_{\lambda, \boldsymbol{\mu}}(X)$  independent of  $f$ , and therefore, giving a representation of  $\mathfrak{t}$  on the finite dimensional Hilbert space  $\mathbb{C}^{m+1}$ . The unitary equivalence class of  $\rho_{\lambda, \boldsymbol{\mu}}$  is uniquely determined by  $U$ , and hence by the operator  $M^{(\lambda, \boldsymbol{\mu})}$ .

Explicit computation of  $\rho_{\lambda, \boldsymbol{\mu}}$  shows that different pairs  $(\lambda, \boldsymbol{\mu})$  give inequivalent representations, proving the theorem.

## 3. REPRODUCING KERNEL AND IRREDUCIBILITY

In the following theorem  $S$  stands for the matrix with entries  $S_{\ell p} = \delta_{p+1, \ell} \ell$ ,  $T$  its transpose.  $D = D(z\bar{w})$  is diagonal with entries  $D_{\ell\ell} = (1 - z\bar{w})^{m-\ell}$ . Let  $K^{\lambda, \boldsymbol{\mu}}$  be the diagonal matrix with

$$(K^{\lambda, \boldsymbol{\mu}})_{\ell\ell} = \sum_{j=0}^m \mu_j^2 \sum_{j=0}^{\ell} \binom{\ell}{j}^2 \frac{(\ell-j)!}{(2\lambda_j)^{\ell-j}}.$$

**Theorem 3.1.** *The space  $\mathbf{A}^{(\lambda, \boldsymbol{\mu})}$  has a reproducing kernel given by*

$$\mathbf{B}^{(\lambda, \boldsymbol{\mu})}(z, w) = (1 - z\bar{w})^{-2\lambda-m} D(z\bar{w}) e^{\bar{w}S} K^{\lambda, \boldsymbol{\mu}} e^{zT} D(z\bar{w}).$$

*The normalized reproducing kernel in the sense of [5] is*

$$\mathbf{B}_0^{(\lambda, \boldsymbol{\mu})}(z, w) = e^{-zT} (K^{\lambda, \boldsymbol{\mu}})^{-1} \mathbf{B}^{(\lambda, \boldsymbol{\mu})}(z, w) (K^{\lambda, \boldsymbol{\mu}})^{-1} e^{-\bar{w}S}.$$

This is proved by computing the effect of the map  $\Gamma$  on the well known reproducing kernels of the spaces  $A^{(\lambda_j)}$  and using the identity

$$(2) \quad \mathbf{B}^{(\lambda, \boldsymbol{\mu})}(z, w) = J(g, z)\mathbf{B}^{(\lambda, \boldsymbol{\mu})}(gz, gw)J(g, w)^*, \quad g \in \tilde{G}.$$

**Theorem 3.2.** *The operators  $M^{(\lambda, \boldsymbol{\mu})}$ ,  $\lambda > \frac{m}{2}; \mu_1, \dots, \mu_m > 0$  are irreducible.*

One proves this by first refining some arguments of [5] to show that any orthogonal projection commuting with  $M$  corresponds to an orthogonal projection  $P$  in  $\mathbb{C}^{m+1}$  such that  $P\mathbf{B}_0^{(\lambda, \boldsymbol{\mu})}(z, w) = \mathbf{B}_0^{(\lambda, \boldsymbol{\mu})}(z, w)P$  for all  $z, w \in \mathbb{D}$ . Such a  $P$  then commutes with all coefficients of the power series development  $\mathbf{B}_0^{(\lambda, \boldsymbol{\mu})}$  at  $z = w = 0$ . Using the explicit expressions in Theorem 5 one can show that no non-trivial  $P$  can exist.

#### 4. MORE GENERAL RESULTS

Every Hermitian holomorphic vector bundle homogeneous under  $\tilde{G}$  can be obtained by the process of holomorphic induction [6, Ch. 13] from representations of  $\mathfrak{t}$  on  $\mathbb{C}^{m+1}$  that are Hermitian on the real reductive part of  $\mathfrak{t}$ . Unitary equivalence classes of such representations are in one-to-one correspondence with isomorphism classes of bundles. The  $\rho_{\lambda, \boldsymbol{\mu}}$  occurring above in Section 2 are certain conjugates of the restriction of the standard  $(m+1)$ -dimensional irreducible representation of  $\mathfrak{sl}(2, \mathbb{C})$  to  $\mathfrak{t}$  tensored with a one-dimensional representation. Any finite dimensional representation of  $\mathfrak{t}$  can be written as a tensor product of a one-dimensional representation  $\rho_\lambda$  characterized by a real parameter  $\lambda$  and another one  $\rho_0$  normalized in some way.

Even though the results are less explicit in the general case than for the operators constructed in Sections 1 and 2, we can prove the following.

**Theorem 4.1.** *To any  $\rho_0$  there corresponds a number  $\lambda_0$  such that the homogeneous bundle induced by  $\rho_\lambda \otimes \rho_0$  for  $\lambda > \lambda_0$  is the Cowen - Douglas bundle of a homogeneous operator.*

It is also possible to make statements about irreducibility and inequivalence of such operators.

A number of results in this Note extend to the case of operator tuples of Cowen - Douglas class on bounded symmetric domains in several complex variables.

Details will appear elsewhere.

#### REFERENCES

1. B. Bagchi and G. Misra, *Homogeneous tuples of multiplication operators on twisted Bergman space*, J. Funct. Anal. **136** (1996), 171 – 213.
2. B. Bagchi and G. Misra, *Homogeneous operators and projective representations of the Möbius group: a survey*, Proc. Ind. Acad. Sc.(Math. Sci.) **111** (2001), 415–437.
3. ———, *The homogeneous shifts*, J. Funct. Anal. **204** (2003), 293 –319.
4. M. J. Cowen and R. G. Douglas, *Complex geometry and operator theory*, Acta Math. **141** (1978), 187–261.
5. R. E. Curto and N. Salinas, *Generalized Bergman kernels and the Cowen - Douglas theory*, Amer J. Math. **106** (1984), 447–488.

6. A. A. Kirillov, *Elements of the theory of representations*, Grundlehren der Mathematischen Wissenschaften, Band 220. Springer-Verlag, Berlin-New York, 1976.
7. D. R. Wilkins, *Homogeneous vector bundles and Cowen - Douglas operators*, International J. of Math. **4** (1993), 503 – 520.

LEHMAN COLLEGE, THE CITY UNIVERSITY OF NEW YORK, BRONX, NY 10468

*E-mail address:* `adam.koranyi@lehman.cuny.edu`

INDIAN STATISTICAL INSTITUTE, R. V. COLLEGE POST, BANGALORE 560 059

*E-mail address:* `gm@isibang.ac.in`