

isibang/ms/2013/34
December 21st, 2013
<http://www.isibang.ac.in/~statmath/eprints>

Expansive automorphisms of totally disconnected, locally compact groups

HELGE GLÖCKNER AND C.R.E. RAJA

Indian Statistical Institute, Bangalore Centre
8th Mile Mysore Road, Bangalore, 560059 India

Expansive automorphisms of totally disconnected, locally compact groups

Helge Glöckner and C.R.E. Raja

Abstract

We study automorphisms α of a totally disconnected, locally compact group G which are *expansive* in the sense that $\bigcap_{n \in \mathbb{Z}} \alpha^n(U) = \{1\}$ for some identity neighbourhood $U \subseteq G$. Notably, we prove that the automorphism induced by α on a quotient group G/N of G modulo an α -stable closed normal subgroup N is always expansive. Further results involve the contraction groups $U_\alpha := \{g \in G : \alpha^n(g) \rightarrow 1 \text{ as } n \rightarrow \infty\}$. If α is expansive, then $U_\alpha U_{\alpha^{-1}}$ is an open identity neighbourhood in G . We give examples where $U_\alpha U_{\alpha^{-1}}$ fails to be a subgroup. However, $U_\alpha U_{\alpha^{-1}}$ is an α -stable, nilpotent open subgroup of G if G is a closed subgroup of $\mathrm{GL}_n(\mathbb{Q}_p)$. Further results are devoted to the divisible and torsion parts of U_α , and to the so-called “nub” $U_0 = \overline{U_\alpha} \cap \overline{U_{\alpha^{-1}}}$ of an expansive automorphism.

Classification: 22D05 (primary); 22D45, 22E20, 37A25, 37P20 (secondary)

Key words: Totally disconnected group, locally compact group, locally profinite group, contraction group, expansive automorphism, contractive automorphism, tidy subgroup, Willis’s theory, p -adic Lie group, finite depth, composition series, simple factor, Baumslag-Solitar group, Schlichting completion, nub, closedness, normalizer

Introduction and statement of results

We consider automorphisms $\alpha: G \rightarrow G$ of a totally disconnected locally compact topological group G which are *expansive* in the sense that $\bigcap_{n \in \mathbb{Z}} \alpha^n(U) = \{1\}$ for some identity neighbourhood $U \subseteq G$. Expansive automorphisms of totally disconnected, *compact* groups were studied by [12], [20] and recently in [28]; our goal is to improve the understanding in the case of non-compact groups. Special cases of expansive automorphisms are automorphisms $\alpha: G \rightarrow G$ which are *contractive* in the sense that $\alpha^n(g) \rightarrow 1$ as $n \rightarrow \infty$ for each $g \in G$ (see Remark 1.10). The structure of totally disconnected, locally compact groups admitting contractive automorphisms was

elucidated in [9] (building on earlier work like [22] and [24]), and the results obtained there can also be used as tools in the investigation of expansive automorphisms (as we shall see). Other key ingredients are the structure theory of totally disconnected, locally compact groups ([26], [27]), in which contractive automorphisms play an important role (as worked out in [1]).

Note that every automorphism α of a discrete group G (e.g., $\alpha = \text{id}_G$) is expansive (as we may choose $U = \{1\}$ then). Therefore discrete groups and their automorphisms are part of the theory of expansive automorphisms. As a consequence, groups permitting expansive automorphisms need not have any particular algebraic properties.

Our first main result generalizes [28, Proposition 6.1] (devoted to the case of compact groups). As usual, a subset $H \subseteq G$ is called α -stable if $\alpha(H) = H$.

Theorem A. *Let $\alpha: G \rightarrow G$ be an automorphism of a totally disconnected, locally compact group, $N \subseteq G$ be an α -stable closed normal subgroup and $\bar{\alpha}: G/N \rightarrow G/N$, $gN \mapsto \alpha(g)N$ be the automorphism of G/N induced by α . Then α is expansive if and only if both $\alpha|_N$ and $\bar{\alpha}$ are expansive.*

Composition series play a central role in the study of contractive automorphisms [9]. In the case of expansive automorphisms, composition series need not exist. However, there are series which get as close as possible to such.

Theorem B. *If $\alpha: G \rightarrow G$ is an expansive automorphism of a totally disconnected, locally compact group G , then there exist α -stable closed subgroups*

$$G = G_0 \supseteq G_1 \supseteq \cdots \supseteq G_n = \{1\}$$

of G such that G_j is normal in G_{j-1} for $j \in \{1, \dots, n\}$ and each of the quotient groups G_{j-1}/G_j is discrete, abelian or topologically perfect. Moreover, one can achieve that every α_j -stable closed normal subgroup of G_{j-1}/G_j is discrete or open, where $\alpha_j: G_{j-1}/G_j \rightarrow G_{j-1}/G_j$, $gG_j \mapsto \alpha(g)G_j$.

In addition, one may assume that all abelian, non-discrete factors G_{j-1}/G_j are simple contraction groups with respect to the automorphism or its inverse, or isomorphic to an infinite power $C_p^{\mathbb{Z}}$ of a cyclic group of prime order, endowed with the right-shift (cf. Remark 6.1 and Proposition 6.2). Topological perfectness means that the commutator group is dense.

According to [27], a compact open subgroup $V \subseteq G$ is called *tidy for α* if it has the following properties:

T1 $V = V_+V_-$, where $V_+ := \bigcap_{n=0}^{\infty} \alpha^n(V)$ and $V_- := \bigcap_{n=0}^{\infty} \alpha^{-n}(V)$;

T2 The α -stable subgroups $V_{++} := \bigcup_{n \in \mathbb{N}_0} \alpha^n(V_+)$ and $\bigcup_{n \in \mathbb{N}_0} \alpha^{-n}(V_-) =: V_{--}$ are closed in G .

Note that $V_+ \subseteq \alpha(V_+) \subseteq \alpha^2(V_+) \subseteq \dots$ and $V_- \subseteq \alpha^{-1}(V_-) \subseteq \alpha^{-2}(V_-) \subseteq \dots$ here. Following [28], the intersection U_0 of all subgroups V which are tidy for α is called the *nub* of α ; it is a compact, α -stable subgroup of G .

We mention that a totally disconnected, locally compact group admitting an expansive automorphism α is always metrizable (cf. [13]) and has a second countable, α -stable open subgroup (Lemma 1.1 (a)).

If $\alpha: G \rightarrow G$ is an automorphism of a totally disconnected, locally compact group G , then

$$U_\alpha := \{g \in G: \alpha^n(g) \rightarrow 1 \text{ as } n \rightarrow \infty\}$$

is a subgroup of G called the associated *contraction group*. In general, U_α need not be closed. However, if α is expansive, then the topology on U_α “can be made locally compact,” i.e., it can be refined to a totally disconnected, locally compact group topology τ^* with respect to which $\alpha|_{U_\alpha}$ is contractive (see [23, Proposition 9] for this fact, or our Lemma 2.3). In this way, the structure theory of locally compact contraction groups (see [22], [24] and [9]) becomes available. In particular, the set

$$T_\alpha := \text{tor}(U_\alpha)$$

of all torsion elements in the group U_α and the set

$$D_\alpha := \text{div}(U_\alpha)$$

of divisible elements are α -stable closed subgroups of (U_α, τ^*) , and $(U_\alpha, \tau^*) = D_\alpha \times T_\alpha$ internally as a topological group, if we endow D_α and T_α with the topology induced by (U_α, τ^*) (see [9, Theorem B]).

Theorem C. *Let G be a totally disconnected, locally compact group and $\alpha: G \rightarrow G$ be an automorphism such that U_α can be made locally compact (for example, any expansive automorphism). Then $D_\alpha = \text{div}(U_\alpha)$ is an α -stable closed subgroup of G .*

The nub U_0 of an expansive automorphism need not have an open normalizer (Remark 7.1), in which case not both of U_α and $U_{\alpha^{-1}}$ normalize U_0 . Nonetheless, U_0 is always normalized (and even centralized) by both D_α and $D_{\alpha^{-1}}$.

Theorem D. *Let G be a totally disconnected, locally compact group and $\alpha: G \rightarrow G$ be an automorphism such that U_α can be made locally compact (for example, any expansive automorphism). Then the nub U_0 of α centralizes the divisible part D_α of U_α . Also $\overline{T_\alpha}$ centralizes D_α . Moreover, $D_\alpha \cap \overline{T_\alpha} = \{1\}$, and $\overline{U_\alpha} = D_\alpha \overline{T_\alpha}$ is a closed, α -stable subgroup of G which is isomorphic to $D_\alpha \times \overline{T_\alpha}$ as a topological group.*

We also consider classes of examples. If a Lie group G over a totally disconnected local field \mathbb{K} admits an expansive analytic automorphism, then its Lie algebra $L(G)$ is nilpotent (Proposition 8.1). In the case of p -adic Lie groups for a prime number p , we obtain:

Theorem E. *Let G be a p -adic Lie group which is linear in the sense that there exists an injective continuous homomorphism $G \rightarrow \mathrm{GL}_n(\mathbb{Q}_p)$ for some $n \in \mathbb{N}$. Let $\alpha: G \rightarrow G$ be an expansive automorphism. Then G has an open α -stable subgroup which is nilpotent.*

If $\alpha: G \rightarrow G$ is an expansive automorphism of a totally disconnected, locally compact group G , then $U_\alpha U_{\alpha^{-1}}$ is an open subset of G (see Proposition 1.1). In many examples, $U_\alpha U_{\alpha^{-1}}$ happens to be a subgroup of G (for instance, for all expansive automorphisms of closed subgroups $G \subseteq \mathrm{GL}_n(\mathbb{Q}_p)$, see Proposition 8.8). However, this is not always so which could be seen from Remark 8.7. We also consider the localized completion $G_{p,q}$ of a Baumslag-Solitar group $\mathrm{BS}(p, q) = \langle a, t: ta^p t^{-1} = a^q \rangle$ with primes $p \neq q$, as recently studied in [4]. Then $\mathrm{BS}(p, q) \subseteq G_{p,q}$. We show:

Theorem F. *Let $\alpha: G_{p,q} \rightarrow G_{p,q}$ be conjugation by t . Then α is expansive but $U_\alpha U_{\alpha^{-1}}$ is not a subgroup of $G_{p,q}$.*

Acknowledgement. The second author was supported by the German Academic Exchange Service (DAAD) and the second author wishes to thank DAAD and Institut für Mathematik, Universität Paderborn, Paderborn, Germany for providing the facilities during his stay.

1 Preliminaries and basic facts

We write $\mathbb{N} = \{1, 2, \dots\}$, $\mathbb{N}_0 := \mathbb{N} \cup \{0\}$ and $\mathbb{Z} := \mathbb{N}_0 \cup -\mathbb{N}$. If J is a finite set, we let $\#J$ be its cardinality. We write $X \subseteq Y$ for inclusion of sets, while $X \subset Y$ means that X is a proper subset of Y . As usual, we write $N \triangleleft G$ if N is a normal subgroup of G . All topological groups considered in this article are assumed Hausdorff, and locally compact topological groups are simply called locally compact groups. Totally disconnected, locally compact non-discrete topological fields (like the field of p -adic numbers) will be called local fields (see [25] for further information). See [3] and [21] for basic information on Lie groups over local (and more general complete ultrametric) fields (which we always assume finite-dimensional). If we say that α is an automorphism of a topological group, then we assume that both α and α^{-1} are continuous; similarly, both α and α^{-1} are assumed analytic if α is an automorphism of an analytic Lie group over a local field. We write $\text{Aut}(G)$ for the group of all automorphisms of a topological group G . If a subgroup $N \subseteq G$ is stable under all $\alpha \in \text{Aut}(G)$, then N is called *topologically characteristic*. A topological group G is called *topologically perfect* if its commutator group $[G, G]$ is dense in G . If F is a finite group and X a set, we write $F^X := \prod_{x \in X} F$ for the direct power endowed with the compact product topology. By contrast, $F^{(X)} \subseteq F^X$ is the subgroup of all $(g_x)_{x \in X} \in F^X$ such that $g_x = 1$ for all but finitely many $x \in X$. We shall always endow $F^{(X)}$ with the discrete topology. Surjective, open, continuous homomorphisms between topological groups are called *quotient morphisms*.

If $\alpha: G \rightarrow G$ is an automorphism of a locally compact group, choose a Haar measure λ on G and define the module of α via

$$\Delta_G(\alpha) := \lambda(\alpha(K))/\lambda(K),$$

for any compact subset $K \subseteq G$ with non-empty interior. If $\alpha: G \rightarrow G$ is an automorphism of a totally disconnected, locally compact group, we define the nub U_0 , the contraction group U_α and its subgroups T_α and D_α as explained in the introduction. If $V \subseteq G$ is a compact open subgroup, we shall use the subgroups V_+ , V_{++} , V_- and V_{--} defined there, and abbreviate

$$V_0 := V_+ \cap V_- = \bigcap_{n \in \mathbb{Z}} \alpha^n(V). \quad (1)$$

We shall also need the so-called *Levi factor*

$$M_\alpha := \{g \in G: \{\alpha^n(g): n \in \mathbb{Z}\} \text{ is relatively compact}\};$$

it is known that M_α is an α -stable closed subgroup of G [1, p.224]. The following lemma compiles basic facts concerning expansive automorphisms.

Lemma 1.1 *If α is an expansive automorphism of a totally disconnected, locally compact group G , then the following holds:*

- (a) G is metrizable and has an α -stable, σ -compact open subgroup;
- (b) $V_{--} = U_\alpha$ and $V_{++} = U_{\alpha^{-1}}$ for each compact open subgroup $V \subseteq G$ such that $V_0 = \{1\}$;
- (c) G has a compact open subgroup V such that $V = V_+V_-$ and $V_0 = \{1\}$;
- (d) $U_\alpha U_{\alpha^{-1}}$ is open in G ;
- (e) $\alpha|_H$ is expansive, for each α -stable subgroup $H \subseteq G$.

Proof. (a) Because α is expansive, there exists an identity neighbourhood V such that $\bigcap_{n \in \mathbb{Z}} \alpha^n(V) = \{1\}$. After shrinking V , we may assume that V is compact. Because V is compact and $(\bigcap_{k=-n}^n \alpha^k(V))_{k \in \mathbb{N}}$ a decreasing sequence of closed identity neighbourhoods in V with intersection $\{1\}$, the members of the sequence form a basis of identity neighbourhoods. Hence G is metrizable. The subgroup of G generated by $\bigcup_{n \in \mathbb{Z}} \alpha^n(V)$ is α -stable, open and σ -compact.

(b) By [1, Proposition 3.16], we have $V_{--} = U_\alpha V_0$ and thus $V_{--} = U_\alpha$. Likewise, $V_{++} = U_{\alpha^{-1}}$.

(c) and (d): Using expansiveness and van Dantzig's Theorem [10, Theorem 7.7], we find a compact open subgroup $W \subseteq G$ such that $\bigcap_{n \in \mathbb{Z}} \alpha^n(W) = \{1\}$. By [26, Lemma 1], there exists $m \in \mathbb{N}$ such that $V := \bigcap_{k=1}^m \alpha^k(W)$ satisfies $V = V_+V_-$. Then $V_0 \subseteq W_0 = \{1\}$ and thus $V_0 = \{1\}$, proving (c). The latter entails $U_\alpha = V_{--}$ and $U_{\alpha^{-1}} = V_{++}$, by (b). In particular, $V_- \subseteq U_\alpha$ and $V_+ \subseteq U_{\alpha^{-1}}$, entailing that $V = V_+V_- \subseteq U_\alpha U_{\alpha^{-1}}$. Thus $U_\alpha U_{\alpha^{-1}}$ is an identity neighbourhood. Given $g \in U_\alpha$ and $h \in U_{\alpha^{-1}}$, the map $G \rightarrow G$, $x \mapsto gxh$ is a homeomorphism which takes $U_\alpha U_{\alpha^{-1}}$ onto itself and 1 to gh . Hence $U_\alpha U_{\alpha^{-1}}$ has gh in its interior and thus $U_\alpha U_{\alpha^{-1}}$ is open.

(f) If $V \subseteq G$ is an identity neighbourhood with $\bigcap_{n \in \mathbb{Z}} \alpha^n(V) = \{1\}$, then $V \cap H$ is an identity neighbourhood in H and $\bigcap_{n \in \mathbb{Z}} \alpha^n(V \cap H) = \{1\}$. \square

The first statement of Lemma 1.1 (a) also follows from [13, Lemma 2.4].

1.2 Let α be an automorphism of a totally disconnected, locally compact group G and U_0 be the nub of α . The following facts are useful:

- (a) The closure of U_α in G is $\overline{U_\alpha} = U_\alpha U_0$ (see [1, Corollary 3.30] if G is metrizable; the general case follows with [11]).
- (b) $U_0 = \overline{U_\alpha} \cap \overline{U_{\alpha^{-1}}}$ (see [1, Corollary 3.27] if G is metrizable; the general case follows with [11]).
- (c) $U_0 \cap U_\alpha$ and $U_0 \cap U_{\alpha^{-1}}$ are dense in U_0 (see [28, Theorem 4.1 (v) and Proposition 5.4 (i)]).
- (d) If $N \subseteq G$ is a closed α -stable subgroup, $q: G \rightarrow G/N$ the canonical quotient morphism and $\bar{\alpha}$ the automorphism of G/N induced by α , then $q(U_\alpha) = U_{\bar{\alpha}}$ (see [1, Theorem 3.8] if G is metrizable, [11] in the general case).
- (e) U_α is closed if and only if $U_{\alpha^{-1}}$ is closed, if and only if G has small subgroups tidy for α , i.e., every identity neighbourhood of G contains some tidy subgroup (see [1, Theorem 3.32] if G is metrizable; the general case can be deduced using the techniques from [11]).
- (f) $P_\alpha := \{g \in G: \alpha^{\mathbb{N}_0}(g) \text{ is relatively compact}\}$ normalizes U_α (see [1, Proposition 3.4]). Hence $M_\alpha = P_\alpha \cap P_{\alpha^{-1}}$ and $U_0 \subseteq M_\alpha$ normalize U_α .

The following results help to show that certain automorphisms are expansive.

Proposition 1.3 *Let α be an automorphism of a totally disconnected, locally compact group G . Then the following holds:*

- (a) α is expansive if and only if its restriction $\alpha|_{M_\alpha}$ to the Levi factor is expansive.
- (b) If U_α is closed, then α is expansive if and only if $U_\alpha U_{\alpha^{-1}}$ is open in G , if and only if M_α is discrete.

Proof. (a) In view of Lemma 1.1 (e), we only need to show that if $\alpha|_{M_\alpha}$ is expansive, then so is α . Let $P \subseteq M_\alpha$ be a compact, open identity neighbourhood such that $\bigcap_{n \in \mathbb{Z}} \alpha^n(P) = \{1\}$. There is a compact identity neighbourhood $Q \subseteq G$ such that $Q \cap M_\alpha = P$. If $g \in I := \bigcap_{n \in \mathbb{Z}} \alpha^n(Q)$, then $\alpha^n(g) \in Q$ for each $n \in \mathbb{Z}$, whence $\alpha^{\mathbb{Z}}(g)$ is relatively compact and thus

$g \in M_\alpha$. Hence $I \subseteq M_\alpha$. Since I is α -stable and $I \subseteq Q \cap M_\alpha = P$, we deduce that $I = \bigcap_{n \in \mathbb{Z}} \alpha^n(I) \subseteq \bigcap_{n \in \mathbb{Z}} \alpha^n(P) = \{1\}$. Thus $\bigcap_{n \in \mathbb{Z}} \alpha^n(Q) = \{1\}$ and thus α is expansive.

(b) If α is expansive, then $U_\alpha U_{\alpha^{-1}}$ is open by Lemma 1.1 (d). If U_α is closed, then the set $U_\alpha M_\alpha U_{\alpha^{-1}}$ is open in G and the product map

$$U_\alpha \times M_\alpha \times U_{\alpha^{-1}} \rightarrow U_\alpha M_\alpha U_{\alpha^{-1}}, \quad (x, y, z) \mapsto xyz$$

is a homeomorphism (by 1.2 (e) above and part (f) from the theorem in [5]). Hence $U_\alpha U_{\alpha^{-1}} \cap M_\alpha = \{1\}$. If $U_\alpha U_{\alpha^{-1}}$ is open, this implies that M_α is discrete. If M_α is discrete, then $\alpha|_{M_\alpha}$ is expansive and hence also α , by (a). \square

1.4 If α is an automorphism of a totally disconnected, *compact* group G , then U_α and $U_{\alpha^{-1}}$ are normal in G (since $G = M_\alpha = M_{\alpha^{-1}}$ in 1.2 (f)), and hence so is the nub $U_0 = \overline{U_\alpha} \cap \overline{U_{\alpha^{-1}}}$.

1.5 Recall that a group G is called a torsion group *of finite exponent* if there exists $m \in \mathbb{N}$ such that $g^m = 1$ for all $g \in G$. If such a group is a subgroup of a topological group H , then also $g^m = 1$ for all g in the closure \overline{H} of G in H , and thus also \overline{H} is a torsion group of finite exponent.

1.6 Let α be an expansive automorphism of a totally disconnected, compact group G . Then the following holds:

- (a) The nub U_0 is open in G (see [28, Lemma 5.1]), whence also $\overline{U_\alpha}$ and $\overline{U_{\alpha^{-1}}}$ are open in G (by 1.2 (b)).
- (b) G is a torsion group of finite exponent [By (a) and 1.4, U_0 is a normal subgroup and G/U_0 is finite, hence a torsion group of finite exponent. It therefore suffices to show that U_0 is a torsion group of finite exponent. This is immediate from [28, Proposition 4.4, Theorem 6.2 and Proposition 6.3]].
- (c) Since U_α and $U_{\alpha^{-1}}$ are normal in G , the set $U_\alpha U_{\alpha^{-1}}$ is an open α -stable subgroup of G contained in U_0 . Thus $U_0 = U_\alpha U_{\alpha^{-1}}$ by [28, Corollary 4.3].

Lemma 1.7 *Let α be an automorphism of a totally disconnected, locally compact group G and U_0 be the nub of α . Let $H \subseteq G$ be a closed, α -stable subgroup. Then the following holds:*

(a) Then nub of $\alpha|_H$ is contained in U_0 .

(b) If $U_0 \subseteq H$, then the nub of $\alpha|_H$ coincides with U_0 .

Proof. (a) Let W be the nub of $\alpha|_H$. Using 1.2 (b) twice, we deduce that $W = \overline{U_{\alpha|_H}} \cap \overline{U_{(\alpha|_H)^{-1}}} \subseteq \overline{U_\alpha} \cap \overline{U_{\alpha^{-1}}} = U_0$.

(b) Since $U_0 \subseteq H$, we have $U_\alpha \cap U_0 \subseteq U_\alpha \cap H = U_{\alpha|_H}$ and thus $U_0 = \overline{U_0 \cap U_\alpha} \subseteq \overline{U_{\alpha|_H}}$ (using 1.2 (c)). Likewise, $U_0 \subseteq \overline{U_{(\alpha|_H)^{-1}}}$. Hence $U_0 \subseteq \overline{U_{\alpha|_H}} \cap \overline{U_{(\alpha|_H)^{-1}}} = W$. Since $W \subseteq U_0$ by (a), equality follows. \square

We shall also need certain facts concerning contractive automorphisms.

1.8 Let α be a contractive automorphism of a topological group G .

(a) If $G \neq \{1\}$, then G is infinite and non-discrete. [If $x \in G \setminus \{1\}$, then $\alpha^n(x) \neq 1$ for all n and $\alpha^n(x) \rightarrow 1$, entailing that the topological group G is not discrete and hence infinite.]

(b) If G is locally compact, then α is *compactly contractive*, i.e., for each identity neighbourhood $U \subseteq G$ and compact set $K \subseteq G$ there exists $m \in \mathbb{N}$ such that $\alpha^n(K) \subseteq U$ for all $n \geq m$ (see [22, Lemma 1.4 (iv)]). Moreover, G is non-compact (unless $G = \{1\}$); see [22, 3.1].

1.9 Let α be a contractive automorphism of a totally disconnected, locally compact group G . Then the following holds:

(a) If $G \neq \{1\}$, then $\Delta_G(\alpha^{-1})$ is an integer ≥ 2 (see [9, Proposition 1.1 (e)]).

(b) If $G = G_0 \supset G_1 \supset G_2 \supset \cdots \supset G_n = \{1\}$ is a series of α -stable closed subgroups of G such that G_j is a proper normal subgroup of G_{j-1} for all $j \in \{1, \dots, n\}$, then n is bounded by the number of prime factors of $\Delta_G(\alpha^{-1})$ (see [9, Lemma 3.5]).

(c) $G = \text{div}(G) \times \text{tor}(G)$ as a topological group for a divisible torsion free group $\text{div}(G)$ and a torsion group $\text{tor}(G)$ of finite exponent (cf. [9, Theorem B]).

(d) If $N \subseteq G$ is an α -stable closed normal subgroup, $q: G \rightarrow G/N$ the canonical quotient morphism and $\bar{\alpha}$ the automorphism of G/N induced by α , then $q(\text{div}(G)) = \text{div}(G/N)$ and $q(\text{tor}(G)) = \text{tor}(G/N)$. [The inclusions $q(\text{div}(G)) \subseteq \text{div}(G/N)$ $q(\text{tor}(G)) \subseteq \text{tor}(G/N)$ are clear. Since $G/N = q(\text{div}(G)\text{tor}(G)) = q(\text{div}(G))q(\text{tor}(G))$ and $G/N = \text{div}(G/N) \times \text{tor}(G/N)$, equality follows.]

Remark 1.10 If an automorphism $\alpha: G \rightarrow G$ of a totally disconnected, locally compact group is contractive, then it is also expansive. To see this, let $V \subseteq G$ be any compact neighbourhood of the identity. If $g \in G \setminus \{1\}$, then $G \setminus \{g\}$ is an identity neighbourhood, whence there exists $n \in \mathbb{N}$ such that $\alpha^n(V) \subseteq G \setminus \{g\}$ (see 1.8 (b)). Thus $g \notin \alpha^n(V)$ and we have shown that $\bigcap_{n \in \mathbb{Z}} \alpha^n(V) = \{1\}$.

2 Making contraction groups locally compact

The problem of refining group topologies on contraction groups was studied by Siebert [23]. The following special case is useful for our purposes.

Definition 2.1 Let G be a topological group, with topology τ , and

$$\alpha: (G, \tau) \rightarrow (G, \tau)$$

be a contractive automorphism. We say that (G, α) (or simply G) *can be made locally compact* if there exists a locally compact topology τ^* on G making it a topological group such that $\tau \subseteq \tau^*$ and $\alpha: (G, \tau^*) \rightarrow (G, \tau^*)$ is a contractive automorphism. The topology τ^* is unique (if it exists), as will be recalled presently. We write G^* for G , endowed with the topology τ^* . If τ is totally disconnected, then also τ^* is totally disconnected (as the inclusion map $G^* \rightarrow G$ is continuous).

The following fact is also obtained in [23, Corollary 8] as part of a more general theory.

Lemma 2.2 τ^* is uniquely determined by the properties from Definition 2.1.

Proof. Assume that $\hat{\tau}$ is a group topology on G with the same properties as τ^* . We show that the identity map

$$\phi: (G, \hat{\tau}) \rightarrow (G, \tau^*), \quad x \mapsto x$$

is continuous. Reversing the roles of $\hat{\tau}$ and τ^* , also ϕ^{-1} will be continuous and thus $\hat{\tau} = \tau^*$. Because ϕ is a homomorphism, we need only prove its continuity at 1. By local compactness, there exists a compact identity neighbourhood $V \subseteq (G, \hat{\tau})$. After replacing V with the closure of its interior V^0 , we may assume that V^0 is dense in V . Then V is also compact in (G, τ) . Let

$U \subseteq (G, \tau^*)$ be an arbitrary identity neighbourhood, and $W \subseteq (G, \tau^*)$ be a compact identity neighbourhood such that $WW^{-1} \subseteq U$. Then W is also compact in (G, τ) , hence closed in (G, τ) and hence closed in $(G, \hat{\tau})$. Since $\alpha: (G, \tau^*) \rightarrow (G, \tau^*)$ is contractive, we have $G = \bigcup_{n \in \mathbb{N}_0} \alpha^{-n}(W)$. Hence V is the countable union of the closed subsets $V \cap \alpha^{-n}(W)$, for $n \in \mathbb{N}_0$. By the Baire Category Theorem, there exists $m \in \mathbb{N}_0$ such that $V \cap \alpha^{-m}(W)$ has non-empty interior in V . Because V^0 is dense in V , we deduce that $\alpha^{-m}(W) \cap V^0$ has non-empty interior in V^0 , and so W has non-empty interior W^0 in $(G, \hat{\tau})$. Then $W^0(W^0)^{-1}$ is an identity neighbourhood in $(G, \hat{\tau})$ and hence U is an identity neighbourhood in $(G, \hat{\tau})$. Since $U = \phi^{-1}(U)$, we see that ϕ is continuous at 1. \square

See also [23, Proposition 9] for the following fact.

Lemma 2.3 *Let G be a totally disconnected, locally compact group. If an automorphism $\alpha: G \rightarrow G$ is expansive, then $(U_\alpha, \alpha|_{U_\alpha})$ and $(U_{\alpha^{-1}}, \alpha^{-1}|_{U_{\alpha^{-1}}})$ can be made locally compact.*

Proof. Let $V \subseteq G$ be a compact open subgroup such that $\bigcap_{n \in \mathbb{Z}} \alpha^n(V) = \{1\}$. Then $\alpha^n(V_-) = V_- \cap \bigcap_{k=1}^n \alpha^k(V)$ is open in V_- for each $n \in \mathbb{N}_0$. Since V_- is compact and $\bigcap_{n \in \mathbb{N}_0} \alpha^n(V_-) = \bigcap_{n \in \mathbb{Z}} \alpha^n(V) = \{1\}$, the open subgroups $\alpha^n(V_-)$ form a basis of identity neighbourhoods in V_- , for $n \in \mathbb{N}_0$. If $n \in \mathbb{N}$ and $g \in U_\alpha = V_{--} = \bigcup_{k \in \mathbb{N}_0} \alpha^{-k}(V_-)$ (see Proposition 1.1(c)), then $g \in \alpha^{-k}(V_-)$ for some $k \in \mathbb{N}_0$ and $\alpha^n(V_-)$ is an open subgroup of the topological group $\alpha^{-k}(V_-)$. Hence, there exists $m \in \mathbb{N}_0$ such that $g\alpha^m(V_-)g^{-1} \subseteq \alpha^n(V_-)$. By the preceding, there exists a group topology τ^* on U_α for which $\{\alpha^n(V_-): n \in \mathbb{N}_0\}$ is a basis of identity neighbourhoods. Thus V_- is an open subgroup of (U_α, τ^*) , and the latter group induces the given compact topology on V_- . Thus (U_α, τ^*) is a totally disconnected, locally compact group and α is still continuous (being continuous on the open subgroup V_-) as well as α^{-1} (being continuous on the subgroup $\alpha(V_-)$ which is open in V_-). Since $U_\alpha = \bigcup_{n \in \mathbb{N}_0} \alpha^{-n}(V_-)$ and $(\alpha^n(V_-))_{n \in \mathbb{N}_0}$ is a basis of identity neighbourhoods, it readily follows that the automorphism α of (U_α, τ^*) is contractive. \square

Remark 2.4 U_α can also be made locally compact if α is an arbitrary (not necessarily expansive) analytic automorphism of a Lie group G over a local field (see Proposition 13.3(b) in the extended preprint version of [7]).

Example 2.5 The right-shift α is an automorphism of the compact group $G := (\mathbb{Z}_p)^\mathbb{Z}$, where \mathbb{Z}_p is the additive group of p -adic integers. The contraction group U_α is non-trivial, as it is the group of all $(z_n)_{n \in \mathbb{Z}}$ such that $z_n \rightarrow 0$ as $n \rightarrow -\infty$. Then U_α cannot be locally compact, because $\text{tor}(U_\alpha) \subseteq \text{tor}(G) = \{0\}$ and $\text{div}(U_\alpha) \subseteq \text{div}(G) = \{0\}$. Thus, if U_α could be made locally compact, then we would have $U_\alpha = \text{div}(U_\alpha) + \text{tor}(U_\alpha) = \{0\}$, contradicting $U_\alpha \neq \{0\}$.

Lemma 2.6 *Let G be a topological group and $\alpha: G \rightarrow G$ a contractive automorphism such that (G, α) can be made locally compact. Then we have:*

- (a) *$(H, \alpha|_H)$ can be made locally compact for each closed α -stable subgroup H of G (or G^*), and H^* carries the topology induced by G^* .*
- (b) *If $\phi: G \rightarrow H$ is a continuous homomorphism to a topological group H admitting an automorphism $\beta: H \rightarrow H$ such that $\beta \circ \phi = \phi \circ \alpha$, then $\beta|_{\phi(G)}$ is contractive, $\phi(G)$ can be made locally compact and*

$$G^* \rightarrow \phi(G)^*, \quad x \mapsto \phi(x)$$

is a topological quotient map.

Proof. (a) Let τ and τ^* be as in Definition 2.1. Since $\tau \subseteq \tau^*$, H is closed in G^* (in either case) and hence H is a locally compact group in the topology σ on H induced by G^* , which is finer than the topology induced by G and turns $\alpha|_H$ into a contractive automorphism of (H, σ) . Thus $H^* = (H, \sigma)$.

(b) Let σ be the topology on $\phi(G)$ turning $G^* \rightarrow (\phi(G), \sigma)$, $x \mapsto \phi(x)$ into a quotient map. Then σ is finer than the topology induced on $\phi(G)$ by H . Moreover, $(\phi(G), \sigma) \cong G^*/\ker \phi$ is locally compact. Since $\beta|_{\phi(G)} \circ \phi = \phi \circ \alpha: G^* \rightarrow (\phi(G), \sigma)$ is continuous, the map $\beta|_{\phi(G)}$ is continuous with respect to the quotient topology σ . Also $(\beta|_{\phi(G)})^{-1}$ is continuous, by an analogous argument. Finally, $\beta|_{\phi(G)}: (\phi(G), \sigma) \rightarrow (\phi(G), \sigma)$ is contractive: Since $\beta^n \circ \phi = \phi \circ \alpha^n$, this follows from the facts that $\alpha: G^* \rightarrow G^*$ is contractive and $\phi: G^* \rightarrow \phi(G)^*$ is continuous. \square

The following observation is crucial for many of our arguments.

Proposition 2.7 *Let α be an expansive automorphism of a totally disconnected, locally compact group G and*

$$G = G_0 \supseteq G_1 \supseteq \cdots \supseteq G_n$$

be α -stable closed subgroups of G such that G_j is normal in G_{j-1} for all $j \in \{1, \dots, n\}$. Let J be the set of all $j \in \{1, \dots, n\}$ such that G_{j-1}/G_j is not discrete. Then

$$\#J \leq \ell_\alpha + \ell_{\alpha^{-1}},$$

where ℓ_α is the number of prime factors of $\Delta_{U_\alpha^*}(\alpha^{-1}|_{U_\alpha^*})$ and $\ell_{\alpha^{-1}}$ is the number of prime factors of $\Delta_{U_{\alpha^{-1}}^*}(\alpha|_{U_{\alpha^{-1}}^*})$.

Proof. Let J_α (resp., $J_{\alpha^{-1}}$) be the set of all $j \in \{1, \dots, n\}$ such that $U_\alpha \cap G_j \subset U_\alpha \cap G_{j-1}$ (resp., $U_{\alpha^{-1}} \cap G_j \subset U_{\alpha^{-1}} \cap G_{j-1}$). If $j \in \{1, \dots, n\} \setminus (J_\alpha \cup J_{\alpha^{-1}})$, then $U_\alpha \cap G_j = U_\alpha \cap G_{j-1}$ and $U_{\alpha^{-1}} \cap G_j = U_{\alpha^{-1}} \cap G_{j-1}$. Since α is expansive, $(U_\alpha \cap G_{j-1})(U_{\alpha^{-1}} \cap G_{j-1})$ is open in G_{j-1} (see 1.1 (d) and (e)). We deduce that G_j is open in G_{j-1} and thus $j \notin J$. Hence $J \subseteq J_\alpha \cup J_{\alpha^{-1}}$, whence $\#J \leq \#J_\alpha + \#J_{\alpha^{-1}} \leq \ell_\alpha + \ell_{\alpha^{-1}}$ (using 1.9 (b) in the last step). \square

We shall use a simple fact.

Lemma 2.8 . *Let K be a compact group and $D \subseteq K$ be a subgroup which is divisible. Then also the closure \overline{D} is divisible. If K is totally disconnected, then $D = \{1\}$.*

Proof. For each $m \in \mathbb{N}$, the map $f_m: \overline{D} \rightarrow \overline{D}$, $g \mapsto g^m$ is continuous and hence has compact image. As the image contains D by hypothesis, we see that $f_m(\overline{D}) = \overline{D}$. Thus \overline{D} is divisible.

If K is totally disconnected, then K is a pro-finite group. In particular, the homomorphisms $f: K \rightarrow F$ to finite groups F separate points on \overline{D} . But each $f(\overline{D})$ is both finite and divisible and therefore the trivial group. Hence also $\overline{D} = \{1\}$. \square

Lemma 2.9 *Let α be an automorphism of a totally disconnected, locally compact group G . If U_α can be made locally compact (e.g., if α is expansive), then the following holds:*

- (a) $U_\alpha \cap U_0 = T_\alpha \cap U_0$;
- (b) $U_0 = \overline{T_\alpha \cap U_0}$;
- (c) $\overline{T_\alpha} \cap U_\alpha = T_\alpha$;
- (d) $\overline{T_\alpha} = T_\alpha U_0$.

If both U_α and $U_{\alpha^{-1}}$ can be made locally compact, then we also have:

$$(e) \quad U_0 = \overline{T_\alpha} \cap \overline{T_{\alpha^{-1}}}.$$

Proof. (a) By Lemma 2.6 (a), $U_\alpha \cap U_0$ can be made locally compact. Thus α restricts to a contractive automorphism β of $(U_\alpha \cap U_0)^*$, enabling us to write $(U_\alpha \cap U_0)^* = D_\beta T_\beta$. Since U_0 is compact and totally disconnected, its divisible subgroup D_β has to be trivial, by Lemma 2.8. Thus $U_\alpha \cap U_0 = U_\beta = T_\beta \subseteq T_\alpha$ and hence $U_\alpha \cap U_0 = T_\alpha \cap U_0$.

(b) Since $U_0 = \overline{U_\alpha} \cap \overline{U_0}$ (see 1.2 (c)), the assertion is immediate from (a).

(c) $T_\alpha \subseteq \overline{T_\alpha} \cap U_\alpha$ is trivial. Because T_α is a torsion group of finite exponent (see 1.9 (c)), also $\overline{T_\alpha}$ is a torsion group, see 1.5. Let β the restriction of α to the closed α -stable subgroup $\overline{T_\alpha} \cap U_\alpha^*$ of U_α^* . Then $\overline{T_\alpha} \cap U_\alpha = D_\beta T_\beta$. Since D_β is torsion-free (see 1.9 (c)) and $\overline{T_\alpha}$ a torsion group, $D_\beta = \{1\}$ follows. Thus $\overline{T_\alpha} \cap U_\alpha = T_\beta \subseteq T_\alpha$.

(d) We have $\overline{T_\alpha} \subseteq \overline{U_\alpha} = U_\alpha U_0$ (see 1.2 (a)). Since $U_0 \subseteq \overline{T_\alpha}$ by (b), we deduce that $\overline{T_\alpha} = (U_\alpha \cap \overline{T_\alpha}) U_0 = T_\alpha U_0$ (using (c) for the last equality).

(e) We have $U_0 = \overline{U_\alpha} \cap \overline{U_{\alpha^{-1}}} \supseteq \overline{T_\alpha} \cap \overline{T_{\alpha^{-1}}}$ (see 1.2 (b)) and $U_0 = \overline{T_\alpha} \cap \overline{U_0} \cap \overline{T_{\alpha^{-1}} \cap U_0} \subseteq \overline{T_\alpha} \cap \overline{T_{\alpha^{-1}}}$ (using (b)). \square

Lemma 2.10 *Let α be an automorphism of a locally compact group G and $H \subseteq G$ be an α -stable subgroup such that $\alpha|_H$ is contractive. Then the closure $\overline{H} \subseteq G$ is σ -compact.*

Proof. Let $K \subseteq \overline{H}$ be a compact identity neighbourhood. Then $\bigcup_{n \in \mathbb{Z}} \alpha^n(K)$ is a σ -compact subset of \overline{H} and generates a σ -compact subgroup S of \overline{H} . Since S is an α -stable open subgroup of \overline{H} and $\alpha|_H$ is contractive, we have $H \subseteq S$ and thus $S = \overline{H}$ (since S is closed). Hence $\overline{H} = S$ is σ -compact. \square

3 Proof of Theorem A

Let G be a totally disconnected, locally compact group, α be an expansive automorphism of G and $N \subseteq G$ be an α -stable, closed normal subgroup. Let $q: G \rightarrow G/N$ be the canonical quotient morphism and $\bar{\alpha}$ be the automorphism of G/N induced by α (determined by $\bar{\alpha} \circ q = q \circ \alpha$). Then $\alpha|_N$ is

expansive, by Lemma 1.1 (e). We show also that $\bar{\alpha}$ is expansive. By Proposition 1.3 (a), we need only to show that $\bar{\alpha}$ restricts to an expansive automorphism of $M_{\bar{\alpha}}$. After replacing G with $q^{-1}(M_{\bar{\alpha}})$ (which is closed since $M_{\bar{\alpha}}$ is closed), we may assume that $G/N = M_{\bar{\alpha}}$. Let U be a subgroup of G/N tidy for $\bar{\alpha}$. Then $U = U_+ = U_-$ as $\bar{\alpha}^{\mathbb{Z}}(g)$ is relatively compact for each $g \in U$ (cf. [26, Lemma 9]) and thus U is an $\bar{\alpha}$ -stable, compact open subgroup of G/N . After replacing G with $q^{-1}(U)$, we may assume that G/N is compact. Using [28, Proposition 5.1] and the metrizability of G/N , we find a descending sequence $(H_n)_{n \in \mathbb{N}}$ of $\bar{\alpha}$ -stable closed normal subgroups H_n of G/N such that $\bar{\alpha}$ induces an expansive automorphism α_n on $(G/N)/H_n$ for each $n \in \mathbb{N}$ and G/N is the projective limit $G/N = \varprojlim (G/N)/H_n$. Set $L_n := q^{-1}(H_n)$; then $(L_n)_{n \in \mathbb{N}}$ is a descending sequence of α -stable closed normal subgroups of G , with $\bigcap_{n \in \mathbb{N}} L_n = N$.

There exists $m \in \mathbb{N}$ such that L_n is open in L_m for all $n \geq m$. Indeed, if this was wrong, we could find a subsequence $(L_{n_k})_{k \in \mathbb{N}}$ such that, for each $k \in \mathbb{N}$, the normal subgroup $L_{n_{k+1}}$ is not open in L_{n_k} . This contradicts Proposition 2.7.

After passing to a subsequence, we may assume that L_n is open in L_1 for each $n \in \mathbb{N}$. Hence L_n contains both $U_\alpha \cap L_1$ and $U_{\alpha^{-1}} \cap L_1$. As a consequence, $N = \bigcap_{n \in \mathbb{N}} L_n$ contains both $U_\alpha \cap L_1$ and $U_{\alpha^{-1}} \cap L_1$. Hence N is open in L_1 (and in each L_n), using that $\alpha|_{L_1}$ is expansive and thus $(U_\alpha \cap L_1)(U_{\alpha^{-1}} \cap L_1)$ an open subset of L_1 (see 1.1 (d) and (e)). This implies that the compact group $H_1 \cong L_1/N$ is discrete and hence a finite group. Since $H_1 \supseteq H_2 \supseteq \dots$ with $\bigcap_{n \in \mathbb{N}} H_n = \{1\}$, we deduce that $H_n = \{1\}$ for some n . Since $\bar{\alpha}$ corresponds to the expansive automorphism α_n on $(G/N)/H_n \cong G/N$, we see that $\bar{\alpha}$ is expansive.

Conversely, assume that both $\alpha|_N$ and $\bar{\alpha}: G/N \rightarrow G/N$ are expansive.¹ Then there is an open identity neighbourhood $P \subseteq G/N$ such that $\bigcap_{n \in \mathbb{Z}} \bar{\alpha}^n(P) = \{1\}$, and an open identity neighbourhood $Q \subseteq N$ such that $\bigcap_{n \in \mathbb{Z}} \alpha^n(Q) = \{1\}$. After shrinking Q , we may assume that $Q \subseteq q^{-1}(P)$. Then $Q = N \cap V$ for some open identity neighbourhood $V \subseteq G$. After replacing V with $V \cap q^{-1}(P)$, we may assume that $V \subseteq q^{-1}(P)$. We have $N = q^{-1}(\bigcap_{n \in \mathbb{Z}} \bar{\alpha}^n(P)) = \bigcap_{n \in \mathbb{Z}} \alpha^n(q^{-1}(P))$, entailing that $I := \bigcap_{n \in \mathbb{Z}} \alpha^n(V)$ is an α -stable subset of N .

¹Compare also [28, Proposition 6.1]. The compactness of G assumed there is inessential for this part of the proof of [28, Proposition 6.1].

Since $I = I \cap N \subseteq V \cap N = Q$, we deduce that $I = \bigcap_{n \in \mathbb{Z}} \alpha^n(I) \subseteq \bigcap_{n \in \mathbb{Z}} \alpha^n(Q) = \{1\}$. Hence $I = \{1\}$ and α is expansive. \square

4 Proof of Theorem B

The following lemma is useful.

Lemma 4.1 *Let α be an expansive automorphism of a totally disconnected, locally compact group G such that every α -stable, closed normal subgroup $N \subseteq G$ is open or discrete. Let $C \subseteq G$ be the topologically characteristic subgroup² of G generated by $U_\alpha \cup U_{\alpha^{-1}}$. Let $\overline{[C, C]}$ be the closure of the commutator group of C . Then C is an open normal subgroup of G , and one of the following cases occurs:*

- (a) $\overline{[C, C]}$ is open in G , in which case $C = \overline{[C, C]}$ is topologically perfect.
- (b) $\overline{[C, C]}$ is discrete.

Proof. C is an open subgroup of G as it contains the open set $U_\alpha U_{\alpha^{-1}}$ (see 1.1 (d)). The closed subgroup $\overline{[C, C]}$ is topologically characteristic in C , whence it is topologically characteristic in G and hence α -stable and normal. Therefore $\overline{[C, C]}$ is open or discrete. If $\overline{[C, C]}$ is open, then it contains $U_\alpha \cup U_{\alpha^{-1}}$. Hence $\overline{[C, C]} = C$, using that C is the smallest topologically characteristic subgroup of G which contains $U_\alpha \cup U_{\alpha^{-1}}$. \square

Proof of Theorem B. Define ℓ_α and $\ell_{\alpha^{-1}}$ as in Proposition 2.7. For every series

$$\Sigma: G = G_0 \triangleright G_1 \triangleright \cdots \triangleright G_n = \{1\}$$

of α -stable closed subgroups of G , let J_Σ be the set of all $j \in \{1, \dots, n\}$ such that G_{j-1}/G_j is not discrete. Then $J_\Sigma \leq \ell_\alpha + \ell_{\alpha^{-1}}$, by Proposition 2.7, entailing that the maximum

$$m := \max_{\Sigma} J_\Sigma$$

over all series Σ exists. Let $\Sigma: G = G_0 \triangleright G_1 \triangleright \cdots \triangleright G_n = \{1\}$ be a series with $J_\Sigma = m$. Let $N \subseteq G_{j-1}$ be an α -stable closed normal subgroup with $G_j \subseteq N$. If neither G_{j-1}/N nor N/G_j was discrete, we would have $J_{\Sigma \cup \{N\}} = J_\Sigma + 1$,

²Thus C is the subgroup generated by $\bigcup_{\beta \in \text{Aut}(G)} \beta(U_\alpha \cup U_{\alpha^{-1}})$.

a contradiction. Thus N will be open in G_{j-1} or G_j open in N .

For $j \in J_\Sigma$, let $q_j: G_{j-1} \rightarrow G_{j-1}/G_j$ be the canonical quotient morphism, $\alpha_j: G_{j-1}/G_j \rightarrow G_{j-1}/G_j$ be the automorphism induced by α and $C_j \subseteq G_{j-1}/G_j$ be the topologically characteristic subgroup generated by $U_{\alpha_j} \cup U_{\alpha_j^{-1}}$. If $\overline{[C_j, C_j]}$ is open in G_{j-1}/G_j , define $M_j := N_j := q_j^{-1}(C_j)$; thus $G_{j-1}/M_j \cong (G_{j-1}/G_j)/C_j$ and $M_j/N_j = \{1\}$ are discrete and $N_j/G_j \cong C_j$ is topologically perfect. If $\overline{[C_j, C_j]}$ is discrete, we define $M_j := q_j^{-1}(C_j)$ and $N_j := q_j^{-1}(\overline{[C_j, C_j]})$ (by Lemma 4.1, only these two cases can occur). Then $G_{j-1}/M_j \cong (G_{j-1}/G_j)/C_j$ is discrete, $M_j/N_j \cong C_j/\overline{[C_j, C_j]}$ is abelian and $N_j/G_j \cong \overline{[C_j, C_j]}$ is discrete. Hence

$$\Sigma' := \Sigma \cup \bigcup_{j \in J_\Sigma} \{M_j, N_j\}$$

is a series of α -stable closed subnormal subgroups such that all non-discrete subfactors are abelian or topologically perfect. Since $\#J_{\Sigma'} = \#J_\Sigma$ is maximal, all non-discrete subfactors of Σ' have the property that all stable closed normal subgroups are open or discrete. \square

5 Proof of Theorem C

After replacing G with the closure of D_α in G , we may assume that $G = \overline{D_\alpha}$. Since U_0 normalizes U_α (see 1.2(f)), we deduce from $G = \overline{U_\alpha} = U_\alpha U_0$ (see 1.2(a)) that all of G normalizes U_α . Thus $U_\alpha \triangleleft G$. Since $T_\alpha = \text{tor}(U_\alpha)$ is characteristic in U_α , we deduce that also T_α is a normal subgroup of G and hence also $\overline{T_\alpha}$. Let $q: G \rightarrow G/\overline{T_\alpha}$ be the canonical quotient morphism and $\bar{\alpha}: G/\overline{T_\alpha} \rightarrow G/\overline{T_\alpha}$ be the automorphism induced by α . Since T_α is a torsion group of finite exponent, also $\overline{T_\alpha}$ is a torsion group (see 1.5). Therefore $D_\alpha \cap \overline{T_\alpha} = \{1\}$, using that D_α is torsion-free (see 1.9(c)). Hence

$$q|_{D_\alpha}: D_\alpha \rightarrow G/\overline{T_\alpha}, \quad g \mapsto g\overline{T_\alpha}$$

is injective. Since $U_\alpha = D_\alpha T_\alpha$, using 1.2(d) we obtain $q(D_\alpha) = q(U_\alpha) = U_{\bar{\alpha}}$, where $U_{\bar{\alpha}}$ can be made locally compact by Lemma 2.6 (b). Being a homomorphic image of a divisible group, $U_{\bar{\alpha}}$ is divisible and hence coincides with $D_{\bar{\alpha}}$. If we can show that $D_{\bar{\alpha}}$ is closed, then

$$q|_{D_\alpha^*}: D_\alpha^* \rightarrow D_{\bar{\alpha}}$$

will be a continuous bijective homomorphism between σ -compact locally compact groups and hence an isomorphism of topological groups (see [10, Theorem 5.29]). After replacing G with $G/\overline{T_\alpha}$, we may therefore assume that $T_\alpha = \{1\}$, $U_\alpha = D_\alpha$ and D_α is dense in G . Since $U_0 = \overline{U_0} \cap \overline{T_\alpha} = \{1\}$ by Lemma 2.9 (b), we deduce with 1.2 (a) that $D_\alpha = U_\alpha = U_\alpha U_0 = G$ is closed in G . \square

Corollary 5.1 *Let G be a totally disconnected, locally compact group and α be an automorphism of G such that U_α and $U_{\alpha^{-1}}$ can be made locally compact (e.g., any expansive automorphism). Then $D_\alpha \cap D_{\alpha^{-1}} = \{1\}$.*

Proof. Since D_α and $D_{\alpha^{-1}}$ are closed and α -stable, their intersection $H := D_\alpha \cap D_{\alpha^{-1}}$ is a totally disconnected, locally compact contraction group for both $\alpha|_H$ and $\alpha^{-1}|_H$. Hence $H = \{1\}$. Indeed, if $H \neq \{1\}$ then both $\Delta_H(\alpha|_H)$ and $\Delta_H(\alpha|_H^{-1}) = \Delta_H(\alpha|_H)^{-1}$ would be integers ≥ 2 (see 1.9 (a)), which is impossible. \square

6 Abelian expansion groups

We show that, after passing to a refinement if necessary, only abelian, non-discrete groups of a special form will occur in Theorem B.

Remark 6.1 In the situation of Theorem B, let I be the set of all indices $j \in \{1, \dots, n\}$ such that G_{j-1}/G_j is abelian and non-discrete. Let α_j be the automorphism of G_{j-1}/G_j induced by α and $q_j: G_{j-1} \rightarrow G_{j-1}/G_j$ be the quotient homomorphism, for $j \in I$. Then $U_{\alpha_j}U_{\alpha_j^{-1}}$ is an open α_j -stable subgroup of G_{j-1}/G_j and hence $H_j := q_j^{-1}(U_{\alpha_j}U_{\alpha_j^{-1}})$ is an α -stable open normal subgroup of G_{j-1} . Then G_{j-1}/H_j is discrete and all stable, closed, proper subgroups of H_j/G_j are discrete. After inserting the H_j into the series for all $j \in I$, we may thus assume without loss of generality that all abelian, non-discrete subfactors G_{j-1}/G_j have the property that all of their α_j -stable, closed, proper subgroups are discrete, and that $G_{j-1}/G_j = U_{\alpha_j}U_{\alpha_j^{-1}}$.

Let G_j be a topological group and $\alpha_j \in \text{Aut}(G_j)$ for $j \in \{1, 2\}$. We say that (G_1, α_1) and (G_2, α_2) are isomorphic if there exists an isomorphism $\phi: G_1 \rightarrow G_2$ of topological groups such that $\alpha_2 \circ \phi = \phi \circ \alpha_1$.

Proposition 6.2 *Let $A \neq \{1\}$ be an abelian, totally disconnected, locally compact group and $\alpha: A \rightarrow A$ be an expansive automorphism. Assume that $A = U_\alpha U_{\alpha^{-1}}$ and assume that every α -stable proper closed subgroup of A is discrete. Then there exists a prime number p such that (A, α) is isomorphic to one of the following:*

- (a) \mathbb{Q}_p^n for some $n \in \mathbb{N}$, together with a contractive linear automorphism $\beta: \mathbb{Q}_p^n \rightarrow \mathbb{Q}_p^n$ not admitting non-trivial proper β -stable vector subspaces;
- (b) \mathbb{Q}_p^n for some $n \in \mathbb{N}$, together with β^{-1} for a contractive linear automorphism $\beta: \mathbb{Q}_p^n \rightarrow \mathbb{Q}_p^n$ not admitting non-trivial proper β -stable vector subspaces;
- (c) $C_p^{(-\mathbb{N})} \times C_p^{\mathbb{N}_0}$ with the right-shift;
- (d) $C_p^{(-\mathbb{N})} \times C_p^{\mathbb{N}_0}$ with the left-shift;
- (e) $C_p^{\mathbb{Z}}$ with the right-shift.

Proof. Let D_α be the divisible part and T_α be the torsion part of U_α , and define $D_{\alpha^{-1}}$ and $T_{\alpha^{-1}}$ analogously. If $D_\alpha \neq \{1\}$, then $D_\alpha = D_\alpha^*$ is an α -stable closed subgroup (see Theorem C) which is non-discrete (see 1.8 (a)) and thus $A = D_\alpha$. By 1.8 (a) and the hypotheses, D_α is a divisible simple contraction group and hence of the form described in (a) (see [9, Theorem A]). Likewise, A is of the form described in (b) whenever $D_{\alpha^{-1}} \neq \{1\}$.

Throughout the rest of the proof, assume that $D_\alpha = D_{\alpha^{-1}} = \{1\}$. Then $A = U_\alpha U_{\alpha^{-1}} = T_\alpha T_{\alpha^{-1}}$.

Since the nub U_0 of α is an α -stable closed subgroup of A , it either is all of A or discrete. Being also compact, it is finite in the latter case, and thus $\{1\}$ is an open α -stable (normal) subgroup of U_0 . Now [28, Corollary 4.4] (proper such do not exist) shows that $U_0 = \{1\}$.

Case $U_0 = \{1\}$: Then $T_\alpha = U_\alpha = U_\alpha U_0$ and $T_{\alpha^{-1}} = U_{\alpha^{-1}} = U_{\alpha^{-1}} U_0$ are closed α -stable subgroups of A (using 1.2 (a)). If $T_\alpha \neq \{1\}$, then T_α is non-discrete. Hence $T_\alpha = A$ by the hypotheses, and this is a simple contraction group which is a torsion group and hence of the form described in (c) (see [9, Theorem A]). Likewise, A is of the form described in (d) if $T_{\alpha^{-1}} \neq \{1\}$.

Case $A = U_0$: Then A is compact and is irreducible in the sense of [28, Definition 6.1] as all its proper α -stable closed (normal) subgroups are finite, and moreover A is infinite (as U_α or $U_{\alpha^{-1}}$ is non-trivial and hence non-discrete, being a contraction group). Hence, by [28, Proposition 6.3], (A, α)

is isomorphic to the right-shift of $F^{\mathbb{Z}}$ for a finite simple group F . Since A is abelian, $F \cong C_p$ for some p and thus A is of the form described in (e). \square

Remark 6.3 Let G be a totally disconnected, locally compact group and $\alpha: G \rightarrow G$ be an expansive automorphism. If G is abelian, then the map

$$\pi: U_\alpha^* \times U_{\alpha^{-1}}^* \rightarrow G, \quad (x, y) \mapsto xy$$

is a continuous, open homomorphism with discrete kernel. For non-abelian G , the map still has open image (see Lemma 1.1 (d)), is a local homeomorphism, and equivariant with respect to the natural left and right actions of U_α and $U_{\alpha^{-1}}$, respectively.

[To see this, let $V \subseteq G$ be a compact open subgroup such that $V_+ \cap V_- = \{1\}$ and $V = V_+ V_-$ (see Lemma 1.1 (c)). Then V_- and V_+ are open subgroups of U_α^* and $U_{\alpha^{-1}}^*$, respectively (see proof of Lemma 2.3). Then $\pi(V_+ \times V_-) = V$ is open in G and $\pi|_{V_+ \times V_-}$ is injective, as $vw = v'w'$ for $v, v' \in V_+$, $w, w' \in V_-$ implies $v^{-1}v' = w(w')^{-1} \in V_+ \cap V_- = \{1\}$ and thus $v = v'$ and $w = w'$. Since $V_+ \times V_-$ is compact, π restricts to a homeomorphism $V_+ \times V_- \rightarrow V$. Since $\pi(gv, wh) = g\pi(v, w)h$ for $g \in U_\alpha$, $h \in U_{\alpha^{-1}}$ and $(v, w) \in V_+ \times V_-$, also $\pi|_{gV_+ \times V_- h}$ is a homeomorphism onto an open set.]

Remark 6.4 It can happen that U_α is closed for an expansive automorphism α of a totally disconnected, locally compact group G , but $U_{\bar{\alpha}}$ is not closed for the induced automorphism $\bar{\alpha}$ on G/N for some α -stable closed normal subgroup $N \subseteq G$. The following example also illustrates Remark 6.3.

Given a non-trivial finite abelian group $(F, +)$, consider the restricted products $H_1 := F^{(-\mathbb{N})} \times F^{\mathbb{N}_0}$ and $H_2 := F^{-\mathbb{N}} \times F^{(\mathbb{N}_0)}$, with $V_1 := F^{\mathbb{N}_0}$ and $V_2 := F^{-\mathbb{N}}$, respectively, as compact open subgroups. Let α be the right-shift on $G := H_1 \times H_2$ (i.e., on both H_1 and H_2). Then α is an automorphism and it is expansive as $\bigcap_{n \in \mathbb{Z}} \alpha^n(V_1 \times V_2) = \{0\}$. Moreover, $U_\alpha = H_1$ and $U_{\alpha^{-1}} = H_2$ are closed. Also, let $\bar{\alpha}$ be the right-shift on $F^{\mathbb{Z}}$. Then

$$q: G \rightarrow F^{\mathbb{Z}}, \quad (f, g) \mapsto f + g$$

is a continuous surjective homomorphism. Restricted to the compact open subgroup $V_1 \times V_2$, the map q is an isomorphism of topological groups. Hence q is open, has discrete kernel, and is a quotient morphism. Finally, $U_{\bar{\alpha}} = F^{(-\mathbb{N})} \times F^{\mathbb{N}_0}$ is a dense proper subgroup in $F^{\mathbb{Z}}$. Hence $U_{\bar{\alpha}}$ is not closed in $F^{\mathbb{Z}} \cong G/\ker(q)$.

Another property can be observed.

Proposition 6.5 *Let G be a totally disconnected, locally compact group that is abelian, and $\alpha: G \rightarrow G$ be an expansive automorphism. Then the torsion subgroup $\text{tor}(G)$ is closed in G .*

Proof. Since $V := U_\alpha U_{\alpha^{-1}}$ is an open subgroup of G , we need only show that $V \cap \text{tor}(G) = \text{tor}(V)$ is closed. After replacing G with its α -stable subgroup V , we may therefore assume that $G = U_\alpha U_{\alpha^{-1}}$. Since D_α and $D_{\alpha^{-1}}$ are torsion-free (see 1.9 (c)) and $D_\alpha \cap D_{\alpha^{-1}} = \{1\}$ by Corollary 5.1, we deduce that $D_\alpha D_{\alpha^{-1}}$ is isomorphic to $D_\alpha \times D_{\alpha^{-1}}$ as an abstract group and hence torsion-free. Hence $D_\alpha D_{\alpha^{-1}} \cap T_\alpha T_{\alpha^{-1}} = \{1\}$. Combining this with $G = U_\alpha U_{\alpha^{-1}} = D_\alpha D_{\alpha^{-1}} T_\alpha T_{\alpha^{-1}}$, we see that

$$G = (D_\alpha D_{\alpha^{-1}}) \times (T_\alpha T_{\alpha^{-1}}) = D_\alpha \times D_{\alpha^{-1}} \times T_\alpha T_{\alpha^{-1}}$$

internally as an abstract group. Now $T_\alpha T_{\alpha^{-1}} \subseteq \text{tor}(G) \subseteq \overline{\text{tor}(G)}$. If $T_\alpha T_{\alpha^{-1}}$ was a proper subset of $\text{tor}(G)$, then the torsion group $\text{tor}(G)/T_\alpha T_{\alpha^{-1}}$ would be isomorphic to a non-trivial subgroup of $G/T_\alpha T_{\alpha^{-1}} \cong D_\alpha D_{\alpha^{-1}}$, which is torsion-free. The contradiction shows that $T_\alpha T_{\alpha^{-1}} = \text{tor}(G) = \overline{\text{tor}(G)}$. \square

7 Proof of Theorem D

Since $U_0 \subseteq \overline{U_\alpha}$, we may replace G with $\overline{U_\alpha}$ without changing the nub (see 1.7), nor $\overline{T_\alpha}$, nor D_α . We may therefore assume that U_α is dense in G . Since U_0 normalizes U_α (see 1.2 (f)), U_α is a normal subgroup of $G = U_\alpha U_0$ (exploiting 1.2 (a)). Hence also the characteristic subgroups D_α and T_α of U_α are normal in G . Therefore also $\overline{T_\alpha}$ is normal in G . Since $\overline{T_\alpha}$ is a torsion group (see 1.9 (c) and 1.5) and D_α torsion-free (see 1.9 (c)), we see that $D_\alpha \cap \overline{T_\alpha} = \{1\}$. Moreover, using that $T_\alpha U_0 = \overline{T_\alpha}$ by Lemma 2.9 (d), we obtain $G = U_\alpha U_0 = D_\alpha T_\alpha U_0 = D_\alpha \overline{T_\alpha}$. Hence $G = D_\alpha \times \overline{T_\alpha}$ as an abstract group. In particular, D_α centralizes $\overline{T_\alpha}$. Thus D_α also centralizes $U_0 \subseteq \overline{T_\alpha}$. Since $\overline{T_\alpha}$ is σ -compact by Lemma 2.10, also $D_\alpha \times \overline{T_\alpha}$ is a σ -compact locally compact group. Because also G is locally compact and the product map $\pi: D_\alpha \times \overline{T_\alpha} \rightarrow G$, $(x, y) \mapsto xy$ is a continuous isomorphism of abstract groups, we deduce with [10, 5.29] that π is an isomorphism of topological groups. \square

Remark 7.1 We mention that the nub U_0 of an expansive automorphism $\alpha: G \rightarrow G$ need not have an open normalizer in G . To see this, let F be a finite group which is a semidirect product $F = N \rtimes H$ of a normal subgroup N and a subgroup H which is not normal in F (e.g., F might be the dihedral group $C_3 \rtimes C_2$). Let G be the group of all $(n_k, h_k)_{k \in \mathbb{Z}} \in F^{\mathbb{Z}}$ such that $(n_k)_{k \in \mathbb{Z}} \in N^{(-\mathbb{N})} \times N^{\mathbb{N}_0} =: M$. Thus $G = M \rtimes H^{\mathbb{Z}}$ as an abstract group. Endow G with the topology making it the direct product topological space of the restricted product M and the compact group $H^{\mathbb{Z}}$. Then G is a topological group, being the ascending union of the open subgroups

$$H^{\{k \in \mathbb{Z}: k < -m\}} \times F^{\{k \in \mathbb{Z}: k \geq -m\}}$$

for $m \in \mathbb{N}$, which are topological groups. The right-shift α is an automorphism of G . We have $U_\alpha = M \rtimes (H^{(-\mathbb{N})} \times H^{\mathbb{N}_0})$ and $U_{\alpha^{-1}} = H^{-\mathbb{N}} \times H^{\mathbb{N}_0}$. Thus $\overline{U_\alpha} = G$, $\overline{U_{\alpha^{-1}}} = H^{\mathbb{Z}}$ and $U_0 = \overline{U_\alpha} \cap \overline{U_{\alpha^{-1}}} = H^{\mathbb{Z}}$ (using 1.2 (b)). Since H is not normal in F , we see that $U_0 = H^{\mathbb{Z}}$ is not normal in G . If the normalizer $N_G(U_0)$ was open in G , then (being α -stable), it would contain the dense subgroup U_α of G and hence coincide with G (a contradiction). Thus $N_G(U_0)$ is not open.

8 Example: p -adic Lie groups

Let \mathbb{K} be a local field and $|\cdot|$ be an absolute value on \mathbb{K} defining its topology (see [25]). We pick an algebraic closure $\overline{\mathbb{K}}$ containing \mathbb{K} and use the same symbol, $|\cdot|$, for the unique extension of the absolute value on \mathbb{K} to an absolute value on $\overline{\mathbb{K}}$ (see [18, Theorem 16.1]). If E is a finite-dimensional \mathbb{K} -vector space and $\beta: E \rightarrow E$ a \mathbb{K} -linear automorphism, we write $R(\beta)$ for the set of all absolute values $|\lambda|$ of zeros λ of the characteristic polynomial of β in $\overline{\mathbb{K}}$. We let $\tilde{E}_\lambda \subseteq E \otimes_{\mathbb{K}} \overline{\mathbb{K}}$ be the generalized eigenspace of $\beta \otimes_{\mathbb{K}} \text{id}_{\overline{\mathbb{K}}}$ for the eigenvalue λ . For $\rho \in R(\beta)$, we let

$$E_\rho := \left(\bigoplus_{|\lambda|=\rho} \tilde{E}_\lambda \right) \cap E.$$

Then $E = \bigoplus_{\rho \in R(\beta)} E_\rho$ (see [15, Chapter II, §1]) and we recall that

$$E = U_\beta \oplus M_\beta \oplus U_{\beta^{-1}}$$

with

$$M_\beta = E_1, \quad U_\beta = \bigoplus_{\rho < 1} E_\rho \quad \text{and} \quad U_{\beta^{-1}} = \bigoplus_{\rho > 1} E_\rho \quad (2)$$

(cf. Lemma 9.5 in the preprint version of [7]).

If G is a Lie group over \mathbb{K} , then its tangent space $L(G) := T_1(G)$ at the identity element carries a natural Lie algebra structure, and $L(\alpha): L(G) \rightarrow L(H)$ is a Lie algebra homomorphism for each \mathbb{K} -analytic homomorphism $\alpha: G \rightarrow H$ between \mathbb{K} -analytic Lie groups. We abbreviate $\text{Ad}(g) := L(I_g)$, where $I_g: G \rightarrow G, x \mapsto gxg^{-1}$ for $g \in G$ (cf. [21] for further information).

Proposition 8.1 *Let α be an analytic automorphism of a Lie group G over a local field. Consider the following conditions:*

- (a) α is expansive;
- (b) $\beta := L(\alpha): L(G) \rightarrow L(G)$ is expansive;
- (c) $1 \notin R(\beta)$;
- (d) $L(G)$ is a nilpotent Lie algebra.

Then (a) \Rightarrow (b), (b) \Leftrightarrow (c) and (c) \Rightarrow (d). If U_α is closed (e.g., if G is a p -adic Lie group), then (a), (b) and (c) are equivalent.

Proof. (a) \Rightarrow (b): By contraposition. If (b) is false, then β is not expansive. To deduce that α is not expansive, let $V \subseteq G$ be an identity neighbourhood. Since U_β is a vector subspace of $L(G)$ by (2) and hence closed, using Proposition 1.3 (b) we deduce that M_β is not discrete and hence a non-trivial vector subspace (in view of (2)). But then G contains a so-called centre manifold W around the fixed point 1 of α , which can be chosen as a submanifold of G contained in V that is stable under α and satisfies $T_1(W) = M_\alpha$ (whence $W \neq \{1\}$); see Proposition 13.3. (a) and Theorem 1.10 (c) in the preprint version of [7]. Then $\{1\} \neq W \subseteq \bigcap_{n \in \mathbb{Z}} \alpha^n(V)$, and thus α is not expansive.

(b) \Leftrightarrow (c): Since U_β is closed, β is expansive if and only if $M_\beta = L(G)_1$ is discrete. Since $L(G)_1$ is a vector space, the latter holds if and only if $L(G)_1 = \{0\}$, i.e., $1 \notin R(\beta)$.

(c) \Rightarrow (d). If (c) holds, then β is a Lie algebra automorphism of $L(G)$ and $|\lambda| \neq 1$ for all eigenvalues λ of $\beta \otimes_{\mathbb{K}} \text{id}_{\overline{\mathbb{K}}}$ in $\overline{\mathbb{K}}$, entailing that none of the λ

is a root of unity. Hence $L(G)$ is nilpotent (see Exercise 21 (b) among the exercises for Part I of [3], §4).

To prove the final assertion, we recall that U_α and $U_{\alpha^{-1}}$ are immersed Lie subgroups of G with Lie algebras U_β and $U_{\beta^{-1}}$, respectively (see Proposition 13.2 in the preprint version of [7]). If $1 \notin R(\beta)$, then $L(G) = U_\beta \oplus U_{\beta^{-1}}$, entailing that the product map $U_\alpha \times U_{\alpha^{-1}} \rightarrow G$, $(x, y) \mapsto xy$ has invertible differential at $(1, 1)$ (the addition map $U_\beta \times U_{\beta^{-1}} \rightarrow L(G) = U_\beta \oplus U_{\beta^{-1}}$). Thus, by the inverse function theorem [21], $U_\alpha U_{\alpha^{-1}}$ is an identity neighbourhood in G and hence open. If U_α is closed (which holds in the p -adic case by [24, Theorem 3.5 (ii)]), Proposition 1.3 (b) now shows that α is expansive. \square

8.2 For each continuous homomorphism $\theta: \mathbb{Q}_p \rightarrow \mathrm{GL}_n(\mathbb{Q}_p)$, there exists a nilpotent $n \times n$ -matrix $x \in \mathbb{Q}_p^{n \times n}$ such that $\theta(t) = \exp(tx)$ for all $t \in \mathbb{Q}_p$, using the matrix exponential function [17, Theorem 1.1]. Thus $\theta'(0) = x$ uniquely determines θ , and so does $\theta|_W$ for any 0-neighbourhood $W \subseteq \mathbb{Q}_p$.

Lemma 8.3 *Let α be a contractive automorphism of a p -adic Lie group G . Then the following holds:*

- (a) *For each $g \in G$, there is a unique continuous homomorphism $\theta_g: \mathbb{Q}_p \rightarrow G$ such that $\theta_g(1) = g$. Moreover, $\{\theta'_g(0) : g \in G\} = L(G)$.*
- (b) *If $\mathfrak{h} \subseteq L(G)$ is an $L(\alpha)$ -stable Lie subalgebra, then there exists an α -stable Lie subgroup H of G with $L(H) = \mathfrak{h}$.*

Proof. (a) Let $*$: $L(G) \times L(G) \rightarrow L(G)$ be the Campbell-Hausdorff multiplication on the nilpotent Lie algebra $L(G)$. Because (G, α) and $((L(G), *), L(\alpha))$ are locally isomorphic contraction groups, they are isomorphic (see [24, Proposition 2.2]). The nilpotent group $(L(G), *)$ inherits unique divisibility from $(L(G), +)$, since ng (in the vector space $L(G)$) coincides with g^n (in $(L(G), *)$). It is clear from this that $\theta_g(t) = tg$ is the unique continuous homomorphism $\mathbb{Q}_p \rightarrow (L(G), *)$ with $\theta_g(1) = g$. It satisfies $g = \theta'_g(0)$.

(b) We may work with the isomorphic group $(L(G), *)$ instead of G . Now $H := \mathfrak{h}$ is an $L(\alpha)$ -stable Lie subgroup of $(L(G), *)$ with Lie algebra \mathfrak{h} . \square

Lemma 8.4 *Let G be a linear p -adic Lie group. Assume that G is generated by $\bigcup_{\theta \in \Theta} \theta(\mathbb{Q}_p)$ for a set Θ of continuous homomorphisms $\theta: \mathbb{Q}_p \rightarrow G$, and $L(G)$ is generated by $\{\theta'(0): \theta \in \Theta\}$ as a Lie algebra. Then the centre of G coincides with the kernel of $\text{Ad}: G \rightarrow \text{Aut}(L(G))$.*

Proof. Let $g \in G$. For each $\theta \in \Theta$, the map $I_g \circ \theta: \mathbb{Q}_p \rightarrow G$, $t \mapsto g\theta(t)g^{-1}$ is a continuous homomorphism such that $(I_g \circ \theta)'(0) = \text{Ad}(g)\theta'(0)$. Thus, by 8.2, $I_g \circ \theta = \theta$ if and only if $\text{Ad}(g)\theta'(0) = \theta'(0)$. Since $\bigcup_{\theta \in \Theta} \theta(\mathbb{Q}_p)$ generates G , we see that $g \in Z(G)$ if and only if $\text{Ad}(g)\theta'(0) = \theta'(0)$ for all $\theta \in \Theta$. The latter is equivalent to $\text{Ad}(g)(x) = x$ for all $x \in L(G)$, because $\{x \in L(G): \text{Ad}(g)(x) = x\}$ is a Lie subalgebra of $L(G)$ and $L(G)$ is generated by $\theta'(0)$ for $\theta \in \Theta$ by hypothesis. \square

Proof of Theorem E. After replacing G with an open subgroup, we may assume that G is generated by $U_\alpha \cup U_{\alpha^{-1}}$ (see Lemma 1.1 (d)). We prove that G is nilpotent in this case, by induction on the dimension $\dim(G)$ of G as a p -adic manifold. If $\dim(G) = 0$, then G is discrete, whence $U_\alpha = U_{\alpha^{-1}} = \{1\}$ and $G = \langle U_\alpha \cup U_{\alpha^{-1}} \rangle = \{1\}$ is nilpotent.

Now assume that $\dim(G) > 0$. After replacing G with an isomorphic group, we may assume that G is a subgroup of $\text{GL}_n(\mathbb{Q}_p)$ for some $n \in \mathbb{N}$, and that the inclusion map $G \rightarrow \text{GL}_n(\mathbb{Q}_p)$ is continuous (but not necessarily a homeomorphism onto its image). Then $L(G)$ is a non-zero nilpotent Lie algebra (see (a) \Rightarrow (d) in Proposition 8.1) and so it has centre $Z(L(G)) \neq \{0\}$. The centre is $L(\alpha)$ -stable, and the restriction β of $L(\alpha)$ to the centre is expansive (like $L(\alpha)$). Hence $Z(L(G)) = U_\beta \oplus U_{\beta^{-1}}$. After replacing α with α^{-1} if necessary, we may assume that $U_{\beta^{-1}} \neq \{0\}$. According to Lemma 8.3 (b), there is an α -stable Lie subgroup $H \subseteq U_{\alpha^{-1}}$ with $L(H) = U_\beta$. We claim that H is in the centre $Z(G)$ of G . If this is true, then $Z(G)$ has positive dimension. Thus $G/Z(G)$ is a Lie group of dimension $\dim(G/Z(G)) < \dim(G)$, and it is a linear Lie group as it injects into $\text{Aut}(L(G))$, by Lemma 8.4.³ By induction, $G/Z(G)$ is nilpotent and hence so is G .

To prove the claim, let $h \in H$ and $\theta: \mathbb{Q}_p \rightarrow H \subseteq U_{\alpha^{-1}}$ be a continuous homomorphism with $\theta(1) = h$ (see Lemma 8.3 (a)). Then $x := \theta'(0) \in$

³Let Θ be the set of continuous homomorphisms from \mathbb{Q}_p to U_α or $U_{\alpha^{-1}}$. By $G = \langle U_\alpha \cup U_{\alpha^{-1}} \rangle$ and Lemma 8.3 (a), the first hypothesis of Lemma 8.4 is satisfied. Since $L(G) = L(U_\alpha) + L(U_{\alpha^{-1}})$ and $L(U_\alpha) \cup L(U_{\alpha^{-1}}) = \{\theta'(0): \theta \in \Theta\}$ by Lemma 8.3 (a) and “(a) \Rightarrow (c)” in Lemma 8.1, also the second hypothesis of Lemma 8.4 is satisfied.

$L(H) \subseteq Z(L(G))$, entailing that $\text{ad}(x) := [x, \bullet] = 0$. Now $\theta(t) = \exp(tx)$ for all $t \in \mathbb{Q}_p$, by 8.2. For $|t|$ small, $\text{Ad}(\theta(t)) = \text{Ad}(\exp(tx)) = e^{t \text{ad}(x)} = \text{id}_{L(G)}$, using Corollary 3 in [3, Chapter III, §4, no. 4]. Thus $\text{Ad} \circ \theta = \text{id}_{L(G)}$, by the uniqueness assertion of 8.2, applied to $\text{Ad} \circ \theta: \mathbb{Q}_p \rightarrow \text{Aut}(L(G))$. In particular, $\text{Ad}(h) = \text{Ad}(\theta(1)) = \text{id}_{L(G)}$ and thus $h \in Z(G)$, by Lemma 8.4. \square

8.5 If G is a totally disconnected, locally compact group which is a nilpotent group, let $\{1\} = Z_0 \triangleleft Z_1 \triangleleft \cdots \triangleleft Z_n = G$ be its ascending central series defined recursively via $Z_k := q_k^{-1}(Z(G/Z_{k-1}))$, where $q_k: G \rightarrow G/Z_{k-1}$ is the canonical quotient morphism. Let α be an expansive automorphism of G and α_k the induced automorphism of G_k/G_{k-1} .

Proposition 8.6 *If $Z_k/Z_{k-1} = U_{\alpha_k}U_{\alpha_k^{-1}}$ for all $k \in \{1, \dots, n\}$ in the situation of 8.5, then $G = U_\alpha U_{\alpha^{-1}}$. In particular, $U_\alpha U_{\alpha^{-1}}$ is a subgroup of G .*

Proof. If $n = 0$, then $G = \{1\} = U_\alpha U_{\alpha^{-1}}$. If $n \geq 1$, let β be the expansive automorphism of $G/Z(G)$ induced by α , and $q: G \rightarrow G/Z(G)$ be the canonical quotient morphism. Then $Z_1 = Z(G) = (U_\alpha \cap Z(G))(U_{\alpha^{-1}} \cap Z(G))$ by the hypotheses and $G/Z(G) = U_\beta U_{\beta^{-1}}$ by induction. Since $q(U_\alpha U_{\alpha^{-1}}) = U_\beta U_{\beta^{-1}} = G/Z(G)$, we have $G = U_\alpha U_{\alpha^{-1}} Z(G) = U_\alpha U_{\alpha^{-1}} (U_\alpha \cap Z(G))(U_{\alpha^{-1}} \cap Z(G)) = U_\alpha (U_\alpha \cap Z(G)) U_{\alpha^{-1}} (U_{\alpha^{-1}} \cap Z(G)) = U_\alpha U_{\alpha^{-1}}$. \square

Note that we can easily achieve that $G/Z_{n-1} = U_{\alpha_n} U_{\alpha_n^{-1}}$ after replacing G with its open subgroup generated by $U_\alpha \cup U_{\alpha^{-1}}$. However, the hypotheses on Z_k/Z_{k-1} for $k < n$ cannot always be achieved by passing to an open subgroup (as the following example illustrates).

Remark 8.7 The following example shows that even for nilpotent p -adic Lie groups with an expansive automorphism α , the set $U_\alpha U_{\alpha^{-1}}$ may fail to be a subgroup.

Let $H = \mathbb{Q}_p^3$ be the 3-dimensional p -adic Heisenberg group whose binary operation is given by

$$(x_1, y_1, z_1)(x_2, y_2, z_2) = (x_1 + x_2, y_1 + y_2, z_1 + z_2 + x_1 y_2)$$

for all $(x_1, y_1, z_1), (x_2, y_2, z_2) \in H$. Let $N = \{(0, 0, z) \in H : |z| \leq 1\}$. Then N is a compact central subgroup of H . Identify $G = H/N$ with $\mathbb{Q}_p \times \mathbb{Q}_p \times (\mathbb{Q}_p/\mathbb{Z}_p)$ as a set. Define $\alpha: G \rightarrow G$ by

$$\alpha(x, y, z + \mathbb{Z}_p) = (px, p^{-1}y, z + \mathbb{Z}_p)$$

for all $(x, y, z + \mathbb{Z}_p) \in G$. Then α is a continuous automorphism of the p -adic Lie group G with $M_\alpha = \{(0, 0, z + \mathbb{Z}_p) : z \in \mathbb{Q}_p\}$, $U_\alpha = \{(x, 0, 0) : x \in \mathbb{Q}_p\}$ and $U_{\alpha^{-1}} = \{(0, y, 0) : y \in \mathbb{Q}_p\}$. Since M_α is discrete, α is an expansive automorphism. As $[U_\alpha, U_{\alpha^{-1}}] = \{(0, 0, z + \mathbb{Z}_p) : z \in \mathbb{Q}_p\}$ and $U_\alpha U_{\alpha^{-1}} = \{(x, y, xy + \mathbb{Z}_p) : x, y \in \mathbb{Q}_p\}$, we get that $U_\alpha U_{\alpha^{-1}}$ is not a subgroup.

Proposition 8.8 *Let G be a closed subgroup of $\mathrm{GL}_n(\mathbb{Q}_p)$ and α be an expansive automorphism of G . Then $U_\alpha U_{\alpha^{-1}}$ is an open (unipotent algebraic) subgroup of G .*

Proof. Replacing G by the group generated by U_α and $U_{\alpha^{-1}}$, we may assume by Theorem E that G is a closed nilpotent subgroup of $\mathrm{GL}_n(\mathbb{Q}_p)$. Let \mathbb{G} be the Zariski closure of G . Then \mathbb{G} is defined over \mathbb{Q}_p and \mathbb{G} is nilpotent (cf. Proposition 1.3 (b) and Corollary 1 in 2.4 of [2]). Since U_α and $U_{\alpha^{-1}}$ consists of one-parameter (unipotent) subgroups, \mathbb{G} is Zariski-connected. This implies that the set of unipotent elements form a subgroup \mathbb{G}_u , known as the unipotent radical (cf. Theorem 10.6 of [2]). Since U_α and $U_{\alpha^{-1}}$ consists of one-parameter (unipotent) subgroups, $U_\alpha, U_{\alpha^{-1}} \subseteq \mathbb{G}_u$. This implies that $\mathbb{G} = \mathbb{G}_u$, that is \mathbb{G} is an unipotent algebraic group, hence \mathbb{G} is defined over \mathbb{Q}_p (cf. 4.5 of [2] and the fact that \mathbb{Q}_p -closed and defined over \mathbb{Q}_p are same as characteristic of \mathbb{Q}_p is zero) and $G \subseteq \mathbb{G}(\mathbb{Q}_p)$.

For $i \geq 1$, let $D_i = [\mathbb{G}(\mathbb{Q}_p), D_{i-1}]$ with $D_0 = \mathbb{G}(\mathbb{Q}_p)$ and $G_i = \overline{[G, G_{i-1}]}$ with $G_0 = G$. Then D_{k+1} is trivial for some $k \geq 1$ as \mathbb{G} is unipotent and $G_i \subset D_i$. Thus, G_k is a closed α -stable subgroup of D_k which is a vector space. Let V be the maximal vector subspace of G_k . Then V is a closed α -stable central subgroup of G . The automorphism $\beta: G_k/V \rightarrow G_k/V$ defined by $\beta(x + V) = \alpha(x) + V$ for $x \in G_k$ is expansive. Since V is the maximal vector subspace of G_k and G_k is a closed subgroup of the p -adic vector space D_k , we get that G_k/V is a compact subgroup of the p -adic vector space D_k/V . Since the automorphism group of a compact p -adic analytic group is compact, compact p -adic analytic groups do not admit expansive automorphisms unless finite, hence $V = G_k$. This implies that $G_k = D_k$ and $G_k = V = (U_\alpha \cap V)(U_{\alpha^{-1}} \cap V)$. Since G/G_k is a closed subgroup of $\mathbb{G}(\mathbb{Q}_p)/D_k$ which is a linear (p -adic algebraic) group, the result follows by induction. \square

Remark 8.9 In the case of linear p -adic Lie groups, even if $U_\alpha U_{\alpha^{-1}}$ is an open subgroup for an expansive automorphism, the following example shows

that it is not possible to have either of U_α or $U_{\alpha^{-1}}$ to normalize the other.

Let H be the 3-dimensional p -adic Heisenberg group defined as in Remark 8.7. For $i = 1, 2$, define $\alpha_i: H \rightarrow H$ by

$$\alpha_1(x, y, z) = (px, p^{-2}y, p^{-1}z), \quad \alpha_2(x, y, z) = (p^2x, p^{-1}y, pz)$$

for $(x, y, z) \in H$. Let $G = H \times H$ and $\alpha = \alpha_1 \times \alpha_2$. Then

$$U_\alpha = \{(x, 0, 0) : x \in \mathbb{Q}_p\} \times \{(a, 0, c) : a, c \in \mathbb{Q}_p\}$$

and

$$U_{\alpha^{-1}} = \{(0, y, z) : y, z \in \mathbb{Q}_p\} \times \{(0, b, 0) \mid b \in \mathbb{Q}_p\}.$$

Thus $U_\alpha U_{\alpha^{-1}} = G$. Since $\{(x, 0, 0) : x \in \mathbb{Q}_p\}$ and $\{(0, y, 0) : y \in \mathbb{Q}_p\}$ are not normal subgroups of H , neither U_α or $U_{\alpha^{-1}}$ normalize the other.

9 Example: Baumslag-Solitar groups

Throughout this section, we fix primes $p \neq q$. We let

$$\text{BS}(p, q) := \langle a, t \mid ta^p t^{-1} = a^q \rangle$$

be the Baumslag-Solitar group. Then $\langle a \rangle \cap g \langle a \rangle g^{-1}$ has finite index in $\langle a \rangle$ for each $g \in \text{BS}(p, q)$, and $\bigcap_g g \langle a \rangle g^{-1} = \{1\}$, hence the Schlichting completion $G_{p,q}$ of $\text{BS}(p, q)$ can be formed, which is a certain totally disconnected, locally compact group in which $\text{BS}(p, q)$ is dense, and in which $K := \overline{\langle a \rangle}$ is a compact open subgroup (see [4], cf. [19] and [8]). We are interested in the inner automorphism

$$\alpha: G_{p,q} \rightarrow G_{p,q}, \quad x \mapsto txt^{-1}.$$

Proof of Theorem F. By [4, Proposition 8.1], K contains an open subgroup $V \cong \mathbb{Z}_p \times \mathbb{Z}_q$ and K/V is a cyclic group of order dividing $\gcd(p, q) = 1$. Thus $K = V \cong \mathbb{Z}_p \times \mathbb{Z}_q$. After multiplication with a unit, We may assume that the isomorphism takes a to $(1, 1)$.

Let $G = \mathbb{Z} \ltimes (\mathbb{Q}_p \times \mathbb{Q}_q)$ be the semidirect product of \mathbb{Z} and $\mathbb{Q}_p \times \mathbb{Q}_q$ given by $(n, u, v)(m, u', v') = (n+m, u + (q/p)^n u', v + (q/p)^n v')$ for all $n, m \in \mathbb{Z}$, $u, u' \in \mathbb{Q}_p$ and $v, v' \in \mathbb{Q}_q$. The isomorphism $K \cong \mathbb{Z}_p \times \mathbb{Z}_q$ gives a homomorphism from $\langle a \rangle$ to G . Since $(1, 0, 0)(0, p, p)(-1, 0, 0) = (0, q, q)$, sending $t \mapsto (1, 0, 0)$ yields a group homomorphism $\phi: \text{BS}(p, q) \rightarrow G$. Since $\phi|_{\langle a \rangle}$ is a continuous

homomorphism, ϕ extends to a continuous homomorphism of $G_{p,q}$ into G which would also be denoted by ϕ . Since $\phi|_K$ is an isomorphism, $\ker(\phi)$ is discrete. Moreover, as $\phi(G_{p,q})$ contains both $(1, 0, 0)$ and $\mathbb{Z}_p \times \mathbb{Z}_q$, ϕ is surjective.

Let β be the inner automorphism of G given by $(1, 0, 0)$. Then $\phi \circ \alpha = \beta \circ \phi$ and β is expansive. Since the kernel of ϕ is discrete, expansiveness of α follows from Theorem A. As the open subgroup $K \cong \mathbb{Z}_p \times \mathbb{Z}_q$ satisfies an ascending chain condition on closed subgroups (see, e.g., [6, Proposition 3.2]), U_α is closed by [24, Lemma 3.2].

In case $U_\alpha U_{\alpha^{-1}}$ is a group, we will now show that ϕ is an isomorphism which would lead to a contradiction as $G_{p,q}$ is not solvable but G is solvable.⁴ Suppose $N := U_\alpha U_{\alpha^{-1}}$ is a group. Then $\phi|_N$ is an isomorphism of N with $\mathbb{Q}_p \times \mathbb{Q}_q$ (using that $U_\alpha \cong \mathbb{Q}_q$ and $U_{\alpha^{-1}} \cong \mathbb{Q}_p$).⁵ Now the group generated by t and N is an open subgroup of G containing both t and a , hence $G_{p,q} = \langle t, N \rangle = \langle t \rangle N$ (as t normalizes N). This implies that ϕ is an isomorphism. \square

References

- [1] Baumgartner, U. and G. A. Willis, *Contraction groups and scales of automorphisms of totally disconnected locally compact groups*, Israel J. Math. **142** (2004), 221–248.
- [2] Borel, A., "Linear algebraic groups". Second edition. Graduate Texts in Mathematics, 126. Springer-Verlag, New York, 1991.
- [3] Bourbaki, N., "Lie Groups and Lie Algebras, Chapters 1–3," Springer-Verlag, Berlin, 1989.
- [4] Elder, M. and G. A. Willis, *Totally disconnected groups from Baumslag-Solitar groups*, preprint, [arXiv:1301.4775](https://arxiv.org/abs/1301.4775).
- [5] Glöckner, H., *Contraction groups for tidy automorphisms of totally disconnected groups*, Glasg. Math. J. **47** (2005), 329–333.
- [6] Glöckner, H., *Locally compact groups built up from p -adic Lie groups, for p in a given set of primes*, J. Group Theory **9** (2006), 427–454.

⁴Alternatively, $\text{BS}(p, q)$ would be a finitely generated linear group then and hence residually finite by [14]. But $\text{BS}(p, q)$ is not residually finite for primes $p \neq q$, see [16].

⁵In fact, $V = K$ is tidy for α with $V_- \cong \mathbb{Z}_q$, $V_+ \cong \mathbb{Z}_p$ and $V_0 = \{1\}$ (see [4]), whence $U_\alpha = V_{--} \cong \mathbb{Q}_q$ and $U_{\alpha^{-1}} = V_{++} \cong \mathbb{Q}_p$ (cf. Lemma 1.1 (b)).

- [7] Glöckner, H., *Invariant manifolds for analytic dynamical systems over ultrametric fields*, Expo. Math. **31** (2013), 116–150 (expanded preprint version: [arXiv:0808.2709v2](https://arxiv.org/abs/0808.2709v2)).
- [8] Glöckner, H. and G. A. Willis, *Topologization of Hecke pairs and Hecke C^* -algebras*, Topol. Proc. **26** (2001–2002), 565–591.
- [9] Glöckner, H. and G. A. Willis, *Classification of the simple factors appearing in composition series of totally disconnected contraction groups*, J. Reine Angew. Math. **643** (2010), 141–169.
- [10] Hewitt, E. and K. A. Ross, “Abstract Harmonic Analysis,” Vol. I, Springer, 2nd Edition, New York, 1979.
- [11] Jaworski, W., *On contraction groups of automorphisms of totally disconnected locally compact groups*, Israel J. Math. **172** (2009), 1–8.
- [12] Kitchens, B. P., *Expansive dynamics on zero-dimensional groups*, Ergodic Theory Dynam. Systems **7** (1987), 249–261.
- [13] Lam, P., *On expansive transformation groups*, Trans. Amer. Math. Soc. **150** (1970), 131–138.
- [14] Mal’cev, A. I., *On the faithful representation of infinite groups by matrices*. Transl. Amer. Math. Soc. (2), **45** (1965), 1–18.
- [15] Margulis, G. A., “Discrete Subgroups of Semisimple Lie Groups,” Springer, Berlin, 1991.
- [16] Meskin, S., *Nonresidually finite one-relator groups*, Trans. Amer. Math. Soc. **164** (1972), 105–114.
- [17] Ratner, M., *Raghunathan’s conjectures for Cartesian products of real and p -adic Lie groups*, Duke Math. J. **77** (1995), 275–382.
- [18] Schikhof, W. H., “Ultrametric Calculus,” Cambridge University Press, 1984.
- [19] Schlichting, G., *On the periodicity of group operations*, pp. 507–517 in: Cheng, Kai Nah and Leong, Yu Kiang (eds.), “Group Theory” (Singapore 1987), de Gruyter, Berlin, 1989.
- [20] Schmidt, K., “Dynamical systems of algebraic origin,” Progress in Mathematics, 128. Birkhäuser Verlag, Basel, 1995.
- [21] Serre, J.-P., “Lie Algebras and Lie Groups,” Springer, Berlin, 1992.
- [22] Siebert, E., *Contractive automorphisms on locally compact groups*, Math. Z. **191** (1986), 73–90.

- [23] Siebert, E., *Semistable convolution semigroups and the topology of contraction groups*, pp. 325–343 in: “Probability Measures on Groups, IX” (Oberwolfach, 1988), Lecture Notes in Math. **1379**, Springer, Berlin, 1989.
- [24] Wang, J. S. P., *The Mautner phenomenon for p -adic Lie groups*, Math. Z. **185** (1984), 403–412.
- [25] Weil, A., “Basic Number Theory,” Springer, Berlin, 2nd ed., 1973.
- [26] Willis, G. A., *The structure of totally disconnected, locally compact groups*, Math. Ann. **300** (1994), 341–363.
- [27] Willis, G. A., *Further properties of the scale function on a totally disconnected group*, J. Algebra **237** (2001), 142–164.
- [28] Willis, G. A., *The nub of an automorphism of a totally disconnected, locally compact group*, Ergodic Theory Dyn. Systems, in print (DOI: <http://dx.doi.org/10.1017/etds.2012.172>)

Helge Glöckner, Universität Paderborn, Institut für Mathematik,
Warburger Str. 100, 33098 Paderborn, Germany;

e-mail: glockner@math.upb.de

C. R. E. Raja, Stat-Math Unit, Indian Statistical Institute (ISI),
8th Mile Mysore Road, Bangalore 560 059, India;

e-mail: creraja@isibang.ac.in