



# Sums Involving Moments of Reciprocals of Binomial Coefficients

Hacène Belbachir and Mourad Rahmani  
University of Sciences and Technology Houari Boumediene  
Faculty of Mathematics  
P. O. Box 32  
El Alia  
Bab-Ezzouar 16111  
Algiers  
Algeria

[hacenebelbachir@gmail.com](mailto:hacenebelbachir@gmail.com)  
[mrahmani@usthb.dz](mailto:mrahmani@usthb.dz)

B. Sury  
Statistics & Mathematics Unit  
Indian Statistical Institute  
8th Mile Mysore Road  
Bangalore 560059  
India

[sury@isibang.ac.in](mailto:sury@isibang.ac.in)

## Abstract

We investigate sums of the form  $\sum_{0 \leq k \leq n} k^m \binom{n}{k}^{-1}$ . We establish a recurrence relation and compute its ordinary generating function. As application we give the asymptotic expansion. The results extend the earlier works by various authors. In the last section, we establish that  $\sum_{0 \leq k \leq n} \frac{k^m}{n^m} \binom{n}{k}^{-1}$  tends to 1 as  $n \rightarrow \infty$  and that  $\sum_{0 \leq k \leq n-m} k^m \binom{n}{k}^{-1}$  tends to  $m!$  as  $n \rightarrow \infty$ .

# 1 Introduction

For all nonnegative integers  $n, m$ , let

$$S_n^{(m)} := \sum_{k=0}^n k^m \binom{n}{k}^{-1}. \quad (1)$$

There are several papers in the literature dealing with sums involving inverses of binomial coefficients [2, 9, 11, 12, 15, 16, 17, 19]. The cases  $m = 0$  and  $m = 1$  were intensively studied.

In 1947, Staver [13], using the identity

$$\sum_{k=0}^n k \binom{n}{k}^{-1} = \sum_{k=0}^n (n-k) \binom{n}{k}^{-1},$$

obtained a relation between  $S_n^{(1)}$  and  $S_n^{(0)}$ :

$$S_n^{(1)} = \frac{n}{2} S_n^{(0)}, \quad (2)$$

and established a recurrence relation for  $S_n^{(0)}$

$$S_{n+1}^{(0)} = \frac{n+2}{2(n+1)} S_n^{(0)} + 1. \quad (3)$$

Staver also proved for the first time the well-known formula

$$S_n^{(0)} = \frac{n+1}{2^{n+1}} \sum_{k=1}^{n+1} \frac{2^k}{k}. \quad (4)$$

For applications of (4), see Nedemeyer and Smorodinsky [5], Mansour and West [7], and for a probabilistic application, see Letac [4, p. 14]. Finally, for the asymptotic expansion, see Comtet [1, p. 294] and Yang and Zhao [18].

In 1981, using induction and the relation

$$\binom{n}{k}^{-1} = \binom{n-1}{k-1}^{-1} - \frac{n-k}{n-k+1} \binom{n}{k-1}^{-1}$$

Rockett [9] proved (3) and (4).

In 1993, Sury [14] connected the inverse binomial coefficients to the beta function as follows

$$\binom{n}{k}^{-1} = (n+1) \int_0^1 x^k (1-x)^{n-k} dx$$

and proved the relation (4) (see also [16]). Some years later, Mansour [6], generalized the idea of Sury [14], and gave an approach based on calculus to obtain the generating function for related combinatorial identities.

**Theorem 1** (Mansour [6]). Let  $r, n \geq k$  be any nonnegative integer numbers, and let  $f(n, k)$  be given by

$$f(n, k) = \frac{(n+r)!}{n!} \int_{u_1}^{u_2} p^k(t) q^{n-k}(t) dt,$$

where  $p(t)$  and  $q(t)$  are two functions defined on  $[u_1, u_2]$ . Let  $\{a_n\}_{n \geq 0}$  and  $\{b_n\}_{n \geq 0}$  be any two sequences, and let  $A(x), B(x)$  be the corresponding ordinary generating functions. Then

$$\sum_{n=0}^{\infty} \left[ \sum_{k=0}^n f(n, k) a_k b_{n-k} \right] x^n = \frac{d^r}{dx^r} \left[ x^r \int_{u_1}^{u_2} A(xp(t)) B(xq(t)) dt \right]. \quad (5)$$

In particular,

$$\sum_{n \geq 0} S_n^{(0)} x^n = \frac{2}{(x-1)(x-2)} - \frac{2 \ln(1-x)}{(x-2)^2},$$

and

$$\sum_{n \geq 0} S_n^{(1)} x^n = -\frac{x(3x-4)}{(x-1)^2(x-2)^2} + \frac{2x \ln(1-x)}{(x-2)^3}.$$

We shall use the following well-known basic tools [3]:

- (i) The Stirling numbers of the second kind  $\left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\}$  (A008277), can be defined by the generating function

$$\prod_{j=1}^k \frac{x}{1-jx} = \sum_{n \geq k} \left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\} x^n.$$

The most basic recurrence relation is

$$\left\{ \begin{smallmatrix} n+1 \\ k \end{smallmatrix} \right\} = \left\{ \begin{smallmatrix} n \\ k-1 \end{smallmatrix} \right\} + k \left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\},$$

with  $\left\{ \begin{smallmatrix} n \\ 1 \end{smallmatrix} \right\} = \left\{ \begin{smallmatrix} n \\ n \end{smallmatrix} \right\} = 1$ . An important relation involving  $\left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\}$  is

$$x^n = \sum_{k=0}^n (-1)^{n+k} \left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\} x(x+1) \cdots (x+k-1). \quad (6)$$

- (ii) The Eulerian numbers  $\left\langle \begin{smallmatrix} n \\ k \end{smallmatrix} \right\rangle$  (A008292) are defined by

$$\left\langle \begin{smallmatrix} n \\ k \end{smallmatrix} \right\rangle = \sum_{i=0}^k (-1)^i (k-i)^n \binom{n+1}{i}, \quad n \geq k \geq 1.$$

Which also satisfy the recursive relation

$$\left\langle \begin{smallmatrix} n \\ k \end{smallmatrix} \right\rangle = k \left\langle \begin{smallmatrix} n-1 \\ k \end{smallmatrix} \right\rangle + (n-k+1) \left\langle \begin{smallmatrix} n-1 \\ k-1 \end{smallmatrix} \right\rangle,$$

with  $\left\langle \begin{smallmatrix} 1 \\ 1 \end{smallmatrix} \right\rangle = 1$ .

(iii) The Worpitzky numbers  $W_{n,k}$  ([A028246](#)), are defined by

$$W_{n,k} = \sum_{i=0}^k (-1)^{i+k} (i+1)^n \binom{k}{i}.$$

They can also be expressed through the Stirling numbers of the second kind as follows

$$W_{n,k} = k! \left\{ \begin{matrix} n+1 \\ k+1 \end{matrix} \right\}. \quad (7)$$

The Worpitzky numbers satisfy the recursive relation

$$W_{n,k} = (k+1)W_{n-1,k} + kW_{n-1,k-1} \quad (n \geq 1, k \geq 1). \quad (8)$$

Some simple properties related to these three remarkable sequences are

$$\sum_{k=0}^n \left\langle \begin{matrix} n \\ k \end{matrix} \right\rangle x^k = \sum_{k=0}^n (x-1)^{n-k} kW_{n-1,k-1}, \quad (9)$$

$$\sum_{k=0}^n \binom{n}{k} \left\{ \begin{matrix} k \\ t \end{matrix} \right\} = \left\{ \begin{matrix} n+1 \\ t+1 \end{matrix} \right\}, \quad (10)$$

and

$$\sum_{k=0}^n \left\langle \begin{matrix} n \\ k \end{matrix} \right\rangle \binom{k+1}{t} = W_{n,n-t}. \quad (11)$$

## 2 Some results on $S_n^{(2)}$

In this section, we give some results concerned  $S_n^{(2)}$ . Applying Theorem 1, for  $a_n = n^2$  and  $b_n = 1$ , we obtain, for  $|x| < 1$ ,

$$A(x) = \frac{x(x+1)}{(1-x)^3},$$

and

$$B(x) = \frac{1}{1-x},$$

from (5) we get the generating function for  $S_n^{(2)}$

$$\begin{aligned} \sum_{n \geq 0} S_n^{(2)} x^n &= \frac{d}{dx} \left[ x \int_0^1 \frac{xt(xt+1)}{(1-xt)^3(1-x+xt)} dt \right] \\ &= \frac{2x(2x^3 - 3x^2 - 3x + 5)}{(x-2)^3(x-1)^3} - 2(x^2 + 2x - 2) \frac{\ln(1-x)}{(x-2)^4}. \end{aligned} \quad (12)$$

In addition, we have a relation between  $S_n^{(2)}$  and  $S_n^{(0)}$  given by the following theorem.

**Theorem 2.** *If  $n$  is a nonnegative integer, then  $S_n^{(2)}$  satisfies the recursion relation*

$$S_{n+1}^{(2)} = \frac{(n-1)(n+2)^2}{2(n-2)(n+1)^2} S_n^{(2)} + \frac{(n+2)(n^2-2n-2)}{2(n-2)}, \quad (13)$$

and

$$S_n^{(2)} = \frac{1}{4}(n+1)(n-2)S_n^{(0)} + \frac{1}{2}(n+1)^2. \quad (14)$$

*Proof.* We use the WZ method [8]. Denote the summand in  $S_n^{(2)}$  by  $L(n, k) := k^2 \binom{n}{k}^{-1}$ , by the Zeilberger's Maple package EKHAD, we construct the function

$$G(n, k) = (n^2k^2 + n + 3nk - nk^2 + 2 + 2k - 2k^2)(n+1-k) \binom{n}{k}^{-1}$$

such that

$$(n-1)(n+2)^2 L(n, k) - 2(n-2)(n+1)^2 L(n+1, k) = G(n, k+1) - G(n, k).$$

By summing the above telescoping equation over  $k$  from 0 to  $n-1$ , we obtain the following recurrence relation

$$(n-1)(n+2)^2 \sum_{k=0}^{n-1} L(n, k) - 2(n-2)(n+1)^2 \sum_{k=0}^{n-1} L(n+1, k) = G(n, n) - G(n, 0),$$

that we can rewrite

$$(n-1)(n+2)^2 S_n^{(2)} - (n-1)(n+2)^2 n^2 - 2(n-2)(n+1)^2 S_{n+1}^{(2)} + 2(n-2)(n+1)^2 \left( \frac{n^2}{n+1} + (n+1)^2 \right) = n^3(n-1),$$

as desired. We prove the relation (14) by induction on  $n$ , the result clearly holds for  $n=0$ , we now show that the formula for  $n+1$  follows from (13) and induction hypothesis

$$S_{n+1}^{(2)} = \frac{(n-1)(n+2)^2}{2(n-2)(n+1)^2} \left( \frac{1}{4}(n+1)(n-2)S_n^{(0)} + \frac{1}{2}(n+1)^2 \right) + \frac{(n+2)(n^2-2n-2)}{2(n-2)}.$$

Applying (3), we obtain

$$S_{n+1}^{(2)} = \frac{1}{4}(n+2)(n-1)S_{n+1}^{(0)} + \frac{1}{2}(n+2)^2.$$

This completes the proof.  $\square$

Using the relation  $\sum_{k=0}^n k^3 \binom{n}{k}^{-1} = \sum_{k=0}^n (n-k)^3 \binom{n}{k}^{-1}$  and relations (2) and (14), we easily obtain the following identity between  $S_n^{(3)}$  and  $S_n^{(0)}$ .

**Corollary 3.** *For any nonnegative integers  $n$ , we have*

$$S_n^{(3)} = \frac{1}{8}n(n^2-3n-6)S_n^{(0)} + \frac{3}{4}n(n+1)^2.$$

### 3 Generalization

Before stating the main result of this section, we need a lemma.

**Lemma 4.**

$$S_{n+j}^{(0)} = (n+j+1) \left( \frac{1}{2^j (n+1)} S_n^{(0)} + \sum_{r=0}^{j-1} \frac{1}{2^r (n+j-r+1)} \right), \quad (15)$$

*Proof.* We proceed by induction on  $j$ . For  $j = 0$  the identity (15) holds. Now suppose that (15) holds for some  $j$  and replace  $n$  by  $n+1$  in (15), then using relation (3) the result follows.  $\square$

**Theorem 5.** For any nonnegative integers  $m$  and  $n$ , we have

$$S_n^{(m)} = \sum_{j=0}^m (-1)^{m+j} \binom{n+j}{n} W_{m,j} \left( S_{n+j}^{(0)} - \frac{1}{(n+j)!} \sum_{r=0}^{j-1} r! (n+j-r)! \right), \quad (16)$$

with the usual convention that the empty sum is 0.

*Proof.* We can write  $S_n^{(m)}$  as follows

$$\begin{aligned} S_n^{(m)} &= \sum_{k=0}^n ((k+1) - 1)^m \binom{n}{k}^{-1}, \\ &= \sum_{k=0}^n \binom{n}{k}^{-1} \sum_{i=0}^m (-1)^{m-i} \binom{m}{i} (k+1)^i, \end{aligned}$$

and with (6), we obtain

$$\begin{aligned} S_n^{(m)} &= \sum_{k=0}^n \binom{n}{k}^{-1} \sum_{i=0}^m (-1)^{m-i} \binom{m}{i} \sum_{j=0}^i (-1)^{i+j} \left\{ \begin{matrix} i \\ j \end{matrix} \right\} (k+1) \cdots (k+j), \\ &= \sum_{k=0}^n \sum_{i=0}^m \sum_{j=0}^i \frac{1}{n!} (-1)^{m+j} \binom{m}{i} \left\{ \begin{matrix} i \\ j \end{matrix} \right\} k! (k+1) \cdots (k+j) (n-k)!. \end{aligned}$$

After some rearrangement,

$$\begin{aligned} S_n^{(m)} &= \sum_{k=0}^n \sum_{i=0}^m \sum_{j=0}^i (-1)^{m+j} \binom{m}{i} \left\{ \begin{matrix} i \\ j \end{matrix} \right\} (n+1) \cdots (n+j) \binom{n+j}{k+j}^{-1}, \\ &= \sum_{i=0}^m \sum_{j=0}^i (-1)^{m+j} \binom{m}{i} \left\{ \begin{matrix} i \\ j \end{matrix} \right\} j! \frac{(n+j)!}{n! j!} \sum_{r=j}^{n+j} \binom{n+j}{r}^{-1}. \end{aligned}$$

Now, from (10) and (11), the result holds.  $\square$

Setting  $m = 4$  in (16), we have the following

**Corollary 6.** *If  $n$  is a nonnegative integer, then*

$$S_n^{(4)} = \frac{1}{16} (n+1) (n^3 - 7n^2 - 2n + 16) S_n^{(0)} + \frac{1}{8} (7n - 8) (n+1)^3.$$

**Theorem 7.** *For any nonnegative integers  $m$  and  $n$*

$$S_{n+1}^{(m)} = \delta_{0m} + \frac{1}{n+1} \sum_{j=0}^{m+1} \binom{m+1}{j} S_n^{(j)}, \quad (17)$$

where  $\delta_{ij}$  is the Kronecker symbol.

*Proof.* Recall that  $\binom{n+1}{k} = \frac{n+1}{k} \binom{n}{k-1}$ , we have

$$\begin{aligned} \sum_{k=0}^{n+1} \frac{k^m}{\binom{n+1}{k}} &= \delta_{0m} + \sum_{k=1}^{n+1} k^m \binom{n+1}{k}^{-1} \\ &= \delta_{0m} + \frac{1}{n+1} \sum_{k=1}^{n+1} k^{m+1} \binom{n}{k-1}^{-1} \\ &= \delta_{0m} + \frac{1}{n+1} \sum_{k=0}^n (k+1)^{m+1} \binom{n}{k}^{-1} \\ &= \delta_{0m} + \frac{1}{n+1} \sum_{j=0}^{m+1} \binom{m+1}{j} \sum_{k=0}^n k^j \binom{n}{k}^{-1}. \end{aligned}$$

This proves the theorem. □

Setting  $m = 1$  in (17) and using (14), we have the following

**Corollary 8.** *If  $n$  is a nonnegative integer, then*

$$S_{n+1}^{(1)} = \frac{n+2}{2n} S_n^{(1)} + \frac{1}{2} (n+1).$$

## 4 Ordinary generating function

We apply Theorem 1, for  $a_n = n^m$  ( $m \geq 1$ ),  $b_n = 1$ , and for  $|x| < 1$  we have

$$\begin{aligned} A(x) &= \frac{1}{(1-x)^{m+1}} \sum_{k=0}^m \left\langle m \middle| k \right\rangle x^{k+1} = \sum_{k=0}^m \frac{(-1)^{m+k}}{(1-x)^{k+1}} W_{m,k}, \\ B(x) &= \sum_{n \geq 0} x^n = \frac{1}{1-x}. \end{aligned}$$

From (5), we get

$$\sum_{n \geq 0} S_n^{(m)} x^n = \frac{d}{dx} \left[ x \int_0^1 \frac{\sum_{k=0}^m \langle m \rangle_k (xt)^{k+1}}{(1-xt)^{m+1} (1-x+xt)} dt \right]. \quad (18)$$

Making the substitution  $xt = y$  in the right-hand side of (18), we obtain

$$\sum_{n \geq 0} S_n^{(m)} x^n = \frac{d}{dx} \left[ \int_0^x \frac{\sum_{k=0}^m \langle m \rangle_k y^{k+1}}{(1-y)^{m+1} (1-x+y)} dy \right].$$

Since the degree of the denominator is at least one higher than that of the numerator, this fraction decomposes into partial fractions of the form

$$\frac{\sum_{k=0}^m \langle m \rangle_k y^{k+1}}{(1-y)^{m+1} (1-x+y)} = \frac{\alpha^{(m)}(x)}{1-x+y} + \sum_{s=0}^m \frac{\alpha_s^{(m)}(x)}{(1-y)^{m-s+1}}. \quad (19)$$

We note in passing that (19) is equivalent to

$$\begin{aligned} \sum_{k=0}^m \langle m \rangle_k y^{k+1} &= (1-y)^{m+1} \alpha^{(m)}(x) + (1-x+y) \sum_{s=0}^m (1-y)^s \alpha_s^{(m)}(x) \\ &= \sum_{k=0}^m (-1)^{m+k} y (1-y)^{m-k} W_{m-1, k-1}. \end{aligned} \quad (20)$$

For  $y = 1$  and using the fact that  $W_{p,p} = p!$  for  $p \geq 0$ , we immediately obtain the well-known identity

$$\sum_{k=0}^m \langle m \rangle_k = m!.$$

Next, if we set  $y = 0$  for  $|x| < 1$ , we obtain the following relation between  $\alpha^{(m)}(x)$  and  $\alpha_s^{(m)}(x)$

$$\sum_{s=0}^m \alpha_s^{(m)}(x) = \frac{\alpha^{(m)}(x)}{x-1}. \quad (21)$$

**Proposition 9.** *For  $m \geq 1$ , we have*

$$\begin{aligned} \alpha_s^{(m)}(x) &= \sum_{k=0}^m \sum_{i=0}^s \frac{(-1)^{i+s}}{(2-x)^{i+1}} \langle m \rangle_k \binom{k+1}{s-i} \\ &= \sum_{j=m-s}^m \frac{(-1)^{m+j}}{(2-x)^{s-m+1+j}} W_{m,j}. \end{aligned} \quad (22)$$



and

$$\begin{aligned}
\alpha^{(m)}(x) &= \frac{1}{(2-x)^{m+1}} \sum_{k=0}^m \langle m \rangle \binom{m}{k} (x-1)^{k+1} \\
&= \sum_{j=0}^m \frac{(-1)^{m+j}}{(2-x)^{j+1}} W_{m,j}, \\
&= \alpha_m^{(m)}(x).
\end{aligned} \tag{23}$$

*Proof.* We verify that (22) and (23) satisfy (20). Denote the right-hand side of (20) by  $R^{(m)}(y)$

$$R^{(m)}(y) = (1-y)^{m+1} \alpha^{(m)}(x) + (1-x+y) \sum_{s=0}^m (1-y)^s \alpha_s^{(m)}(x).$$

After some rearrangement, we get

$$\begin{aligned}
R^{(m)}(y) &= \frac{(1-y)^{m+1}}{(2-x)^{m+1}} \sum_{k=0}^m \langle m \rangle \binom{m}{k} (x-1)^{k+1} + \\
&\quad (1-x+y) \sum_{s=0}^m (1-y)^s \sum_{k=0}^m \sum_{i=0}^s \frac{(-1)^{i+s}}{(2-x)^{i+1}} \langle m \rangle \binom{k+1}{s-i} \\
&= \sum_{k=0}^m \langle m \rangle \binom{m}{k} \left[ \frac{(1-y)^{m+1}}{(2-x)^{m+1}} (x-1)^{k+1} + \right. \\
&\quad \left. \frac{1-x+y}{2-x} \sum_{s=0}^m \frac{(1-y)^s}{(2-x)^s} \sum_{j=0}^s \binom{k+1}{j} (-1)^j (2-x)^j \right],
\end{aligned}$$

using binomial formula, we obtain

$$\begin{aligned}
R^{(m)}(y) &= \sum_{k=0}^m \langle m \rangle \binom{m}{k} \left[ \frac{(1-y)^{m+1}}{(2-x)^{m+1}} \sum_{j=0}^{k+1} \binom{k+1}{j} (2-x)^j (-1)^j + \right. \\
&\quad \sum_{s=0}^m \frac{(1-y)^s}{(2-x)^s} \sum_{j=0}^s \binom{k+1}{j} (-1)^j (2-x)^j - \\
&\quad \left. \sum_{s=0}^m \frac{(1-y)^{s+1}}{(2-x)^{s+1}} \sum_{j=0}^s \binom{k+1}{j} (-1)^j (2-x)^j \right].
\end{aligned}$$

Now, for  $k \leq m$

$$\begin{aligned}
R^{(m)}(y) &= \sum_{k=0}^m \langle m \rangle \left[ \sum_{s=m+1}^{m+1} \frac{(1-y)^s}{(2-x)^s} \sum_{j=0}^{m+1} \binom{k+1}{j} (2-x)^j (-1)^j + \right. \\
&\quad \left. \sum_{s=0}^m \frac{(1-y)^s}{(2-x)^s} \sum_{j=0}^s \binom{k+1}{j} (-1)^j (2-x)^j - \right. \\
&\quad \left. \sum_{s=0}^m \frac{(1-y)^{s+1}}{(2-x)^{s+1}} \sum_{j=0}^s \binom{k+1}{j} (-1)^j (2-x)^j \right] \\
&= \sum_{k=0}^m \langle m \rangle \left[ \sum_{s=0}^{m+1} \frac{(1-y)^s}{(2-x)^s} \sum_{j=0}^s \binom{k+1}{j} (-1)^j (2-x)^j - \right. \\
&\quad \left. \sum_{s=1}^{m+1} \frac{(1-y)^s}{(2-x)^s} \sum_{j=0}^{s-1} \binom{k+1}{j} (-1)^j (2-x)^j \right] \\
&= \sum_{k=0}^m \langle m \rangle \left[ 1 + \sum_{s=1}^{m+1} \frac{(1-y)^s}{(2-x)^s} \left( \sum_{j=0}^s \binom{k+1}{j} (-1)^j (2-x)^j - \right. \right. \\
&\quad \left. \left. \sum_{j=0}^{s-1} \binom{k+1}{j} (-1)^j (2-x)^j \right) \right].
\end{aligned}$$

Finally,

$$R^{(m)}(y) = \sum_{k=0}^m \langle m \rangle \left( 1 + \sum_{s=1}^{k+1} \binom{k+1}{s} (y-1)^s \right) = \sum_{k=0}^m \langle m \rangle y^{k+1}.$$

Now, according to (11) and (7), we have

$$\begin{aligned}
\alpha_s^{(m)}(x) &= \sum_{i=0}^s \frac{(-1)^{i+s}}{(2-x)^{i+1}} \sum_{k=0}^m \langle m \rangle \binom{k+1}{s-i} \\
&= \sum_{i=0}^s \frac{(-1)^{i+s} (m-s+i)!}{(2-x)^{i+1}} \left\{ \begin{matrix} m+1 \\ m-s+i+1 \end{matrix} \right\} \\
&= \sum_{i=0}^s \frac{(-1)^{i+s}}{(2-x)^{i+1}} W_{m, m-s+i} \\
&= \sum_{j=m-s}^m \frac{(-1)^{m+j}}{(2-x)^{s-m+1+j}} W_{m, j},
\end{aligned}$$

on the other side, it follows from (9) that

$$\begin{aligned}
\alpha^{(m)}(x) &= \sum_{j=0}^{m-1} (-1)^{m+j+1} \frac{(j+1)(x-1)}{(2-x)^{j+2}} W_{m-1,j} \\
&= \sum_{j=0}^{m-1} \frac{(-1)^{m+j}}{(2-x)^{j+1}} (j+1) W_{m-1,j} - \sum_{j=0}^{m-1} \frac{(-1)^{m+j}}{(2-x)^{j+2}} (j+1) W_{m-1,j} \\
&= \sum_{j=0}^m \frac{(-1)^{m+j}}{(2-x)^{j+1}} (j+1) W_{m-1,j} + \sum_{j=0}^m \frac{(-1)^{m+j}}{(2-x)^{j+1}} j W_{m-1,j-1}.
\end{aligned}$$

Using (8), we get  $\alpha^{(m)}(x)$  as desired. This completes the proof.  $\square$

Now, Integrating the right-hand side of (19) over  $y$ , we obtain

$$\begin{aligned}
\int_0^x \frac{\sum_{k=0}^m \langle m \rangle_k y^{k+1}}{(1-y)^{m+1} (1-x+y)} dy = \\
-2\alpha^{(m)}(x) \ln(1-x) + \sum_{s=0}^{m-1} \frac{\alpha_s^{(m)}(x)}{s-m} (1 - (1-x)^{s-m}). \quad (24)
\end{aligned}$$

By differentiating (24) we get the ordinary generating function of  $S_n^{(m)}$

$$\begin{aligned}
\sum_{n \geq 0} S_n^{(m)} x^n &= -2 \frac{d}{dx} \alpha^{(m)}(x) \ln(1-x) + \frac{\alpha^{(m)}(x)}{1-x} + \\
&\sum_{s=0}^{m-1} \frac{d}{dx} \frac{\alpha_s^{(m)}(x)}{s-m} (1 - (1-x)^{s-m}) + \sum_{s=0}^m \alpha_s^{(m)}(x) (1-x)^{s-m-1} \quad (25)
\end{aligned}$$

with

$$\frac{d}{dx} \alpha_s^{(m)}(x) = \sum_{j=m-s}^m \frac{(s-m+j+1)(-1)^{m+j}}{(2-x)^{s-m+j+2}} W_{m,j}.$$

With Proposition 9, we can now rewrite (25) as follows

**Theorem 10.** *For any real number  $x$  such that  $|x| < 1$  and for all positive integers  $m \geq 1$ , we have*

$$\begin{aligned}
\sum_{n \geq 0} S_n^{(m)} x^n &= \left( \sum_{j=0}^m \frac{2(j+1)(-1)^{m+j+1}}{(2-x)^{j+2}} W_{m,j} \right) \ln(1-x) \\
&+ \sum_{0 \leq j \leq s \leq m-1} \frac{(-1)^j W_{m,m-j}}{(2-x)^{s-j+1}} \left( \frac{(s-j+1)(1-(1-x)^{s-m})}{(s-m)(2-x)} + (1-x)^{s-m-1} \right) \\
&+ \frac{1}{1-x} \sum_{j=0}^m \frac{2(-1)^{j+m}}{(2-x)^{j+1}} W_{m,j}.
\end{aligned}$$

## 5 Asymptotic expansion

Yang and Zhao [18] proved recently the asymptotic expansions for  $S_n^{(0)}$  and  $S_n^{(1)}$  of the following type:

$$S_n^{(0)} \sim 2 + \frac{2}{n-1} - \frac{1}{2^{n-1}}, n \rightarrow \infty, \quad (26)$$

and

$$S_n^{(1)} \sim \frac{n}{2} \left( 2 + \frac{2}{n-1} - \frac{1}{2^{n-1}} \right), n \rightarrow \infty.$$

However, there are finer asymptotics which we proceed to discuss now. We note that the following claim is consistent with Theorem 7.

$$S_n^{(m)} \sim n^m, \quad n \rightarrow \infty.$$

It is not difficult to observe that  $T_n^{(m)} := \sum_{k=0}^{n-m} k^m \binom{n}{k}^{-1}$  also converges as  $n \rightarrow \infty$  and, we prove alongside the above asymptotic formula that this limit is  $m!$ .

**Theorem 11.** *For  $m > 0$ , we have*

$$\lim_{n \rightarrow \infty} T_n^{(m)} = m!,$$

and

$$\lim_{n \rightarrow \infty} \frac{S_n^{(m)}}{n^m} = 1.$$

*Proof.* Let us denote the sum  $S_n^{(0)}$  by  $S_n$  for simplicity just in this proof. One has the recurrence relation  $S_n = \frac{n+1}{2n} S_{n-1} + 1$  (see [14], for instance). Using this, it can be shown that

$$S_n \rightarrow 2,$$

$$n(S_n - 2) \rightarrow 2,$$

$$n(n(S_n - 2) - 2) = n^2 S_n - 2n^2 - 2n \rightarrow 4,$$

and so on. More generally, there are constants  $a_0, a_1, a_2, \dots$  such that, recursively

$$u_n^{(k)} = n u_n^{(k-1)} - a_{k-1},$$

$$u_n^{(k)} \rightarrow 0 \text{ as } n \rightarrow \infty,$$

where  $u_n^{(0)} = S_n - 2$ .

In other words, for all  $k \geq 1$ ,

$$n^k S_n - (2n^k + a_0 n^{k-1} + \dots + a_{k-2} n) \rightarrow a_k,$$

as  $n \rightarrow \infty$ , for some  $a_k$ .

In fact, the sequence  $a_0, a_1, \dots$  is  $2, 4, 16, 88, 616, 5224, \dots$  is described by the generating function

$$\sum_{n \geq 0} a_n \frac{x^n}{n!} = \frac{2}{(2 - e^x)^2}.$$

The constants can also be defined as

$$a_k = \sum_{r=0}^k (r+2) W_{k,r} - (r+1) W_{k-1,r},$$

or recursively as

$$a_k = 4 + \sum_{r=1}^k \left( \binom{k+1}{r} + \binom{k}{r} \right) (-1)^r (2 - a_0 + a_1 - \dots \pm a_{r-1}) + (-1)^{k+1} (2 - a_0 + \dots \pm a_{k-1}).$$

This is seen from

$$\begin{aligned} 2n^{k+1} S_n &= n^k (2n S_n) \\ &= n^k ((n+1) S_{n-1} + 2n) \\ &= (n^{k+1} + n^k) S_{n-1} + 2n^{k+1} \\ &= \sum_{r=1}^k \left( \binom{k+1}{r} + \binom{k}{r} \right) (n-1)^r S_{n-1} + 2n^{k+1}. \end{aligned}$$

Write

$$X^m = c_0 + c_1(X+1) + c_2(X+1)(X+2) + \dots + c_m(X+1)(X+2)\dots(X+m), \quad (27)$$

where  $c_i$ 's depend on  $m$  ( $c_m = 1$ ). We have

$$\begin{aligned} T_n^{(m)} &= \sum_{k=0}^{n-m} (c_0 + c_1(k+1) + c_2(k+1)(k+2) + \dots + c_m(k+1)(k+2)\dots(k+m)) \binom{n}{k}^{-1} \\ &= c_0 \sum_{k=0}^{n-m} \binom{n}{k}^{-1} + c_1(n+1) \sum_{k=0}^{n-m} \binom{n+1}{k+1}^{-1} + \dots + c_m(n+1)\dots(n+m) \sum_{k=0}^{n-m} \binom{n+m}{k+m}^{-1} \\ &= c_0 \left( S_n - \sum_{k=n-m+1}^n \binom{n}{k}^{-1} \right) + c_1(n+1) \left( S_{n+1} - \sum_{k=n-m+2}^{n+1} \binom{n+1}{k}^{-1} - 1 \right) + \dots \\ &\quad \dots + c_m(n+1)\dots(n+m) \left( S_{n+m} - \sum_{k=n+1}^{n+m} \binom{n+m}{k}^{-1} - \sum_{k=0}^{m-1} \binom{n+m}{k}^{-1} \right). \end{aligned}$$

A computation using the above result

$$n^k S_n - (2n^k + a_0 n^{k-1} + \dots + a_{k-1} n + a_k) \rightarrow 0,$$

shows

$$\lim_{n \rightarrow \infty} T_n^{(m)} = 0!c_0 + 1!c_1 + 2!c_2 + \cdots + (m-1)!c_{m-1} + 2(m!)c_m.$$

This can be shown by induction on  $m$ . The proof is finished by observing that  $c_m = 1$  and

$$0!c_0 + 1!c_1 + 2!c_2 + \cdots + m!c_m = 0,$$

which follows by looking at the constant term of (27). This finishes the proof of the first assertion. For the second one, we proceed similarly to the above discussion. We have

$$\begin{aligned} S_n^{(m)} &= \sum_{k=0}^n (c_0 + c_1(k+1) + c_2(k+1)(k+2) + \cdots + (k+1)(k+2)\cdots(k+m)) \binom{n}{k}^{-1} \\ &= c_0 S_n + c_1(n+1)(S_{n+1} - 2) + c_2(n+1)(n+2) \left( S_{n+2} - 2 \sum_{k=0}^1 \binom{n+2}{k}^{-1} \right) + \cdots \\ &\quad \cdots + (n+1)\cdots(n+m) \left( S_{n+m} - 2 \sum_{k=0}^{m-1} \binom{n+m}{k}^{-1} \right). \end{aligned}$$

Thus, looking at the largest degree factor, we obtain

$$S_n^{(m)} \sim n^m.$$

□

A corollary of the above proof is that the sum

$$S_n^{(0)} = 2 + \frac{2}{n} + \frac{4}{n^2} + \frac{16}{n^3} + \frac{88}{n^4} + \frac{616}{n^5} + \frac{5224}{n^6} + \cdots,$$

as  $n \rightarrow \infty$ .

## 6 Acknowledgment

The authors are indebted to the referee for making constructive suggestions for improvement. The asymptotic result  $S_n^{(m)} \sim n^m$  was pointed out by the referee and we have revised the paper to replace the earlier asymptotics by it. The referee has a rather different approach to the results of this paper and we look forward to seeing his proof in print soon.

The first and the second author were partially supported by CMEP project 09 MDU 765.

## References

- [1] L. Comtet, *Advanced Combinatorics*, Reidel, 1974.
- [2] P. Juan, The sum of inverses of binomial coefficients revisited. *Fib. Quart.* **35** (1997), 342–345.

- [3] R. Graham, D. Knuth, and O. Patashnik, *Concrete Mathematics*, Addison-Wesley Publishing Company, 1994.
- [4] G. Letac, *Problèmes de probabilité*, Presses Universitaires de France, 1970.
- [5] F. Nedemeyer and Y. Smorodinsky, Resistances in the multidimensional cube. *Quantum* **7** (1) (Sept./Oct. 1996), 12–15, 63.
- [6] T. Mansour, Combinatorial identities and inverse binomial coefficients. *Adv. in Appl. Math.* **28** (2002), 196–202.
- [7] T. Mansour and J. West, Avoiding 2-letter signed patterns. *Semm. Lothar. Combin.* **49** (2002/04), Art. B49a.
- [8] M. Petkovsek, H. S. Wilf and D. Zeilberger, *A = B*, A. K. Peters, 1996.
- [9] M. A. Rockett, Sums of the inverses of binomial coefficients. *Fib. Quart.* **19** (1981), 433–437.
- [10] N. J. A. Sloane, *The On-Line Encyclopedia of Integer Sequences*. Published electronically at <http://www.research.att.com/~njas/sequences>, 2010.
- [11] A. Sofo, General properties involving reciprocals of binomial coefficients. *J. Integer Sequences* **9** (2004), [Article 06.4.5](#).
- [12] R. Sprugnoli, Sums of the reciprocals of central binomial coefficients. *Integers* **6** (2006), Paper A27.
- [13] T. B. Staver. Om summasjon av potenser av binomialkoeffisientene. *Norsk Mat. Tidssker* **29** (1947), 97–103.
- [14] B. Sury, Sum of the reciprocals of the binomial coefficients. *European J. Combin.* **14** (1993), 351–353.
- [15] B. Sury, Tianming Wang, and Feng-Zhen Zhao. Some identities involving reciprocals of binomial coefficients. *J. Integer Sequences* **7** (2004), [Article 04.2.8](#).
- [16] T. Trif, Combinatorial sums and series involving inverses of binomial coefficients. *Fib. Quart.* **38** (2000), 79–84.
- [17] J.-H. Yang and F.-Z. Zhao. Sums involving the inverses of binomial coefficients. *J. Integer Sequences* **9** (2006), [Article 06.4.2](#).
- [18] J.-H. Yang and F.-Z. Zhao. The asymptotic expansions of certain sums involving inverse of binomial coefficient. *International Mathematical Forum* **5** (2010) 761–768.
- [19] F.-Z. Zhao and T. Wang. Some results for sums of the inverses of binomial coefficients. *Integers* **5** (2005), Paper A22.

---

2000 *Mathematics Subject Classification*: Primary 11B65; Secondary 05A10, 05A16.

*Keywords*: Binomial coefficient, recurrence relation, generating function, asymptotic expansion.

---

(Concerned with sequences [A008277](#), [A008292](#), and [A028246](#).)

---

Received December 1 2010; revised versions received December 8 2010; May 28 2011. Published in *Journal of Integer Sequences*, June 10 2011.

---

Return to [Journal of Integer Sequences home page](#).