

INVARIANT SUBSPACES OF VOICULESCU'S CIRCULAR OPERATOR

K. DYKEMA AND U. HAAGERUP

1 Introduction

The invariant subspace problem relative to a von Neumann algebra $\mathcal{M} \subseteq B(\mathcal{H})$ asks whether every operator $T \in \mathcal{M}$ has a proper, nontrivial invariant subspace $\mathcal{H}_0 \subseteq \mathcal{H}$ such that the orthogonal projection p onto \mathcal{H}_0 is an element of \mathcal{M} ; equivalently, it asks whether there is a projection $p \in \mathcal{M}$, $p \notin \{0, 1\}$, such that $Tp = pTp$. Even when \mathcal{M} is a II_1 -factor, this invariant subspace problem remains open. In this paper we show that the circular operator and each circular free Poisson operator (defined below) has a continuous family of invariant subspaces relative to the von Neumann algebra it generates. These operators arise naturally in free probability theory (see the book [VDN]), and each generates the von Neumann algebra II_1 -factor $L(F_2)$ associated to the nonabelian free group on two generators.

Given a von Neumann algebra \mathcal{M} with normal faithful state ϕ , a circular operator is $y = (x_1 + ix_2)/\sqrt{2} \in \mathcal{M}$, where x_1 and x_2 are centered semicircular elements having the same second moments and that are free with respect to ϕ . For specificity, we will always take circular elements as having the normalization $\phi(y^*y) = 1$, which is equivalent to $\phi(x_i^2) = 1$. Voiculescu found [V2] a matrix model for a circular element, showing that if $X(n)$ is a random matrix whose entries are i.i.d. complex $(0, 1/n)$ -Gaussian random variables then $X(n)$ converges in $*$ -moments as $n \rightarrow \infty$ to a circular element, meaning that

$$\lim_{n \rightarrow \infty} \tau_n(X(n)^{\epsilon(1)} X(n)^{\epsilon(2)} \cdots X(n)^{\epsilon(k)}) = \phi(y^{\epsilon(1)} y^{\epsilon(2)} \cdots y^{\epsilon(k)})$$

for every $k \in \mathbf{N}$ and for every choice of $\epsilon(j)$ being “ $*$ ” or no symbol, where τ_n is the expectation of the normalized trace and where y is a circular

K.D. is partially supported by U.S. NSF grant DMS 0070558. U.H. is a member of MaPhySto, Centre for Mathematical Physics and Stochastics, which is funded by a grant from The Danish National Research Foundation.

element. Using the matrix model, Voiculescu showed that if $(y_{ij})_{1 \leq i, j \leq N}$ is a $*$ -free family of circular elements in a von Neumann algebra \mathcal{M} with respect to a normal faithful state ϕ , then the matrix $y = \frac{1}{\sqrt{N}}(y_{ij})_{1 \leq i, j \leq N} \in M_N(\mathcal{M})$ is circular with respect to the state ϕ_N given by $\phi_N((x_{ij})_{1 \leq i, j \leq N}) = \frac{1}{N} \sum_{i=1}^N \phi(x_{ii})$. Furthermore, he showed that the polar decomposition of a circular operator y is $y = ub$ where u is a Haar unitary (i.e. a unitary satisfying $\phi(u^k) = 0$ for every integer $k > 0$), where b is a quarter-circular element, (i.e. having moments $\phi(b^k) = \frac{1}{\pi} \int_0^2 t^k \sqrt{4-t^2}$) and where u and b are $*$ -free. These and results of a similar nature have been instrumental in applications of free probability to the study of the free group factors $L(F_n)$ and related factors; some of the first of these were [V1], [R1], [Dy1], [R2], [Dy2].

Voiculescu's matrix model for the circular element is the starting point for finding invariant subspaces. Combined with a result of Dyson, it leads to upper triangular matrix models for the circular operator, namely, a sequence $Y(n)$ of upper triangular random matrices whose $*$ -moments converge to those of a circular operator. In these models, the elements of $Y(n)$ that are above the diagonal are complex $(0, 1/n)$ -Gaussian random variables, and we show that a number of different choices are possible for the diagonal entries. From these matrix models we show that for any $N \geq 2$ a circular operator can be realized as an $N \times N$ upper triangular matrix

$$\frac{1}{\sqrt{N}} \begin{pmatrix} a_1 & b_{12} & b_{13} & \cdots & b_{1,N-1} & b_{1N} \\ 0 & a_2 & b_{23} & \cdots & b_{2,N-1} & b_{2N} \\ 0 & 0 & a_3 & \ddots & \vdots & b_{3N} \\ \vdots & \vdots & \ddots & \ddots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 & a_{N-1} & b_{N-1,N} \\ 0 & 0 & \cdots & 0 & 0 & a_N \end{pmatrix} \quad (1)$$

with entries in some W^* -noncommutative probability space, where the collection of $N(N+1)/2$ nonzero entries is $*$ -free, where the entries b_{ij} lying strictly above the diagonal are circular elements and where the entries a_j on the diagonal are *circular free Poisson elements* (a_j being circular free Poisson of parameter j). These latter are generalizations of the circular operator (in the family of R-diagonal elements introduced by Nica and Speicher [NS]) which are quite natural from the perspective of free probability theory.

DEFINITION 1.1. *Let (A, ψ) be a W^* -noncommutative probability space with ψ faithful and let $c \geq 1$. A circular free Poisson element of parameter*

c in (A, ψ) is an element of the form UH_c where $U, H_c \in A$, U is a Haar unitary, $H_c \geq 0$, the pair $\{U, H_c\}$ is $*$ -free and H_c^2 has moments equal to those of a free Poisson distribution¹ with parameter c . Thus, letting $a = (1 - \sqrt{c})^2$ and $b = (1 + \sqrt{c})^2$, the moments of H_c^2 are equal to those of the measure ν_c that is supported on $[a, b]$, is absolutely continuous with respect to Lebesgue measure and has density

$$\frac{d\nu_c}{d\lambda}(t) = \frac{\sqrt{(b-t)(t-a)}}{2\pi t} 1_{[a,b]}(t).$$

We hasten to point out that a circular element z with normalization $\psi(z^*z) = 1$ is nothing other than a circular free Poisson element of parameter $c = 1$.

The spectrum of a circular free Poisson element of parameter c has been found by Haagerup and Larsen [HL] to be the annulus centered at zero with radii $\sqrt{c-1}$ and \sqrt{c} , if $c > 1$, whereas the spectrum of the circular operator is the disk of radius 1. In the realization (1) of the circular operator, we have that the diagonal entry a_j is circular free Poisson of parameter j . Therefore, the spectrum of the diagonal entry a_j increases in modulus as j increases, and the spectra of a_j and a_k overlap only if $|j - k| \leq 1$. These properties of the realization (1) allow general techniques for upper triangular operators to be applied in order to find invariant subspaces of the circular operator y . It turns out that for every $0 < r < 1$ there is a unique projection $p \in \mathcal{M} = \{y\}''$ such that

- (i) $yp = pyp$
- (ii) $\sigma(yp) = \{z \in \mathbf{C} \mid |z| \leq r\}$
- (iii) $\sigma((1-p)y) = \{z \in \mathbf{C} \mid r \leq |z| \leq 1\}$

where in (ii) (respectively (iii)) the spectrum is computed relative to the algebra $p\mathcal{M}p$, (respectively $(1-p)\mathcal{M}(1-p)$).

In fact, the techniques outlined above can be employed with very little extra effort to find invariant subspaces for every circular free Poisson operator, and the proof is presented in this generality throughout.

In §2, a theory is developed proving the existence of invariant subspaces, relative to the generated von Neumann algebras, of upper triangular operators, the spectra of whose diagonal entries satisfy certain conditions. In §3, we consider upper triangular random matrices whose entries strictly above

¹We must point out that the formula found in [VDN, p.35] for the free Poisson distribution has some errors. The formula for \mathcal{R}_μ found there is correct, but the formulae for $G_\mu(z)$ and for the density are incorrect.

the diagonal are i.i.d. complex Gaussian random variables which are independent of the diagonal entries. The general theme of the results in §3 is that the diagonal entries may be changed in certain ways but that as matrix size tends to infinity, the limit $*$ -moments remain the same. In §4, we generalize asymptotic freeness results which Voiculescu originally proved [V2] for Gaussian random matrices and constant diagonal matrices; we allow the diagonal matrices to be random, subject to certain conditions. In §5, the random matrix results of the previous two sections together with results of Dyson and others are used to find various upper triangular matrix models for circular free Poisson elements, and these are in turn used to find an upper triangular realization of the same, as in (1). In §6, this upper triangular realization of the circular free Poisson element is fed into the machinery of §2 to find invariant subspaces.

2 An Invariant Subspace for an Upper Triangular Operator

Suppose \mathcal{H} is a Hilbert space and $T : \mathcal{H} \rightarrow \mathcal{H}$ is a bounded operator. In this section we are concerned with invariant subspaces \mathcal{H}_0 for T such that the projection from \mathcal{H} onto \mathcal{H}_0 lies in the von Neumann algebra generated by T . It is easy to see (Lemma 2.1) that for every $r \geq 0$ the set

$$\mathcal{H}_r(T) = \overline{\{\xi \in \mathcal{H} \mid \limsup_{k \rightarrow \infty} \|T^k \xi\|^{1/k} \leq r\}} \quad (2)$$

is such an invariant subspace \mathcal{H}_0 . The question is whether, for any given operator, these subspaces can be other than $\{0\}$ or \mathcal{H} . We will show (Proposition 2.2) that the answer is yes if T can be written as an upper triangular operator,

$$T = \begin{pmatrix} * & * & * \\ 0 & * & * \\ 0 & 0 & * \end{pmatrix}$$

with respect to a decomposition $\mathcal{H} = \mathcal{H}_1 \oplus \mathcal{H}_2 \oplus \mathcal{H}_3$ under a condition on the spectra of the elements in the upper left-hand and lower right-hand corners of the above matrix.

LEMMA AND DEFINITION 2.1. *Let $T : \mathcal{H} \rightarrow \mathcal{H}$ be a bounded operator on a Hilbert space \mathcal{H} , let $r \geq 0$ and let $\mathcal{H}_r = \mathcal{H}_r(T)$ be the subspace (2) defined above. Then \mathcal{H}_r is an invariant subspace for T such that the orthogonal projection $p_r = p_r(T)$ from \mathcal{H} onto \mathcal{H}_r lies in the von Neumann algebra $\{T\}''$ generated by T .*

Proof. Consider the subset

$$E_r = E_r(T) = \left\{ \xi \in \mathcal{H} \mid \limsup_{k \rightarrow \infty} \|T^k \xi\|^{1/k} \leq r \right\} \quad (3)$$

which is dense in \mathcal{H}_r . To see that \mathcal{H}_r is a closed subspace of \mathcal{H} , it will suffice to show that E_r is a subspace. Let $a \in \mathbb{C}$ and $\xi_1, \xi_2 \in E_r$. Since $\|T^k(a\xi_1)\| = |a| \|T^k \xi_1\|$ and $\|T^k(\xi_1 + \xi_2)\| \leq 2 \max(\|T^k \xi_1\|, \|T^k \xi_2\|)$, it is clear that $a\xi_1 \in E_r$ and $\xi_1 + \xi_2 \in E_r$. Moreover, since $\|T^k(T\xi)\| \leq \|T\| \|T^k(\xi)\|$ it is clear that E_r , and hence also \mathcal{H}_r , is invariant for T .

To show that $p_r \in \{T\}''$ it will be enough to show that $Up_r = p_r U$ whenever U is a unitary operator on \mathcal{H} such that $UT = TU$. Moreover, $Up_r = p_r U$ will follow once we show that $U\xi \in E_r$ for every $\xi \in E_r$. But this holds because

$$\|T^k U\xi\| = \|UT^k \xi\| = \|T^k \xi\|. \quad \square$$

PROPOSITION 2.2. *Let $T : \mathcal{H} \rightarrow \mathcal{H}$ be a bounded operator on a Hilbert space \mathcal{H} . Suppose that e_1, e_2, e_3 are orthogonal projections in \mathcal{H} with $e_1 + e_2 + e_3 = 1$ and that e_1 and $e_1 + e_2$ are invariant for T . This means that T is upper triangular with respect to this decomposition of \mathcal{H} :*

$$T = \begin{pmatrix} e_1 T e_1 & * & * \\ 0 & e_2 T e_2 & * \\ 0 & 0 & e_3 T e_3 \end{pmatrix}.$$

Let $r \geq 0$ and suppose that

$$\sup \{ |\lambda| \mid \lambda \in \sigma(e_1 T e_1) \} \leq r < \inf \{ |\lambda| \mid \lambda \in \sigma(e_3 T e_3) \},$$

where $\sigma(e_j T e_j)$ denotes the spectrum of $e_j T e_j$ acting on $e_j \mathcal{H}$. Then the invariant subspace $\mathcal{H}_r(T)$ and its projection $p_r = p_r(T)$ defined in Lemma and Definition 2.1 satisfy

$$e_1 \leq p_r \leq e_1 + e_2.$$

Proof. If $\xi = e_1 \xi$ then $T^k \xi = (e_1 T e_1)^k \xi$ so $\|T^k \xi\|^{1/k} \leq (\|(e_1 T e_1)^k\| \|\xi\|)^{1/k}$ while $\lim_{k \rightarrow \infty} \|(e_1 T e_1)^k\|^{1/k} \leq r$. This shows $e_1 \leq p_r$.

Suppose $\xi \in \mathcal{H}$ and $e_3 \xi \neq 0$. Then $e_3 T^k \xi = (e_3 T e_3)^k \xi$. As an operator on $e_3 \mathcal{H}$, $e_3 T e_3$ is invertible and its inverse has spectral radius $< r^{-1}$. Therefore $\|T^k \xi\| \geq \|(e_3 T e_3)^k \xi\| \geq \|(e_3 T e_3)^{-k}\|^{-1} \|e_3 \xi\|$ and hence $\limsup_{k \rightarrow \infty} \|T^k \xi\|^{1/k} \geq \lim_{k \rightarrow \infty} \|(e_3 T e_3)^{-k}\|^{-1/k} > r$, so $\xi \notin E_r(T)$. This shows that $E_r(T) \perp e_3 \mathcal{H}$, and therefore that $p_r \leq e_1 + e_2$. \square

Invariant projections $p_r(T)$ and the dense subspaces $E_r(T)$ for some specific operators T are described in §6.

Now we show that for an element x of an abstract W^* -algebra, the projection $p_r(x)$ is defined independently of how the W^* -algebra is represented as a von Neumann algebra acting on a Hilbert space.

LEMMA 2.3. *Let \mathcal{H} and \mathcal{H}' be Hilbert spaces, let $T \in B(\mathcal{H})$ and take $r > 0$. Then*

$$p_r(T) \otimes 1_{\mathcal{H}'} = p_r(T \otimes 1_{\mathcal{H}'}). \quad (4)$$

Proof. Let $E_r(T)$ be given by (3) and let

$$E_r(T \otimes 1) = \{w \in \mathcal{H} \otimes \mathcal{H}' \mid \limsup_{k \rightarrow \infty} \|(T^k \otimes 1_{\mathcal{H}'}w)\|^{1/k} \leq r\}.$$

Letting \odot denote the algebraic tensor product of vector spaces, we clearly have $E_r(T) \odot \mathcal{H}' \subseteq E_r(T \otimes 1)$, so the inequality \leq holds in (4). Given a unit vector $\eta \in \mathcal{H}'$ let $V_\eta : \mathcal{H} \rightarrow \mathcal{H} \otimes \mathcal{H}'$ be $V_\eta(\xi) = \xi \otimes \eta$. Then $TV_\eta^* = V_\eta^*(T \otimes 1_{\mathcal{H}'}),$ so if $w \in E_r(T \otimes 1)$ then $V_\eta^*w \in E_r(T)$ for every η . Therefore $E_r(T \otimes 1) \subseteq \mathcal{H}_r(T) \otimes \mathcal{H}'$ and \geq holds in (4). \square

LEMMA AND DEFINITION 2.4. *If \mathcal{M} is a von Neumann algebra, if \mathcal{H} is a Hilbert space and if $\pi : \mathcal{M} \rightarrow B(\mathcal{H})$ is a normal, faithful $*$ -representation then given $x \in \mathcal{M}$ we have, for $r \geq 0$, the projection*

$$p_r(\pi(x)) \in \{\pi(x)\}'' \subseteq \pi(\mathcal{M}).$$

Let us denote by $p_r(x) \in \mathcal{M}$ the element so that $\pi(p_r(x)) = p_r(\pi(x))$. Then $p_r(x)$ is independent of the choice of \mathcal{H} and π .

Proof. Let \mathcal{H}' be a Hilbert space and $\pi' : \mathcal{M} \rightarrow B(\mathcal{H}')$ a normal, faithful $*$ -representation. Using [D, Theorem 1.4.3], we find a Hilbert space \mathcal{H}'' such that the representations $\pi \otimes 1_{\mathcal{H}''}$ and $\pi' \otimes 1_{\mathcal{H}''}$ are unitarily equivalent, via a unitary $U : \mathcal{H} \otimes \mathcal{H}'' \rightarrow \mathcal{H}' \otimes \mathcal{H}''$. Now applying Lemma 2.3 twice, we have

$$\begin{aligned} U^*(\pi'(p_r(x)) \otimes 1)U &= \pi(p_r(x)) \otimes 1 = p_r(\pi(x)) \otimes 1 = p_r(\pi(x) \otimes 1) \\ &= p_r(U^*(\pi'(x) \otimes 1)U) = U^*p_r(\pi'(x) \otimes 1)U = U^*(p_r(\pi'(x)) \otimes 1)U. \end{aligned}$$

Hence $\pi'(p_r(x)) = p_r(\pi'(x))$. \square

3 Upper Triangular Random Matrices

In this section, we consider upper triangular random matrices whose entries strictly above the diagonal are i.i.d. Gaussian, and we prove that the diagonal entries can be modified in various ways without affecting the limiting $*$ -moments as matrix size increases without bound. Throughout this paper, we consider random matrices whose entries have moments of all orders. Thus, let (Ω, μ) be a usual probability space, let $\mathcal{L} = \bigcap_{1 \leq p < \infty} L^p(\mu)$ and consider the expectation $E(f) = \int f d\mu$. If $S_1, S_2 \subseteq \mathcal{L}$ are sets of random variables, we say that S_1 and S_2 are *independent sets* if the two σ -algebras

generated by $\{f^{-1}(A) \mid f \in S_i, A \text{ Borel subset of } \mathbf{C}\}$ ($i = 1, 2$) are independent with respect to μ , and similarly for families of sets of random variables. The $*$ -algebra of $n \times n$ random matrices is $\mathcal{M}_n = \mathcal{L} \otimes M_n(\mathbf{C})$ and has the trace $\tau_n = E \otimes \text{tr}_n$, where tr_n is the trace on $M_n(\mathbf{C})$ normalized so that $\text{tr}_n(1) = 1$. We fix a system of matrix units in $M_n(\mathbf{C})$, denoted by $(e(i, j; n))_{1 \leq i, j \leq n}$.

NOTATION 3.1. Let $\sigma^2 > 0$ and $n \in \mathbf{N}$.

- (i) On \mathbf{C} , by Lebesgue measure we shall mean $d(\text{Re } z)d(\text{Im } z)$, i.e. normalized so that the unit disk has measure π . On the space $M_n(\mathbf{C})$ of $n \times n$ complex matrices, Lebesgue measure shall mean the product of Lebesgue measure on each of the n^2 complex entries. On the space $M_n^{s.a.}$ of self-adjoint complex $n \times n$ matrices, Lebesgue measure shall mean the product of Lebesgue measure on each of the $n(n-1)/2$ complex entries strictly above the diagonal and Lebesgue measure on each of the n real diagonal entries.
- (ii) We say that a random variable $a \in \mathcal{L}$ is a complex $(0, \sigma^2)$ -Gaussian if $\text{Re } a$ and $\text{Im } a$ are independent real Gaussian random variables each having first moment 0 and second moment $\sigma^2/2$. Thus $E(a) = 0$, $E(|a|^2) = \sigma^2$ and a has density $(\pi\sigma^2)^{-1}e^{-|z|^2/\sigma^2}$ with respect to Lebesgue measure.
- (iii) Given $T \in \mathcal{M}_n$, we will write $T \in \text{UTGRM}(n, \sigma^2)$ (the acronym is for “upper triangular Gaussian random matrix”) if the entries t_{ij} of T ($1 \leq i, j \leq n$) satisfy that $t_{ij} = 0$ whenever $i \geq j$ and that $(t_{ij})_{1 \leq i < j \leq n}$ is an independent family of random variables, each of which is complex $(0, \sigma^2)$ -Gaussian.

Our first result is that if $T(n) \in \text{UTGRM}(n, 1/n)$, if $D(n) \in \mathcal{M}_n$ is a diagonal random matrix that is independent of $T(n)$ and if the joint distribution of the diagonal entries of $D(n)$ is permutation invariant, then in the limit as $n \rightarrow \infty$, the $*$ -moments of $T(n) + D(n)$ depend only on the marginal $*$ -distributions of finite sets of the diagonal entries of $D(n)$.

Theorem 3.2. For every $n \in \mathbf{N}$ let $T(n) \in \text{UTGRM}(n, \frac{1}{n})$ and let

$$D_1(n) = \sum_{i=1}^n a(i; n) \otimes e(i, i; n) \in \mathcal{M}_n$$

$$D_2(n) = \sum_{i=1}^n b(i; n) \otimes e(i, i; n) \in \mathcal{M}_n$$

be diagonal random matrices such that $T(n)$ and $D_1(n)$ are independent matrix-valued random variables and $T(n)$ and $D_2(n)$ are independent

matrix-valued random variables. Let $d\nu_\iota(\lambda_1, \dots, \lambda_n)$ be the joint distribution of the diagonal entries of $D_\iota(n)$. Assume that $d\nu_\iota$ is invariant under all permutations of its n variables ($\iota = 1, 2$). Suppose that the marginal $*$ -distributions of the diagonal entries of $D_1(n)$ are arbitrarily close to those of $D_2(n)$ as $n \rightarrow \infty$; namely suppose that

$$\begin{aligned} \forall p \in \mathbf{N} \quad \forall r_1, s_1, \dots, r_p, s_p \in \mathbf{N} \cup \{0\}, \\ \lim_{n \rightarrow \infty} \left(E(a(1; n)^{r_1} \overline{a(1; n)^{s_1}} \cdots a(p; n)^{r_p} \overline{a(p; n)^{s_p}}) \right. \\ \left. - E(b(1; n)^{r_1} \overline{b(1; n)^{s_1}} \cdots b(p; n)^{r_p} \overline{b(p; n)^{s_p}}) \right) = 0. \end{aligned} \quad (5)$$

Let $Z_\iota(n) = D_\iota(n) + T(n)$. Then

$$\begin{aligned} \forall m \in \mathbf{N} \quad \forall \epsilon(1), \dots, \epsilon(m) \in \{*, \cdot\}, \\ \lim_{n \rightarrow \infty} \left(\tau_n(Z_1(n)^{\epsilon(1)} \cdots Z_1(n)^{\epsilon(m)}) - \tau_n(Z_2(n)^{\epsilon(1)} \cdots Z_2(n)^{\epsilon(m)}) \right) = 0, \end{aligned}$$

where we take $\epsilon(j) = \cdot$ to mean $\epsilon(j)$ is “not $*$ ” or “no symbol.” Therefore if $Z_1(n)$ converges in $*$ -moments as $n \rightarrow \infty$ then so does $Z_2(n)$ and their limit $*$ -moments coincide.

Proof. Write

$$T(n) = \sum_{1 \leq i < j \leq n} t(i, j; n) \otimes e(i, j; n).$$

Let us first fix $n \in \mathbf{N}$, $\iota \in \{1, 2\}$ and let us denote $D_\iota(n)$ simply by D , $T(n)$ by T , $Z_\iota(n)$ by Z and the diagonal entries of D by $d(i; n)$, ($1 \leq i \leq n$). Now each word in Z and Z^* is a sum of words in D , D^* , T and T^* ; hence we will investigate words of the form

$$\begin{aligned} W = (T^{\epsilon(1)} \cdots T^{\epsilon(l(1))}) D^{\kappa(1)} (T^{\epsilon(l(1)+1)} \cdots T^{\epsilon(l(2))}) D^{\kappa(2)} \\ \cdots (T^{\epsilon(l(p-1)+1)} \cdots T^{\epsilon(l(p))}) D^{\kappa(p)} (T^{\epsilon(l(p)+1)} \cdots T^{\epsilon(q)}) \end{aligned} \quad (6)$$

for arbitrary

$$\begin{aligned} p, q \in \mathbf{N} \cup \{0\}, \\ \epsilon(1), \dots, \epsilon(q), \kappa(1), \dots, \kappa(p) \in \{*, \cdot\}, \\ 0 \leq l(1) \leq l(2) \leq \cdots \leq l(p) \leq q. \end{aligned}$$

Now, using the independence of T and D , we see that

$$\begin{aligned} \tau_n(W) = n^{-1} \sum_{i_1, \dots, i_q \in \{1, \dots, n\}} \left(\prod_{j=1}^q G^{\epsilon(j)}(i_j, i_{j+1}) \right) \\ \cdot E(t^{\epsilon(1)}(i_1, i_2; n) t^{\epsilon(2)}(i_2, i_3; n) \cdots t^{\epsilon(q)}(i_q, i_{q+1}; n)) \\ \cdot E(d^{\kappa(1)}(i_{l(1)+1}; n) d^{\kappa(2)}(i_{l(2)+1}; n) \cdots d^{\kappa(p)}(i_{l(p)+1}; n)), \end{aligned} \quad (7)$$

where we have used the convention $i_{q+1} = i_1$, where $G^\epsilon(\cdot, \cdot)$ is defined by

$$G(i_j, i_{j+1}) = \begin{cases} 1 & \text{if } i_j < i_{j+1} \\ 0 & \text{otherwise} \end{cases}$$

$$G^*(i_j, i_{j+1}) = \begin{cases} 1 & \text{if } i_j > i_{j+1} \\ 0 & \text{otherwise} \end{cases}$$

and where

$$t^{\epsilon(j)}(i_j, i_{j+1}; n) = \begin{cases} t(i_j, i_{j+1}; n) & \text{if } \epsilon(j) = \cdot \\ \overline{t(i_{j+1}, i_j; n)} & \text{if } \epsilon(j) = * \end{cases}$$

$$d^{\kappa(j)}(i; n) = \begin{cases} d(i; n) & \text{if } \kappa(j) = \cdot \\ \overline{d(i; n)} & \text{if } \kappa(j) = * \end{cases}$$

Using that the joint distribution of $d(1; n), \dots, d(n; n)$ is invariant under permutation of the n variables, we see that each moment

$$E(d^{\kappa(1)}(i_{l(1)+1}; n) d^{\kappa(2)}(i_{l(2)+1}; n) \cdots d^{\kappa(p)}(i_{l(p)+1}; n)) \quad (8)$$

appearing in (7) is equal to a moment

$$E(d(1; n)^{r(1)} \overline{d(1; n)}^{s(1)} d(2; n)^{r(2)} \overline{d(2; n)}^{s(2)} \cdots d(p; n)^{r(p)} \overline{d(p; n)}^{s(p)}) \quad (9)$$

where

$$\begin{aligned} r(1), s(1), \dots, r(p), s(p) &\in \mathbf{N} \cup \{0\}, \\ r(1) + s(1) + \cdots + r(p) + s(p) &= p, \end{aligned} \quad (10)$$

and by further permutation the moment (8) corresponds to a unique moment of the form (9) if we make the additional stipulation that

$$r(1) + s(1) \geq r(2) + s(2) \geq \cdots \geq r(p) + s(p) \quad (11)$$

and, if $r(j) + s(j) = r(j+1) + s(j+1)$ then $r(j) \geq r(j+1)$.

Hence rearranging the sum in (7) we get

$$\begin{aligned} \tau_n(W) &= n^{-1} \sum_{r(1), s(1), \dots, r(p), s(p)} E(d(1; n)^{r(1)} \overline{d(1; n)}^{s(1)} \cdots d(p; n)^{r(p)} \overline{d(p; n)}^{s(p)}) \\ &\quad \cdot \sum_{i_1, \dots, i_q} \left(\prod_{j=1}^q G^{\epsilon(j)}(i_j, i_{j+1}) \right) E(t^{\epsilon(1)}(i_1, i_2; n) \cdots t^{\epsilon(q)}(i_q, i_{q+1}; n)), \end{aligned} \quad (12)$$

where the first sum is over all $r(1), s(1), \dots, r(p), s(p)$ satisfying (10) and (11), and the second sum is over all $i_1, \dots, i_q \in \{1, \dots, n\}$ such that there is a permutation, σ , of $\{1, \dots, n\}$ for which

$$\begin{aligned} d(\sigma(i_{l(1)+1}); n)^{\kappa(1)} d(\sigma(i_{l(2)+1}); n)^{\kappa(2)} \cdots d(\sigma(i_{l(p)+1}); n)^{\kappa(p)} \\ = d(1; n)^{r(1)} \overline{d(1; n)}^{s(1)} \cdots d(p; n)^{r(p)} \overline{d(p; n)}^{s(p)}. \end{aligned} \quad (13)$$

Let W_i ($i \in \{1, 2\}$) denote the word on the right-hand side of (6) where D is taken to be $D_i(n)$. Then

$$\begin{aligned} \tau_n(W_1) - \tau_n(W_2) \\ = n^{-1} \sum_{r(1), s(1), \dots, r(p), s(p)} \left(E(a(1; n)^{r(1)} \overline{a(1; n)}^{s(1)} \cdots a(p; n)^{r(p)} \overline{a(p; n)}^{s(p)}) \right. \\ \left. - E(b(1; n)^{r(1)} \overline{b(1; n)}^{s(1)} \cdots b(p; n)^{r(p)} \overline{b(p; n)}^{s(p)}) \right) \\ \cdot \sum_{i_1, \dots, i_q} \left(\prod_{j=1}^q G^{\epsilon(j)}(i_j, i_{j+1}) \right) E(t^{\epsilon(1)}(i_1, i_2; n) \cdots t^{\epsilon(q)}(i_q, i_{q+1}; n)). \end{aligned}$$

Now in order to prove the lemma it will suffice to show that $\lim_{n \rightarrow \infty} \tau_n(W_1) - \tau_n(W_2) = 0$. Since by the hypothesis (5), the difference of moments of a and b tends to zero as $n \rightarrow \infty$, it will suffice to show that for every $p, q \in \mathbf{N}$ and every $r(1), s(1), \dots, r(p), s(p) \in \mathbf{N} \cup \{0\}$, the quantity

$$n^{-1} \sum_{i_1, \dots, i_q} \left(\prod_{j=1}^q G^{\epsilon(j)}(i_j, i_{j+1}) \right) E(t^{\epsilon(1)}(i_1, i_2; n) \cdots t^{\epsilon(q)}(i_q, i_{q+1}; n)) \quad (14)$$

remains bounded as $n \rightarrow \infty$, where the sum is over all $i_1, \dots, i_q \in \{1, \dots, n\}$ such that there is a permutation, σ , of $\{1, \dots, n\}$ for which (13) holds. But this follows from the sort of counting arguments used by Voiculescu in [V2]. Indeed, for any $1 \leq i < i' \leq n$ and for any $m, m' \in \mathbf{N} \cup \{0\}$,

$$E(t(i, i'; n)^m \overline{t(i, i'; n)}^{m'}) \neq 0$$

implies $m = m'$. Taken together with the independence assumption on the entries of T , this shows that for any $i_1, \dots, i_q \in \{1, \dots, n\}$, a necessary condition so that

$$E(t^{\epsilon(1)}(i_1, i_2; n) \cdots t^{\epsilon(q)}(i_q, i_{q+1}; n)) \neq 0$$

is that there be a bijection, γ , from $\{1, \dots, q\}$ to itself, without fixed points, such that $\gamma^2 = \text{id}$ and

$$\forall j \in \{1, \dots, q\}, \quad i_{\gamma(j)} = i_{j+1}, \text{ and } i_{\gamma(j)+1} = i_j. \quad (15)$$

Moreover, there is a constant, c_1 , depending only on q , such that for all $n \in \mathbf{N}$ and all choices of i_1, \dots, i_q ,

$$|E(t^{\epsilon(1)}(i_1, i_2; n) \cdots t^{\epsilon(q)}(i_q, i_{q+1}; n))| \leq c_1 n^{-q/2}. \quad (16)$$

If γ is a bijection of $\{1, \dots, q\}$, let $N(\gamma, n)$ be the number of choices of $i_1, \dots, i_q \in \{1, \dots, n\}$ such that (15) holds. There are only finitely many bijections, γ , of $\{1, \dots, q\}$. Hence, in light of the bound (16), in order to show that (14) is bounded as $n \rightarrow \infty$, it will suffice to show that for each bijection γ of $\{1, \dots, q\}$ without fixed points such that $\gamma^2 = \text{id}$, the quantity

$$n^{-1-(q/2)} N(\gamma, n) \quad (17)$$

remains bounded as $n \rightarrow \infty$. However, $N(\gamma, n) \leq n^{d(\gamma)}$, where $d(\gamma)$ is the number of vertices in the quotient graph, G_γ , which is obtained from the q -gon graph by identifying the j th and $\gamma(j)$ th edges with opposite orientations, for every j . The graph G_γ has $q/2$ edges, hence at most $1 + (q/2)$ vertices, which shows that $d(\gamma) \leq 1 + (q/2)$ and hence that (17) remains bounded as $n \rightarrow \infty$. \square

Now we work on results that let us dispense with the permutation invariance assumed for the diagonal matrices of the previous theorem. Let \mathcal{U}_2 denote the group of unitary 2×2 complex matrices.

LEMMA 3.3. *There is a Borel function, $U : \mathbf{C}^3 \rightarrow \mathcal{U}_2$ such that for all $a, b, c \in \mathbf{C}$,*

$$U(a, b, c)^* \begin{pmatrix} a & b \\ 0 & c \end{pmatrix} U(a, b, c) = \begin{pmatrix} c & b \\ 0 & a \end{pmatrix}.$$

Proof. If $a = c$ then let $U(a, b, c) = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$. If $a \neq c$ but $b = 0$, then let $U(a, b, c) = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$. If $a \neq c$ and $b \neq 0$ then let

$$U(a, b, c) = \frac{1}{(|a - c|^2 + |b|^2)^{1/2}} \begin{pmatrix} b & \overline{(a - c)b/\bar{b}} \\ c - a & b \end{pmatrix}.$$

\square

LEMMA 3.4. *Fix $n \in \mathbf{N}$, let $T \in \text{UTGRM}(n, 1/n)$ and let*

$$D = \sum_{i=1}^n d(i) \otimes e(i, i; n) \in \mathcal{M}_n$$

be a diagonal random matrix. Suppose that T and D are independent matrix-valued random variables. Let π be a permutation of $\{1, \dots, n\}$ and let

$$D_\pi = \sum_{i=1}^n d(\pi(i)) \otimes e(i, i; n) \in \mathcal{M}_n. \quad (18)$$

Let $Z = D + T$ and $Z_\pi = D_\pi + T$. Then Z and Z_π have the same $$ -moments with respect to τ_n .*

Proof. We may without loss of generality assume that π is a transposition of neighbors:

$$\begin{aligned}\pi(k) &= k+1 \\ \pi(k+1) &= k \\ \pi(j) &= j \text{ if } j \notin \{k, k+1\}.\end{aligned}$$

We will use Lemma 3.3 to show that there is a unitary random matrix, $V \in \mathcal{M}_n$, such that V^*ZV has the same distribution as Z_π , and this will prove the theorem.

Recall that Ω is the usual probability space underlying our random matrices \mathcal{M}_n . For $\omega \in \Omega$ let $V(\omega)$ be the block diagonal matrix

$$V(\omega) = I_{k-1} \oplus U(d(k)(\omega), t(k, k+1)(\omega), d(k+1)(\omega)) \oplus I_{n-k-1}.$$

By this we mean that $V(\omega)$ has $k-1$ ones down the diagonal, then a 2×2 block that is the matrix U from Lemma 3.3, then $n-k-1$ ones. Let $x(i, j)$ denote the (i, j) th entry of the random matrix V^*ZV . If we write

$$T = \sum_{1 \leq i < j \leq n} t(i, j) \otimes e(i, j; n)$$

then

$$\begin{aligned}x(i, j) &= 0 && \text{if } i > j \\ x(i, i) &= d(i) && \text{if } i \notin \{k, k+1\} \\ x(k, k) &= d(k+1) \\ x(k+1, k+1) &= d(k) \\ x(k, k+1) &= t(k, k+1).\end{aligned}$$

Let

$$\begin{pmatrix} u_{11} & u_{12} \\ u_{21} & u_{22} \end{pmatrix} = U(d(k)(\omega), t(k, k+1)(\omega), d(k+1)(\omega)).$$

If $i < k$ then

$$\begin{aligned}x(i, k) &= t(i, k)u_{11} + t(i, k+1)u_{21} \\ x(i, k+1) &= t(i, k)u_{12} + t(i, k+1)u_{22}\end{aligned}$$

and if $j > k+1$ then

$$\begin{aligned}x(k, j) &= \overline{u_{11}}t(k, j) + \overline{u_{21}}t(k+1, j) \\ x(k+1, j) &= \overline{u_{12}}t(k, j) + \overline{u_{22}}t(k+1, j).\end{aligned}$$

In order to show that V^*ZV and Z_π have the same distribution, it is thus enough to show

- (a) $(x(i, j))_{1 \leq i < j \leq n}$ is an independent family of complex $(0, 1/n)$ -Gaussian random variables;
 (b) $\{x(i, i) \mid 1 \leq i \leq n\}$ and $\{x(i, j) \mid 1 \leq i < j \leq n\}$ are independent sets of random variables.

From the facts that

$$\{d(i) \mid 1 \leq i \leq n\} \cup \{t(k, k+1)\} \quad \text{and} \quad \{t(i, j) \mid 1 \leq i < j \leq n, (i, j) \neq (k, k+1)\} \quad (19)$$

are independent sets of random variables, each $t(i, j)$ is complex $(0, 1/n)$ -Gaussian and U is everywhere unitary and is independent of the lower set in (19), we see that

$$(x(i, j))_{1 \leq i < j \leq n, (i, j) \neq (k, k+1)} \quad (20)$$

is an independent family of complex $(0, 1/n)$ -Gaussian random variables. Moreover, the joint distribution of the family (20) is not changed by conditioning on the values of $d(1), \dots, d(n), t(k, k+1)$. Hence

$$\{d(i) \mid 1 \leq i \leq n\} \cup \{t(k, k+1)\} \quad \text{and} \quad \{x(i, j) \mid 1 \leq i < j \leq n, (i, j) \neq (k, k+1)\}$$

are independent sets of random variables. But this implies that (a) and (b) hold. \square

LEMMA 3.5. Let $D \in \mathcal{M}_n$ be a diagonal random matrix, let $T \in \text{UTGRM}(n, 1/n)$ and let $Z = D + T$. Let μ be the joint distribution of the n random variables, $d(1), d(2), \dots, d(n)$, in that order. For every permutation π of $\{1, \dots, n\}$ let μ_π be the joint distribution of the random variables, $d(\pi(1)), d(\pi(2)), \dots, d(\pi(n))$, in that order. Let A be a nonempty set of permutations of $\{1, \dots, n\}$ and let $|A|$ denote the cardinality of A . Consider the measure on \mathbf{C}^n ,

$$\tilde{\mu} = \frac{1}{|A|} \sum_{\pi \in A} \mu_\pi.$$

Let $\tilde{d}(1), \tilde{d}(2), \dots, \tilde{d}(n) \in \mathcal{L}$ be random variables whose joint distribution is $\tilde{\mu}$ and such that

$$\{\tilde{d}(i) \mid 1 \leq i \leq n\} \quad \text{and} \quad \{t(i, j) \mid 1 \leq i < j \leq n\}$$

are independent sets of random variables. Let

$$\tilde{D} = \sum_{i=1}^n \tilde{d}(i) \otimes e(i, i; n) \in \mathcal{M}_n$$

and let $\tilde{Z} = \tilde{D} + T(n)$. Then Z and \tilde{Z} have the same $*$ -moments with respect to τ_n .

Proof. We may introduce a uniformly distributed A -valued random variable, σ , that is independent of D . Then \tilde{D} has the same distribution as the random matrix, D_σ , which at a point $\omega \in \Omega$ takes the value

$$D_\sigma(\omega) = \sum_{i=1}^n d(\sigma_{(\omega)}(i))_{(\omega)} \otimes e(i, i; n).$$

Now each $*$ -moment of D_σ is the average over $\pi \in A$ of the corresponding $*$ -moments of the matrices D_π described in (18). By Lemma 3.4, each of these is in turn equal to the corresponding $*$ -moment of D . \square

Now we combine Theorem 3.2 and Lemma 3.5 to obtain this section's main result.

Theorem 3.6. *For each $n \in \mathbf{N}$ let $D_1(n), D_2(n) \in \mathcal{M}_n$ be diagonal random matrices, let $T(n) \in \text{UTGRM}(n, 1/n)$ and let $Z_\iota(n) = D_\iota(n) + T(n)$, ($\iota = 1, 2$). Suppose that $Z_1(n)$ converges in $*$ -moments as $n \rightarrow \infty$. For $\iota \in \{1, 2\}$ and $n \in \mathbf{N}$ let $\nu_\iota^{(n)}$ be the measure on \mathbf{C}^n that is the joint distribution of the n diagonal entries of $D_\iota(n)$, and let $\tilde{\nu}_\iota^{(n)}$ be the symmetrization of $\nu_\iota^{(n)}$. Suppose that for every $p \in \mathbf{N}$ and every $r(1), s(1), \dots, r(p), s(p) \in \mathbf{N} \cup \{0\}$,*

$$\lim_{n \rightarrow \infty} \left(\int_{\mathbf{C}^n} \lambda_1^{r(1)} \overline{\lambda_1}^{s(1)} \lambda_2^{r(2)} \overline{\lambda_2}^{s(2)} \dots \lambda_p^{r(p)} \overline{\lambda_p}^{s(p)} d\tilde{\nu}_1^{(n)}(\lambda_1, \dots, \lambda_n) \right. \\ \left. - \int_{\mathbf{C}^n} \lambda_1^{r(1)} \overline{\lambda_1}^{s(1)} \lambda_2^{r(2)} \overline{\lambda_2}^{s(2)} \dots \lambda_p^{r(p)} \overline{\lambda_p}^{s(p)} d\tilde{\nu}_2^{(n)}(\lambda_1, \dots, \lambda_n) \right) = 0.$$

Then also $Z_2(n)$ converges in $$ -moments as $n \rightarrow \infty$, and its limit $*$ -moments are the same as those of $Z_1(n)$.*

4 Asymptotically Free Random Matrices

This section concerns asymptotic freeness of self-adjoint i.i.d. Gaussian random matrices $Y(t, n)$ and certain diagonal random matrices that are independent of the $Y(t, n)$. Voiculescu, in his pioneering paper [V2], proved asymptotic freeness of such $Y(t, n)$ and constant diagonal matrices. Briefly stated, our innovation is to let the diagonal matrices be random. (See [Dy1] and [V3] for some other generalizations.) Using a technique based on the polar decomposition, Voiculescu parlayed his asymptotic freeness results for Gaussian random matrices into asymptotic freeness results for Haar distributed random unitary matrices. In a similar way we prove asymptotic freeness of Haar distributed random unitaries $U(r, n)$ and certain diagonal random matrices that are independent of the $U(r, n)$. Finally, this sec-

tion culminates in a result (Theorem 4.6) about matrix models for ($*$ -free families of) certain R -diagonal elements.

See Notation 3.1 for details of some terms used below.

NOTATION 4.1. (i) For a random matrix $X \in \mathcal{M}_n$, we will write $X \in \text{GRM}(n, \sigma^2)$ (the acronym is for “Gaussian random matrix”) if the entries x_{ij} of X ($1 \leq i, j \leq n$) satisfy that $(x_{ij})_{1 \leq i, j \leq n}$ is an independent family of random variables, each of which is complex $(0, \sigma^2)$ -Gaussian. Thus $X \in \text{GRM}(n, \sigma^2)$ if and only if X has density $(\pi\sigma^2)^{-n^2} \exp(-\frac{1}{\sigma^2} \text{Tr}(X^*X))$ with respect to Lebesgue measure on $M_n(\mathbb{C})$.

(ii) Given $Y \in \mathcal{M}_n$ and $\sigma^2 > 0$, we will write $Y \in \text{SGRM}(n, \sigma^2)$ (“self-adjoint Gaussian random matrix”) if the entries y_{ij} of Y ($1 \leq i, j \leq n$) satisfy that $y_{ij} = \overline{y_{ji}}$ for all i and j , that y_{ij} is complex $(0, \sigma^2)$ -Gaussian if $i \neq j$ and is real $(0, \sigma^2)$ -Gaussian if $i = j$ and that $(y_{ij})_{1 \leq i, j \leq n}$ is an independent family of random variables. Thus $Y \in \text{SGRM}(n, \sigma^2)$ if and only if Y has density $(\pi\sigma^2)^{-n^2/2} \exp(-\frac{1}{\sigma^2} \text{Tr}(Y^*Y))$ with respect to Lebesgue measure on $M_n^{s.a.}$.

(iii) Given $U \in \mathcal{M}_n$, we will write $U \in \text{HURM}(n)$ (“Haar unitary random matrix”) if U is a random unitary matrix distributed according to Haar measure on the $n \times n$ unitary matrices.

We begin with a preliminary result that is essentially just a combination of Theorems 2.1 and 2.2 of [V2], in the case of random diagonal matrices.

LEMMA 4.2. Let S and T be sets. For any $n \in \mathbf{N}$ let $Y(s, n) \in \text{SGRM}(n, 1/n)$ ($s \in S$), and consider diagonal random matrices $D(t, n) \in \mathcal{M}_n$ ($t \in T$). Suppose that for some $t_0 \in T$, $D(t_0, n) = I_n$ ($n \in \mathbf{N}$), that $\{D(t, n) \mid t \in T\}$ is closed under multiplication ($n \in \mathbf{N}$) and that $\{D(t, n) \mid t \in T\}$ converges in moments as $n \rightarrow \infty$. Suppose that for every $n \in \mathbf{N}$

$$(\{D(t, n) \mid t \in T\}, (\{Y(s, n)\})_{s \in S}) \quad (21)$$

is an independent family of sets of matrix-valued random variables. Finally, suppose that for every $m \in \mathbf{N}$, every $s_1, \dots, s_m \in S$ and every $t_1, \dots, t_m \in T$, the quantity

$$|\tau_n(Y(s_1, n)D(t_1, n) \cdots Y(s_m, n)D(t_m, n))| \quad (22)$$

remains bounded as $n \rightarrow \infty$. Then the following are equivalent:

- (i) The family (21) of sets of noncommutative random variables is asymptotically free as $n \rightarrow \infty$.
- (ii) Whenever $m \in \mathbf{N}$ is even, $t_1, \dots, t_m \in T$ are fixed and $\alpha: \{1, \dots, m\} \rightarrow S$ is such that for every $s \in S$, $\alpha^{-1}(s)$ has either two or zero elements,

(a) if $\alpha(1) = \alpha(2)$ then

$$\lim_{n \rightarrow \infty} (\tau_n(Y(\alpha(1), n)D(t_1, n)Y(\alpha(2), n)D(t_2, n) \cdots Y(\alpha(m), n)D(t_m, n)) - \tau_n(D(t_1, n))\tau_n(D(t_2, n)Y(\alpha(3), n)D(t_3, n) \cdots Y(\alpha(m), n)D(t_m, n))) = 0$$

(b) if $\alpha(p) \neq \alpha(p+1)$ for every $1 \leq p \leq m-1$ and if $\alpha(m) \neq \alpha(1)$ then

$$\lim_{n \rightarrow \infty} \tau_n(Y(\alpha(1), n)D(t_1, n)Y(\alpha(2), n)D(t_2, n) \cdots Y(\alpha(m), n)D(t_m, n)) = 0.$$

Proof. We may without loss of generality suppose $S = \mathbf{N}$ and $T = \mathbf{N}$. The implication (i) \implies (ii) follows from the last paragraph of [V2, 2.1] and the fact that the limit moments of each $Y(s, n)$ are those of a centered semicircle law with second moment 1.

To show (ii) \implies (i) we will use [V2, 2.1] and an idea from the proof of [V2, 2.2]. Let ω be a nontrivial ultrafilter on \mathbf{N} . On the algebra, $\mathbf{C}\langle (T_s)_{s \in \mathbf{N}}, (A_t)_{t \in \mathbf{N}} \rangle$, of polynomials in noncommuting variables $(T_s)_{s \in \mathbf{N}}$ and $(A_t)_{t \in \mathbf{N}}$, let ϕ_ω be the tracial linear functional defined by $\phi_\omega(p) = \lim_{n \rightarrow \omega} \tau_n(\pi_n(p))$, where $\pi_n : \mathbf{C}\langle (T_s)_{s \in \mathbf{N}}, (A_t)_{t \in \mathbf{N}} \rangle \rightarrow \mathcal{M}_n$ is the algebra homomorphism defined by $\pi_n(T_s) = Y(s, n)$ and $\pi_n(A_t) = D(t, n)$. Let Δ denote the subalgebra of $\mathbf{C}\langle (T_s)_{s \in \mathbf{N}}, (A_t)_{t \in \mathbf{N}} \rangle$ generated by 1 and $\{A_t \mid t \in \mathbf{N}\}$. We will check that the conditions 1° and 2° of [V2, 2.1] hold for the sequence $(T_s)_{s \in \mathbf{N}}$ and the subalgebra Δ with respect to ϕ_ω . Note that every element of Δ is a linear combination of words of the form $A_{t_1}A_{t_2} \cdots A_{t_k}$ and that $\pi_n(A_{t_1}A_{t_2} \cdots A_{t_k}) = D(t, n)$ for some $t \in \mathbf{N}$. Moreover, $\pi_n(1) = D(1, n)$. Therefore $\pi_n(\Delta) = \text{span}\{D(t, n) \mid t \in \mathbf{N}\}$ and hence condition 1° of [V2, 2.1] follows from the boundedness of (22) as $n \rightarrow \infty$.

To see that condition 2°a of [V2, 2.1] holds, it suffices to see that if $m \in \mathbf{N}$ and if $\alpha : \{1, \dots, m\} \rightarrow \mathbf{N}$ is such that $\alpha^{-1}(\{\alpha(1)\})$ has only one element and if $t_1, \dots, t_m \in \mathbf{N}$ then, $\forall n \in \mathbf{N}$,

$$\tau_n(Y(\alpha(1), n)D(t_1, n)Y(\alpha(2), n)D(t_2, n) \cdots Y(\alpha(m), n)D(t_m, n)) = 0. \quad (23)$$

But this follows from the independence of $Y(\alpha(1), n)$ from all the other matrices appearing in (23) and the fact that all entries of $Y(\alpha(1), n)$ have expectation zero.

Now conditions 2°b and 2°c of [V2, 2.1] follow from the hypotheses 2(a) and 2(b). Therefore, by [V2, 2.1], given an injection $\beta : \mathbf{N} \times \mathbf{N} \rightarrow \mathbf{N}$ and defining

$$X_{m,k} = k^{-1/2} \sum_{j=1}^k T_{\beta(m,j)},$$

the family of sets of noncommutative random variables,

$$(\Delta, (\{X_{m,k}\})_{m=1}^\infty)$$

is asymptotically free with respect to ϕ_ω as $k \rightarrow \infty$. However, using the Gaussianity of the entries of the $Y(s, n)$, we see that for every $k \in \mathbf{N}$, $(\Delta, (\{X_{m,k}\})_{m=1}^\infty)$ has the same moments as $(\Delta, (\{T_m\})_{m=1}^\infty)$. Hence $(\Delta, (\{T_m\})_{m=1}^\infty)$ is free with respect to ϕ_ω . Since ω was arbitrary, and since each $Y(s, n)$ converges in moments as $n \rightarrow \infty$, this implies that

$$(\{D(t, n) \mid t \in \mathbf{N}\}, (\{Y(s, n)\})_{s \in \mathbf{N}})$$

is asymptotically free as $n \rightarrow \infty$. \square

Theorem 4.3. *Let S and T be sets. For $s \in S$ and $n \in \mathbf{N}$ let $Y(s, n) \in \text{SGRM}(n, 1/n)$. For $t \in T$ and $n \in \mathbf{N}$ let*

$$D(t, n) = \sum_{i=1}^n d(i; n, t) \otimes e(i, i; n) \in \mathcal{M}_n$$

be a diagonal random matrix, and suppose, for some t and every n , $D(1, n) = I_n$, that $\{D(t, n) \mid t \in T\}$ is closed under multiplication and that $\{D(t, n) \mid t \in T\}$ converges in moments as $n \rightarrow \infty$. Suppose that for every $n \in \mathbf{N}$

$$(\{D(t, n) \mid t \in T\}, (\{Y(s, n)\})_{s \in S})$$

is an independent family of sets of matrix-valued random variables. Assume further that

- (i) *for every $t \in T$ and every $1 \leq p < \infty$,*

$$\sup_{\substack{n \in \mathbf{N} \\ 1 \leq i \leq n}} \|d(i; n, t)\|_p < \infty;$$

- (ii) *for every $m, n \in \mathbf{N}$, $m \leq n$, every $t_1, \dots, t_m \in T$ and every permutation, σ , of $\{1, \dots, m\}$, the joint distribution of*

$$(d(1; n, t_1), d(2; n, t_2), \dots, d(m; n, t_m))$$

is equal to the joint distribution of

$$(d(\sigma(1); n, t_1), d(\sigma(2); n, t_2), \dots, d(\sigma(m); n, t_m));$$

- (iii) *for every $p \in \mathbf{N}$, every $t_1, \dots, t_p \in T$ and every p -tuple, (i_1, \dots, i_p) , of distinct, positive integers, we have*

$$\lim_{n \rightarrow \infty} \left(E(d(i_1; n, t_1) d(i_2; n, t_2) \cdots d(i_p; n, t_p)) - \prod_{j=1}^p E(d(i_j; n, t_j)) \right) = 0.$$

Then the family

$$(\{D(t, n) \mid t \in T\}, (\{Y(s, n)\}_{s \in S}))$$

of sets of random variables converges in moments and is asymptotically free as $n \rightarrow \infty$.

Proof. We will apply Lemma 4.2. Let us first show that the quantity (22) remains bounded as $n \rightarrow \infty$. Write $a(i, j; n, s)$ for the (i, j) th entry of $Y(s, n)$. We have

$$\begin{aligned} & \tau_n(Y(s_1, n)D(t_1, n) \cdots Y(s_m, n)D(t_m, n)) \\ &= n^{-1} \sum_{i_1, \dots, i_m \in \{1, \dots, n\}} E(d(i_1; n, t_1)d(i_2; n, t_2) \cdots d(i_m; n, t_m)) \\ & \quad \cdot E(a(i_0, i_1; n, s_1)a(i_1, i_2; n, s_2) \cdots a(i_{m-1}, i_m; n, s_m)), \end{aligned} \quad (24)$$

where $i_0 = i_m$. Using the generalized Hölder inequality, we have

$$\begin{aligned} |E(d(i_1; n, t_1) \cdots d(i_m; n, t_m))| &\leq \|d(i_1; n, t_1) \cdots d(i_m; n, t_m)\|_1 \\ &\leq \prod_{j=1}^m \|d(i_j; n, t_j)\|_m. \end{aligned} \quad (25)$$

But by the assumption (i), there is $c_2 > 0$ such that

$$\forall n \in \mathbf{N} \quad \forall i_1, \dots, i_m \in \{1, \dots, n\}, \quad \prod_{j=1}^m \|d(i_j; n, t_j)\|_m \leq c_2. \quad (26)$$

Now consider

$$E(a(i_0, i_1; n, s_1)a(i_1, i_2; n, s_2) \cdots a(i_{m-1}, i_m; n, s_m)). \quad (27)$$

From the nature of the entries $a(i, j; n, s)$, we see that the quantity (27) can be nonzero only if there is a bijection $\gamma : \{1, \dots, m\} \rightarrow \{1, \dots, m\}$ such that $\gamma^2 = \text{id}$, γ has no fixed points and

$$\forall j \in \{1, \dots, m\}, \quad s_j = s_{\gamma(j)}, \quad i_j = i_{\gamma(j)-1}, \quad i_{j-1} = i_{\gamma(j)}.$$

One also calculates

$$\begin{aligned} & |E(a(i_0, i_1; n, s_1)a(i_1, i_2; n, s_2) \cdots a(i_{m-1}, i_m; n, s_m))| \\ & \leq \prod_{j=1}^m \|a(i_{j-1}, i_j; n, s_j)\|_m \leq n^{-m/2} \left(\frac{m}{2}\right)!. \end{aligned}$$

Let us call a bijection, γ , of $\{1, \dots, m\}$ without fixed points such that $\gamma^2 = \text{id}$, a *pairing* of $\{1, \dots, m\}$ and for every pairing γ let

$$\Theta(\gamma) = \{(i_1, \dots, i_m) \in \{1, \dots, n\}^m \mid \forall j \in \{1, \dots, m\}, i_j = i_{\gamma(j)-1}, i_{j-1} = i_{\gamma(j)}\}.$$

From the above estimates we obtain

$$\left| \tau_n(Y(s_1, n)D(t_1, n) \cdots Y(s_m, n)D(t_m, n)) \right| \leq c_2 \left(\frac{m}{2}\right)! n^{-(\frac{m}{2}+1)} \sum_{\gamma} |\Theta(\gamma)|, \quad (28)$$

where the sum is over all pairings, γ of $\{1, \dots, m\}$. To each pairing we associate the quotient graph, G_γ , of the clockwise oriented m -gon graph obtained by identifying with opposite orientation each pair of j th and $\gamma(j)$ th edges. The resulting graph has $m/2$ edges, hence at most $\frac{m}{2} + 1$ vertices. Consequently $|\Theta(\gamma)| \leq n^{\frac{m}{2}+1}$, and the quantity in (28) is bounded by $c_2 \left(\frac{m}{2}\right)!$ times the number of pairings, which is finite and independent of n . This shows that the quantities (22) remain bounded as $n \rightarrow \infty$.

We now show that (ii-a) and (ii-b) of Lemma 4.2 are satisfied. Let α be as described there. Then

$$\begin{aligned} & \tau_n(Y(\alpha(1), n)D(t_1, n) \cdots Y(\alpha(m), n)D(t_m, n)) \\ &= n^{-1} \sum_{i_1, \dots, i_m \in \{1, \dots, n\}} E(d(i_1; n, t_1)d(i_2; n, t_2) \cdots d(i_m; n, t_m)) \\ & \cdot E(a(i_0, i_1; n, \alpha(1))a(i_1, i_2; n, \alpha(2)) \cdots a(i_{m-1}, i_m; n, \alpha(m))) , \end{aligned} \quad (29)$$

where we let $i_0 = i_m$. Consider the clockwise oriented m -gon graph, label the edges consecutively e_1, e_2, \dots, e_m and the vertices v_1, v_2, \dots, v_m so that the vertices of the edge e_j are v_{j-1} and $v_j \pmod{m}$. Let G be the quotient of the m -gon graph obtained by identifying edges j and $\alpha(j)$ with opposite orientation ($1 \leq j \leq m$). The resulting identification of vertices of the m -gon graph gives an equivalence relation \sim on $\{v_1, \dots, v_m\}$ whose equivalence classes $F_1, F_2, \dots, F_{k(G)}$ are precisely the lists of indices labeling the $k(G)$ vertices of G . The expression

$$E(a(i_0, i_1; n, \alpha(1))a(i_1, i_2; n, \alpha(2)) \cdots a(i_{m-1}, i_m; n, \alpha(m))) \quad (30)$$

in (29) is nonzero if and only if whenever $1 \leq p, q \leq m$ and $v_p \sim v_q$ then $i_p = i_q$, and then the value of (30) is $n^{-m/2}$. For each equivalence class $F_j = \{v_{p(1)}, v_{p(2)}, \dots, v_{p(r_j)}\}$ of \sim there is $t'_j \in T$ so that $D(t'_j, n) = D(t_{p(1)}, n)D(t_{p(2)}, n) \cdots D(t_{p(r_j)}, n)$; for $p \in \{1, \dots, n\}$ let $d(p; n, t'_j)$ be the p th diagonal entry of $D(t'_j, n)$. Thus

$$\begin{aligned} & \tau_n(Y(\alpha(1), n)D(t_1, n) \cdots Y(\alpha(m), n)D(t_m, n)) \\ &= n^{-(\frac{m}{2}+1)} \sum_{p_1, \dots, p_{k(G)} \in \{1, \dots, n\}} E(d(p_1; n, t'_1)d(p_2; n, t'_2) \cdots d(p_{k(G)}; n, t'_{k(G)})) . \end{aligned} \quad (31)$$

Using the Hölder inequality estimate (25) and (26), we see that the terms

$$E(d(p_1; n, t'_1)d(p_2; n, t'_2) \cdots d(p_{k(G)}; n, t'_{k(G)}))$$

in (31) are uniformly bounded in modulus. Moreover, because G has $m/2$ edges, it follows that $k(G) \leq \frac{m}{2} + 1$. If α satisfies the hypothesis in 2(b) of Lemma 4.2, then every vertex of the m -gon graph is equivalent to at least one other vertex, so $k(G) \leq m/2$ and the quantity (31) tends to zero as $n \rightarrow \infty$, as required. We have proved that (ii-b) of Lemma 4.2 is satisfied.

Suppose that α satisfies the hypothesis in (ii-a) of Lemma 4.2, namely that $\alpha(1) = \alpha(2)$. We are interested in the limit of the moment (31) as $n \rightarrow \infty$. As the number of terms in the sum (31) where $p_i = p_j$ for some $i \neq j$ becomes negligibly small compared to $n^{\frac{m}{2}+1}$ as $n \rightarrow \infty$, we may in (31) sum over only all distinct choices of $p_1, \dots, p_{k(G)} \in \{1, \dots, n\}$. The assumption $\alpha(1) = \alpha(2)$ implies that v_1 is not equivalent to any other vertex under \sim . Therefore, renumbering if necessary, we may take $F_1 = \{v_1\}$ and hence $t'_1 = t_1$. By hypotheses (ii) and (iii), we have that

$$\begin{aligned} \delta_n &\stackrel{\text{def}}{=} E(d(p_1; n, t'_1)d(p_2; n, t'_2) \cdots d(p_{k(G)}; n, t'_{k(G)})) \\ &\quad - E(d(p_1; n, t_1))E(d(p_2; n, t'_2) \cdots d(p_{k(G)}; n, t'_{k(G)})) \end{aligned}$$

is independent of the choice of distinct $p_1, \dots, p_{k(G)} \in \{1, \dots, n\}$ and that $\delta_n \rightarrow 0$ as $n \rightarrow \infty$. Moreover, an analysis of the quotient graph G similar to that used to obtain (31), and keeping the same notation as in (31), shows that

$$\begin{aligned} \tau_n(D(t_2, n)Y(\alpha(3), n)D(t_3, n) \cdots Y(\alpha(m), n)D(t_m, n)) \\ = n^{-m/2} \sum_{p_2, \dots, p_{k(G)} \in \{1, \dots, n\}} E(d(p_2; n, t'_2) \cdots d(p_{k(G)}; n, t'_{k(G)})) . \end{aligned}$$

But then

$$\begin{aligned} n^{-(\frac{m}{2}+1)} \sum_{p_1, \dots, p_{k(G)} \in \{1, \dots, n\}} E(d(p_1; n, t_1))E(d(p_2; n, t'_2) \cdots d(p_{k(G)}; n, t'_{k(G)})) \\ = \left(n^{-1} \sum_{p_1=1}^n E(d(p_1; n, t_1)) \right) \\ \cdot \left(n^{-m/2} \sum_{p_2, \dots, p_{k(G)} \in \{1, \dots, n\}} E(d(p_2; n, t'_2) \cdots d(p_{k(G)}; n, t'_{k(G)})) \right) \\ = \tau_n(D(t_1, n))\tau_n(D(t_2, n)Y(\alpha(3), n)D(t_3, n) \cdots Y(\alpha(m), n)D(t_m, n)) . \end{aligned}$$

Taking the limit as $n \rightarrow \infty$, we can at will require p_j 's to be distinct and then relax this requirement; using that $\delta_n \rightarrow 0$, we obtain the conclusion

of (ii-a) of Lemma 4.2. \square

Following Voiculescu's proof [V2, Theorem 3.8], we will use polar decomposition to extend the asymptotic freeness result of Theorem 4.3 to include also Haar distributed random unitary matrices.

Theorem 4.4. *Let R, S and T be sets. For every $n \in \mathbf{N}$ and $s \in S$ let $Z(s, n) \in \text{GRM}(n, 1/n)$ and for every $r \in R$ let $U(r, n) \in \text{HURM}(n)$; furthermore, for every $t \in T$ let*

$$D(t, n) = \sum_{i=1}^n d(i; n, t) \otimes e(i, i; n) \in \mathcal{M}_n$$

be diagonal random matrices such that the family $(D(t, n))_{t \in T}$ is closed under multiplication and converges in moments as $n \rightarrow \infty$; assume further that the entries $d(i; n, t)$ satisfy the conditions (i), (ii) and (iii) of the statement of Theorem 4.3. Suppose that

$$((\{Z(s, n)\})_{s \in S}, (\{U(r, n)\})_{r \in R}, \{D(t, n) \mid t \in T\}) \quad (32)$$

is an independent family of sets of matrix-valued random variables. Then the family

$$((\{Z(s, n)^*, Z(s, n)\})_{s \in S}, (\{U(r, n)^*, U(r, n)\})_{r \in R}, \{D(t, n) \mid t \in T\}) \quad (33)$$

of sets of noncommutative random variables converges in moments and is asymptotically free as $n \rightarrow \infty$. Furthermore, each $Z(s, n)$ converges in $$ -moments to a circular element and each $U(r, n)$ converges in $*$ -moments to a Haar unitary, as $n \rightarrow \infty$.*

Proof. We shall use Theorem 4.3 and adapt the proof of [V2, Theorem 3.8] to our situation.

CLAIM 4.4.1. *The family*

$$((\{Z(s, n)^*, Z(s, n)\})_{s \in S}, \{D(t, n) \mid t \in T\})$$

of sets of random variables is asymptotically free as $n \rightarrow \infty$.

Proof. With

$$\begin{aligned} \operatorname{Re} Z(s, n) &= (Z(s, n) + Z(s, n)^*)/2 \\ \operatorname{Im} Z(s, n) &= (Z(s, n) - Z(s, n)^*)/2i, \end{aligned}$$

each of $\operatorname{Re} Z(s, n)$ and $\operatorname{Im} Z(s, n)$ is in $\text{SGRM}(n, 1/2n)$ and

$$((\{\operatorname{Re} Z(s, n)\})_{s \in S}, (\{\operatorname{Im} Z(s, n)\})_{s \in S}, \{D(t, n) \mid t \in T\}) \quad (34)$$

is an independent family of sets of matrix-valued random variables. Thus, by Theorem 4.3, each $\operatorname{Re} Z(s, n)$ and each $\operatorname{Im} Z(s, n)$ converges in moments to a semicircular element and the family (34) is asymptotically free as $n \rightarrow \infty$. This proves Claim 4.4.1. \square

Now take $W(r, n) \in \text{GRM}(n, 1/n)$ so that

$$((\{W(r, n)\})_{r \in R}, (\{Z(s, n)\})_{s \in S}, \{D(t, n) \mid t \in T\})$$

is an independent family of sets of matrix-valued random variables. By Claim 4.4.1,

$$((\{W(r, n)^*, W(r, n)\})_{r \in R}, (\{Z(s, n)^*, Z(s, n)\})_{s \in S}, \{D(t, n) \mid t \in T\}) \quad (35)$$

is asymptotically free as $n \rightarrow \infty$. If $W(r, n) = V(W(r, n)^*W(r, n))^{1/2}$ is the polar decomposition of $W(r, n)$, then V is almost everywhere a unitary, and is distributed according to Haar measure on the group of $n \times n$ unitaries. Therefore, letting $U(r, n)$ be the polar part, V , of $W(r, n)$, these random unitary matrices satisfy the hypotheses of the theorem. We will follow the proof of [V2, Theorem 3.8] to show the asymptotic freeness of (33).

For $\epsilon > 0$ let $Y_\epsilon(r, n) = W(r, n)(\epsilon + W(r, n)^*W(r, n))^{-1/2}$.

CLAIM 4.4.2. *For every $\epsilon > 0$, the family*

$$((\{Y_\epsilon(r, n)^*, Y_\epsilon(r, n)\})_{r \in R}, (\{Z(s, n)^*, Z(s, n)\})_{s \in S}, \{D(t, n) \mid t \in T\}) \quad (36)$$

is asymptotically free as $n \rightarrow \infty$.

Proof. Given $A \in \mathcal{M}_n$ and $1 \leq p < \infty$, let $|A|_p = (\tau_n(A^*A)^{p/2})^{1/p}$; moreover, let $|A|_\infty$ be the essential supremum of the operator norm of A evaluated at points of the underlying probability space. Let q be a noncommutative monomial in $d = 2a + 2b + c$ variables (for nonnegative integers a, b, c), with coefficient equal to 1. Let $0 < \delta \leq 1$. By Step I of the proof of [V2, 3.8], letting f be the function $f(t) = (\epsilon + t)^{-1/2}$, there is a polynomial Q_δ such that, letting

$$A_\delta(r, n) = W(r, n)Q_\delta(W(r, n)^*W(r, n)),$$

we have

$$\limsup_{n \rightarrow \infty} |A_\delta(r, n) - Y_\epsilon(r, n)|_d < \delta.$$

Because $|Y_\epsilon(r, n)|_d \leq |Y_\epsilon(r, n)|_\infty \leq 1$, it follows that $|A_\delta(r, n)|_d < 1 + \delta$.

The assumption (i) of Theorem 4.3 on the entries of $D(t, n)$ implies that for all $p \geq 1$ and for all $t \in T$, $\sup_{n \in \mathbb{N}} |D(t, n)|_p < \infty$. Moreover, the convergence in $*$ -moments as $n \rightarrow \infty$ of $Z(s, n)$ implies that $\sup_{n \geq 1} |Z(s, n)|_p < \infty$ whenever p is an even integer; however, as $|Z(s, n)|_p$ is increasing in p , this holds for all $1 \leq p < \infty$. Fix $r_1, \dots, r_a \in R$, $s_1, \dots, s_b \in S$, $t_1, \dots, t_c \in T$, and let

$$R_1(n, \epsilon) = q((Y_\epsilon(r_i, n))^*_{i=1}^a, (Y_\epsilon(r_i, n))_{i=1}^a, (Z(s_i, n))^*_{i=1}^b, \\ (Z(s_i, n))_{i=1}^b, (D(t_i, n))_{i=1}^c)$$

$$R_2(n, \epsilon, \delta) = q((A_\delta(r_i, n))^*_{i=1}^a, (A_\delta(r_i, n))_{i=1}^a, (Z(s_i, n))^*_{i=1}^b, \\ (Z(s_i, n))_{i=1}^b, (D(t_i, n))_{i=1}^c).$$

We may choose a constant K independent of δ and large enough so that

$$\forall i \in \{1, \dots, b\} \quad \limsup_{n \rightarrow \infty} |Z(s_i, n)|_d < K, \quad (37)$$

$$\forall i \in \{1, \dots, c\} \quad \limsup_{n \rightarrow \infty} |D(t_i, n)|_d < K. \quad (38)$$

Using Hölder's inequality we find

$$\limsup_{n \rightarrow \infty} |R_1(n, \epsilon) - R_2(n, \epsilon, \delta)|_1 \leq 2aK^{2b+c}(1 + \delta)^{2a-1}\delta,$$

and therefore

$$\lim_{\delta \rightarrow 0} \limsup_{n \rightarrow \infty} |\tau_n(R_1(n, \epsilon)) - \tau_n(R_2(n, \epsilon, \delta))| = 0. \quad (39)$$

The asymptotic freeness of

$$(\{A_\delta(r, n)^*, A_\delta(r, n)\}_{r \in R}, (\{Z(s, n)^*, Z(s, n)\}_{s \in S}, \{D(t, n) \mid t \in T\}))$$

follows from that of (35); this together with (39) implies the asymptotic freeness of (36), and claim 4.4.2 is proved. \square

Step III of the proof of [V2, 3.8] shows that for every $\theta > 0$ there is $\epsilon_0 > 0$ such that

$$\limsup_{n \rightarrow \infty} |Y_\epsilon(r, n) - U(r, n)|_d < \theta \quad (40)$$

whenever $0 < \epsilon \leq \epsilon_0$. Let q again be a noncommutative monomial having coefficient equal to 1 and with degree $d = 2a + 2b + c$, and let $r_1, \dots, r_a \in R$, $s_1, \dots, s_b \in S$, $t_1, \dots, t_c \in T$. Let

$$R_3(n) = q((U(r_i, n))^*_{i=1}^a, (U(r_i, n))_{i=1}^a, (Z(s_i, n))^*_{i=1}^b, \\ (Z(s_i, n))_{i=1}^b, (D(t_i, n))_{i=1}^c).$$

Letting K be a constant so that (37) and (38) hold, we easily see using (40) and Hölder's inequality that if $0 < \epsilon \leq \epsilon_0$ then

$$\limsup_{n \rightarrow \infty} |R_1(n, \epsilon) - R_3(n)|_1 \leq 2aK^{2b+c}\theta.$$

Therefore

$$\lim_{\epsilon \rightarrow 0} \limsup_{n \rightarrow \infty} |\tau_n(R_1(n, \epsilon)) - \tau_n(R_3(n))| = 0.$$

This, together with Claim 4.4.2 shows that the family (33) is asymptotically free as $n \rightarrow \infty$ and finishes the proof of the theorem. \square

The following sort of result is standard, but a proof is provided here for completeness.

LEMMA 4.5. *Let (A, ϕ) be a W^* -noncommutative probability space, let B be a unital subalgebra of A , let S be a set and for every $s \in S$ let $v_s \in A$ be a unitary with $\phi(v_s) = 0$. Suppose the family*

$$(B, (\{v_s^*, v_s\})_{s \in S}) \quad (41)$$

of $|S| + 1$ sets of noncommutative random variables is free. Then the family

$$(v_s B v_s^*)_{s \in S} \quad (42)$$

of unital subalgebras of A is free.

Proof. From $*$ -freeness of B and v_s we get $\phi(v_s b v_s^*) = \phi(b)$ for every $b \in B$. Let $n \in \mathbf{N}$ and let $s_1, \dots, s_n \in S$ be such that $s_j \neq s_{j+1}$ for every $j \in \{1, \dots, n-1\}$. For every $j \in \{1, \dots, n\}$ let $b_j \in B$ be such that $\phi(b_j) = 0$. In order for freeness of (42) to hold, it will suffice that

$$\phi((v_{s_1} b_1 v_{s_1}^*)(v_{s_2} b_2 v_{s_2}^*) \cdots (v_{s_n} b_n v_{s_n}^*)) = 0.$$

But the above equality follows directly from freeness of (41). \square

Now we apply the asymptotic freeness results proved previously in this section to give some matrix models for ($*$ -free families of) R -diagonal elements.

Theorem 4.6. *Let S be a set and for every $s \in S$ let $X_s(n) \in \mathcal{M}_n$ and let $\sigma_{s,n}$ be the symmetrized joint distribution of the eigenvalues of $(X_s^*(n)X_s(n))^{1/2}$. Given $p \in \{1, \dots, n\}$, let $\sigma_{s,n}^{(p)}$ be the marginal distribution of $\sigma_{s,n}$ corresponding to p of the variables. Suppose that for a compactly supported measure ρ_s on \mathbf{R}_+ and for every $p \in \mathbf{N}$, $\sigma_{s,n}^{(p)}$ converges in moments as $n \rightarrow \infty$ to the product measure $\times_1^p \rho_s$. Suppose also that for any non-random $n \times n$ unitary matrix U , the distributions of $UX_s(n)$ and of $X_s(n)$ are the same.*

- (i) *Fix $s \in S$. Then $X_s(n)$ converges in $*$ -moments to an element $u_s h_s$ of a noncommutative probability space, where u_s is a Haar unitary, $h_s \geq 0$, h_s has the same moments as the measure ρ_s and where the pair $(\{u_s, u_s^*\}, \{h_s\})$ of sets of noncommutative random variables is free.*
- (ii) *Suppose in addition that*

$$(X_s(n))_{s \in S} \quad (43)$$

is a mutually independent family of matrix-valued random variables and that the joint $*$ -moments of (43) are the same as the joint $*$ -moments of

$$(U_s^{(1)} X_s(n) U_s^{(2)})_{s \in S} \quad (44)$$

whenever $U_s^{(1)}$ and $U_s^{(2)}$ are non-random unitary matrices ($s \in S$). Then the family (43) is asymptotically $*$ -free as $n \rightarrow \infty$.

Proof. For brevity we shall prove parts (i) and (ii) simultaneously; while proving (i), we may from the outset assume that the stronger hypotheses of (ii) hold, because if we require S to be a single element then they will in any case be satisfied. We may write $X_s(n) = V_s(n) H_s(n)$ where $V_s(n)$ is a random unitary matrix and $H_s(n) = (X_s(n)^* X_s(n))^{1/2}$. For every $s \in S$ let $W_s(n)$ be a random unitary matrix so that $D_s(n) = W_s(n)^* H_s(n) W_s(n)$ is diagonal, so that the joint distribution of the diagonal entries of $D_s(n)$ is invariant under all permutations of the n variables and so that

$$(\{D_s(n), V_s(n), W_s(n)\})_{s \in S}$$

is a mutually independent family of sets of matrix-valued random variables. Let $U_s^{(1)}(n), U_s^{(2)}(n) \in \text{HURM}(n)$ be such that

$$((\{H_s(n), V_s(n), W_s(n)\})_{s \in S}, (\{U_s^{(1)}(n)\})_{s \in S}, (\{U_s^{(2)}(n)\})_{s \in S})$$

is a mutually independent family of sets of matrix-valued random variables. It follows from the hypotheses of (ii) that

$$(U_s^{(1)}(n) X_s(n) U_s^{(2)}(n))_{s \in S}$$

has the same joint $*$ -moments as the family (43). We have

$$\begin{aligned} & U_s^{(1)}(n) X_s(n) U_s^{(2)}(n) \\ &= (U_s^{(1)}(n) V_s(n) U_s^{(2)}(n)) (U_s^{(2)}(n)^* W_s(n) D_s(n) W_s(n)^* U_s^{(2)}(n)). \end{aligned}$$

Let

$$\begin{aligned} \tilde{V}_s(n) &= U_s^{(1)}(n) V_s(n) U_s^{(2)}(n) \\ \tilde{W}_s(n) &= U_s^{(2)}(n)^* W_s(n). \end{aligned}$$

Then $\tilde{V}_s(n), \tilde{W}_s(n) \in \text{HURM}(n)$ and

$$((D_s(n))_{s \in S}, (\tilde{V}_s(n))_{s \in S}, (\tilde{W}_s(n))_{s \in S}) \quad (45)$$

is an independent family of matrix-valued random variables.

Let

$$\Delta(n) = \{I_n\} \cup \{D_{s_1}(n) D_{s_2}(n) \cdots D_{s_q}(n) \mid q \in \mathbf{N}, s_1, \dots, s_q \in S\}.$$

By hypothesis, each $D_s(n)$ converges in moments as $n \rightarrow \infty$. Since $\Delta(n)$ forms a commuting family of self-adjoint random matrices, and since the family

$$(D_s(n))_{s \in S}, \quad (46)$$

is independent, it follows that $\Delta(n)$ converges in moments as $n \rightarrow \infty$; moreover, the subfamily (46) converges in $*$ -moments to a family $(d_s)_{s \in S}$ is some W^* -noncommutative probability space (A, ϕ) , where d_s is positive and has the same moments as the measure σ_s , and where for distinct $s_1, \dots, s_m \in S$ and any $k_1, \dots, k_m \in \mathbf{N}$,

$$\phi(d_{s_1}^{k_1} d_{s_2}^{k_2} \cdots d_{s_m}^{k_m}) = \prod_{j=1}^m \phi(d_{s_j}^{k_j}). \quad (47)$$

We shall show that the entries of the set $\Delta(n)$ of diagonal random matrices satisfy the properties (i), (ii) and (iii) in the statement of Theorem 4.3. Let $d_s(i, n)$ denote the i th diagonal entry of $D_s(n)$. Note that $E(d_s(i, n)^k)$ stays bounded (in fact converges) as $n \rightarrow \infty$, for every $k \in \mathbf{N}$ and $s \in S$. This, together with the independence of the family (46), implies condition (i). Condition (ii) follows from the independence of (46) and the fact that the joint distribution of the diagonal entries of each $D_s(n)$ is invariant under permutations of the n variables. Because $\sigma_{s,n}^{(p)}$ converges to a product measure, we have for every $s \in S$, $m \in \mathbf{N}$, $k_1, \dots, k_m \in \mathbf{N}$ and every p -tuple (i_1, i_2, \dots, i_m) of distinct, positive integers, that

$$\lim_{n \rightarrow \infty} \left(E(d_s(i_1, n)^{k_1} d_s(i_2, n)^{k_2} \cdots d_s(i_m, n)^{k_m}) - \prod_{j=1}^m E(d_s(i_j, n)^{k_j}) \right) = 0.$$

This together with the independence of (46) implies condition (iii). Hence we conclude from Theorem 4.4 that

$$(\{D_s(n) \mid s \in S\}, (\{\widetilde{V}_s(n)^*, \widetilde{V}_s(n)\}_{s \in S}, (\{\widetilde{W}_s(n)^*, \widetilde{W}_s(n)\}_{s \in S}))$$

is asymptotically free as $n \rightarrow \infty$. Therefore the family (45) converges in $*$ -moments to a family

$$((d_s)_{s \in S}, (v_s)_{s \in S}, (w_s)_{s \in S})$$

in some W^* -noncommutative probability space, where the joint $*$ -moments of $(d_s)_{s \in S}$ are as described above, where each v_s and each w_s is a Haar unitary and where

$$(\{d_s \mid s \in S\}, (\{v_s^*, v_s\}_{s \in S}, (\{w_s^*, w_s\}_{s \in S})) \quad (48)$$

is free. Therefore, the family (43) converges in $*$ -moments as $n \rightarrow \infty$ to the family

$$(v_s(w_s d_s w_s^*))_{s \in S}.$$

It is clear that $w_s d_s w_s^*$ has the same moments as d_s , namely the same moments as the measure σ_s . From the freeness of (48) and Lemma 4.5, it follows that the family

$$((v_s)_{s \in S}, (w_s d_s w_s^*)_{s \in S})$$

is $*$ -free, and the theorem is proved. \square

5 Upper triangular Representations of Circular Free Poisson Elements

In this section, the random matrix results of §3 and §4 are used, together with results of Dyson and others, to give upper triangular matrix models of circular free Poisson elements, and finally to give an upper triangular realization of a circular free Poisson element. An outline of the contents of this section is as follows: a first intermediate goal is a unitarily invariant matrix model for a circular free Poisson element (Theorem 5.4); next, a result of Dyson is quoted (Theorem 5.5) and used to convert the unitarily invariant matrix model to an upper triangular matrix model for a circular free Poisson element (Corollary 5.6); then the diagonal elements of this upper triangular matrix model are decoupled and desymmetrized so as to yield, in the limit as matrix size increases without bound, a triangular realization of a circular free Poisson element (Theorem 5.10).

The following is due to Bronk [B]; see also [HT, §5].

Theorem 5.1. *Let $c \geq 1$ and let Y be an $n \times n$ random matrix whose density with respect to Lebesgue measure on $M_n(\mathbf{C})$ is*

$$K_{c,n}^{(1)} |\det Y|^{2(c-1)n} \exp(-n \operatorname{Tr}(Y^* Y)),$$

where $K_{c,n}^{(1)}$ is a constant. Then the symmetrized joint distribution of the eigenvalues of $Y^* Y$ has density

$$K_{c,n}^{(2)} \left(\prod_{i=1}^n \lambda_i \right)^{2(c-1)n} \left(\prod_{1 \leq i < j \leq n} (\lambda_i - \lambda_j)^2 \right) \exp \left(-n \sum_{i=1}^n \lambda_i \right) \quad (49)$$

with respect to Lebesgue measure on $(\mathbf{R}_+)^n$, where $K_{c,n}^{(2)}$ is a constant.

The next theorem is a corollary of a result of Hewitt and Savage [HeS].

Theorem 5.2. *Let (Ω, \mathbf{E}) be a standard Borel space. Let σ be a Borel probability measure on the product set $\Omega^{\mathbf{N}} = \prod_{n=1}^{\infty} \Omega$ endowed with the product topology. Let σ_1 and σ_2 be the probability measures on Ω and $\Omega \times \Omega$, respectively, determined by*

$$\sigma_1(A) = \sigma(A \times \Omega \times \Omega \times \cdots), \quad A \in \mathbf{E}$$

$$\sigma_2(A_1 \times A_2) = \sigma(A_1 \times A_2 \times \Omega \times \Omega \times \cdots), \quad A_1, A_2 \in \mathbf{E}.$$

Suppose that

- (i) σ is invariant under all finite permutations of coordinates in $\Omega^{\mathbf{N}}$ (i.e. those permutations leaving all but finitely many coordinates fixed);
- (ii) $\sigma_2 = \sigma_1 \times \sigma_1$.

Then σ is equal to the product measure $\times_{n=1}^{\infty} \sigma_1$.

Proof. Since any noncountable standard Borel space is Borel isomorphic to the unit interval, and since $(\mathbf{N}, 2^{\mathbf{N}})$ is Borel isomorphic to the one-point compactification of \mathbf{N} , it is no loss of generality to assume that Ω is a separable compact Hausdorff space and \mathbf{E} is the Borel σ -algebra associated to this topology.

For any compact set K , let $P(K)$ denote the set of Borel probability measures on K . Consider the following subsets of $P(\Omega^{\mathbf{N}})$:

$$\begin{aligned} \tilde{P} &= \left\{ \bigotimes_{n=1}^{\infty} \mu \mid \mu \in P(\Omega) \right\}, \\ \tilde{S} &= \{ \nu \in P(\Omega^{\mathbf{N}}) \mid \nu \text{ is invariant under all finite permutations} \\ &\quad \text{of the coordinates of } \Omega^{\mathbf{N}} \}. \end{aligned}$$

Clearly $\tilde{P} \subseteq \tilde{S}$. By [HeS, Theorem 7.2], every $\nu \in \tilde{S}$ has a representation

$$\nu = \int_{P(\Omega)} \left(\bigotimes_{n=1}^{\infty} \mu \right) d\rho(\mu),$$

for a unique $\rho \in P(P(\Omega))$. In fact (see [S, Theorem 3.1]), \tilde{P} is the set of extreme points of the compact simplex \tilde{S} .

Now let σ be as in the formulation of the theorem. Using hypothesis (i) we have

$$\sigma = \int_{P(\Omega)} \left(\bigotimes_{n=1}^{\infty} \mu \right) d\rho(\mu)$$

for a unique $\rho \in P(P(\Omega))$. In particular

$$\sigma_1 = \int_{P(\Omega)} \mu d\rho(\mu) \quad \text{and} \quad \sigma_2 = \int_{P(\Omega)} (\mu \times \mu) d\rho(\mu).$$

By the assumption on Ω , the space $C(\Omega)$ of complex valued continuous functions on Ω is a separable Banach space (in the uniform norm), so we may let F be a countable dense subset of $C(\Omega)$. Given $f \in C(\Omega)$ and $\lambda \in P(\Omega)$ let us write

$$\lambda(f) = \int_{\Omega} f d\lambda.$$

With this notation, we have for all $f \in F$,

$$\begin{aligned} & \int_{P(\Omega)} |\mu(f) - \sigma_1(f)|^2 d\rho(\mu) \\ &= \int_{P(\Omega)} (\mu \times \mu)(f \otimes \bar{f}) d\rho(\mu) - 2\operatorname{Re} \left(\overline{\sigma_1(f)} \int_{P(\Omega)} \mu(f) d\rho(\mu) \right) + |\sigma_1(f)|^2 \\ &= \sigma_2(f \otimes \bar{f}) - 2\operatorname{Re} (\overline{\sigma_1(f)} \sigma_1(f)) + |\sigma_1(f)|^2 \\ &= \sigma_2(f \otimes \bar{f}) - |\sigma_1(f)|^2. \end{aligned}$$

But hypothesis (ii) shows that the above quantity is zero. Hence $\mu(f) = \sigma_1(f)$ for all $f \in F$, for ρ -almost all $\mu \in P(\Omega)$. Hence $\mu = \sigma_1$ for ρ -almost all $\mu \in P(\Omega)$, which implies $\rho = \delta_{\sigma_1}$, the Dirac measure at the point σ_1 . Therefore $\sigma = \times_{n=1}^{\infty} \sigma_1$. \square

LEMMA 5.3. *Let $c \geq 1$. Given $n \in \mathbf{N}$ let μ_n be the measure having density (49) with respect to Lebesgue measure on \mathbf{R}_+^n . Given $p \in \{1, 2, \dots, n\}$ let $\mu_n^{(p)}$ be the marginal distribution of μ_n corresponding to the variables $\lambda_1, \dots, \lambda_p$. Fix $p \in \mathbf{N}$. Then the distribution $\mu_n^{(p)}$ converges in the weak* topology as $n \rightarrow \infty$ to the product measure $\times_1^p \tau$, where τ has density with respect to Lebesgue measure*

$$\frac{d\tau}{d\lambda} = \frac{\sqrt{(\lambda - a)(b - \lambda)}}{2\pi\lambda} 1_{[a,b]}(\lambda), \quad (50)$$

with $a = (1 - \sqrt{c})^2$ and $b = (1 + \sqrt{c})^2$. Moreover, if f is a continuous function on $[0, \infty)^p$ with polynomial growth, in the sense that $f(t_1, \dots, t_p) \leq K^{(3)}(1 + t_1^{k_1} t_2^{k_2} \dots t_p^{k_p})$ for some constant $K^{(3)} > 0$ and positive integers k_1, \dots, k_p , then

$$\lim_{n \rightarrow \infty} \int_{\mathbf{R}_+^p} f d\mu_n^{(p)} = \int_{\mathbf{R}_+^p} f d\left(\times_1^p \tau\right). \quad (51)$$

Proof. It will be more convenient to consider the measure σ_n whose density with respect to Lebesgue measure on \mathbf{R}_+^n is

$$K_{c,n}^{(4)} \left(\prod_{i=1}^n \lambda_i \right)^{2(c-1)n} \left(\prod_{1 \leq i < j \leq n} (\lambda_i - \lambda_j)^2 \right) \exp \left(- \sum_{i=1}^n \lambda_i \right),$$

for some constant $K_{c,n}^{(4)}$; thus σ_n is the push forward measure of μ_n under the transformation $(\lambda_1, \lambda_2, \dots) \mapsto (n\lambda_1, n\lambda_2, \dots)$. We will find the limit as $n \rightarrow \infty$ of the marginal distributions, $\sigma_n^{(p)}$, of σ_n corresponding to the variables $\lambda_1, \dots, \lambda_p$. Let $\alpha = 2(c-1)n$ and let $\phi_0^{(\alpha)}, \phi_1^{(\alpha)}, \phi_2^{(\alpha)}, \dots$ be the polynomials obtained via Gram-Schmidt orthonormalization of $1, \lambda, \lambda^2, \dots$

in $L^2([0, \infty), \lambda^\alpha e^{-\lambda} d\lambda)$. Thus $\phi_k^{(\alpha)}(\lambda) = \sqrt{n!/\Gamma(\alpha + n + 1)} L_k^\alpha(\lambda)$, where L_k^α are the (generalized) Laguerre polynomials. Using the Vandermonde determinant we have

$$\begin{aligned} \prod_{1 \leq i < j \leq n} (\lambda_j - \lambda_i) &= \det \begin{pmatrix} 1 & 1 & \cdots & 1 \\ \lambda_1 & \lambda_2 & \cdots & \lambda_n \\ \vdots & \vdots & & \vdots \\ \lambda_1^{n-1} & \lambda_2^{n-1} & \cdots & \lambda_n^{n-1} \end{pmatrix} \\ &= K_{\alpha,n}^{(5)} \det \begin{pmatrix} \phi_0^{(\alpha)}(\lambda_1) & \phi_0^{(\alpha)}(\lambda_2) & \cdots & \phi_0^{(\alpha)}(\lambda_n) \\ \phi_1^{(\alpha)}(\lambda_1) & \phi_1^{(\alpha)}(\lambda_2) & \cdots & \phi_1^{(\alpha)}(\lambda_n) \\ \vdots & \vdots & & \vdots \\ \phi_{n-1}^{(\alpha)}(\lambda_1) & \phi_{n-1}^{(\alpha)}(\lambda_2) & \cdots & \phi_{n-1}^{(\alpha)}(\lambda_n) \end{pmatrix} \end{aligned}$$

for some constant $K_{\alpha,n}^{(5)}$. Therefore, the density of σ_n with respect to Lebesgue measure on \mathbf{R}_+^n is

$$D_n(\lambda_1, \dots, \lambda_n) = K_{c,n}^{(6)} \left(\prod_{i=1}^n \lambda_i \right)^\alpha \left(\sum_{\pi \in S_n} \text{sign}(\pi) \prod_{i=1}^n \phi_{\pi(i)-1}^{(\alpha)}(\lambda_i) \right)^2 \exp \left(- \sum_{i=1}^n \lambda_i \right),$$

for a constant $K_{c,n}^{(6)}$. Writing

$$\prod_{i=1}^n \phi_{\pi(i)-1}^{(\alpha)}(\lambda_i) = \phi_{\pi(1)-1}^{(\alpha)}(\lambda_1) \otimes \phi_{\pi(2)-1}^{(\alpha)}(\lambda_2) \otimes \cdots \otimes \phi_{\pi(n)-1}^{(\alpha)}(\lambda_n)$$

and noting that as π ranges over the permutation group S_n these form an orthonormal family with respect to the measure $(\prod_{i=1}^n \lambda_i)^\alpha \exp(-\sum_{i=1}^n \lambda_i) d\lambda_1 \cdots d\lambda_n$ on \mathbf{R}_+^n , we find $K_{c,n}^{(6)} = (n!)^{-1}$. Moreover, the density with respect to Lebesgue measure on \mathbf{R}_+ of the marginal distribution $\sigma_n^{(1)}$ is

$$D_{n,1}(\lambda_1) \stackrel{\text{def}}{=} \int_{\mathbf{R}_+^{n-1}} D_n(\lambda_1, \dots, \lambda_n) d\lambda_2 \cdots d\lambda_n = \frac{1}{n} \left(\sum_{k=0}^{n-1} \phi_k^{(\alpha)}(\lambda_1)^2 \right) \lambda_1^\alpha e^{-\lambda_1}.$$

But then the treatment in §6 of [HT] shows that $\mu_n^{(1)}$ converges in the weak* topology and in moments as $n \rightarrow \infty$ to τ .

The density with respect to Lebesgue measure on \mathbf{R}_+ of the marginal distribution $\sigma_n^{(2)}$ is

$$D_{n,2}(\lambda_1) \stackrel{\text{def}}{=} \int_{\mathbf{R}_+^{n-2}} D_n(\lambda_1, \dots, \lambda_n) d\lambda_3 \cdots d\lambda_n$$

$$\begin{aligned}
&= \frac{1}{n(n-1)} (\lambda_1 \lambda_2)^\alpha e^{-(\lambda_1 + \lambda_2)} \\
&\quad \cdot \sum_{\substack{0 \leq k, \ell \leq n-1 \\ k \neq \ell}} \phi_k^{(\alpha)}(\lambda_1) \phi_\ell^{(\alpha)}(\lambda_2) (\phi_k^{(\alpha)}(\lambda_1) \phi_\ell^{(\alpha)}(\lambda_2) - \phi_\ell^{(\alpha)}(\lambda_1) \phi_k^{(\alpha)}(\lambda_2)) \\
&= \frac{n}{n-1} D_{n,1}(\lambda_1) D_{n,1}(\lambda_2) \\
&\quad - \frac{1}{n(n-1)} (\lambda_1 \lambda_2)^\alpha e^{-(\lambda_1 + \lambda_2)} \left(\sum_{j=0}^{n-1} \phi_j^{(\alpha)}(\lambda_1) \phi_j^{(\alpha)}(\lambda_2) \right)^2.
\end{aligned}$$

We thus have, as elements of $C_0(\mathbf{R}_+^2)^*$,

$$\begin{aligned}
\|\sigma_n^{(2)} - \sigma_n^{(1)} \otimes \sigma_n^{(1)}\| &\leq \frac{1}{n-1} \int_0^\infty \int_0^\infty D_{n,1}(\lambda_1) D_{n,1}(\lambda_2) d\lambda_1 d\lambda_2 \\
&\quad + \frac{1}{n(n-1)} \sum_{j=0}^{n-1} \int_0^\infty \int_0^\infty \phi_j^{(\alpha)}(\lambda_1)^2 \phi_j^{(\alpha)}(\lambda_2)^2 (\lambda_1 \lambda_2)^\alpha e^{-(\lambda_1 + \lambda_2)} d\lambda_1 d\lambda_2 \\
&= \frac{2}{n-1}. \quad (52)
\end{aligned}$$

Since we know that $\mu_n^{(1)}$ converges in the weak* topology as $n \rightarrow \infty$ to τ , it follows from (52) that $\mu_n^{(2)}$ converges in weak* topology as $n \rightarrow \infty$ to $\tau \times \tau$.

Consider the measures $\tilde{\mu}_n = \mu_n \times \delta_0 \times \delta_0 \times \cdots$ on $[0, \infty)^{\mathbf{N}}$ and let ν be a w* cluster point in $C_0([0, \infty)^{\mathbf{N}})^*$ of these. Let $\nu^{(p)}$ be the marginal distribution of ν corresponding to the first p coordinates of $[0, \infty)^{\mathbf{N}}$. Then from what we have proved above we have

- ν is invariant under finite permutations of the coordinates in $[0, \infty)^{\mathbf{N}}$;
- $\nu^{(1)} = \tau$;
- $\nu^{(2)} = \tau \times \tau$.

Hence, by Theorem 5.2, $\nu = \times_{p=1}^\infty \nu^{(p)}$. Since ν was an arbitrary cluster point of $(\tilde{\mu}_n)_{n=1}^\infty$ it follows that $\tilde{\mu}_n$ converges in weak* topology to $\times_{p=1}^\infty \tau$ as $n \rightarrow \infty$. Therefore, for all $p \in \mathbf{N}$, the marginal distribution $\nu_n^{(p)}$ converges in weak* topology to the measure $\times_1^p \tau$ as $n \rightarrow \infty$.

It remains to show that (51) holds whenever f is of polynomial growth.

CLAIM 5.3.1. *Let $p \in \mathbf{N}$ and let h be a positive continuous function on $[0, \infty)^p$. Then $\liminf_{n \rightarrow \infty} \int h d\mu_n^{(p)} \geq \int h d\nu^{(p)}$.*

Proof. Choose $h_j \in C_0([0, \infty)^p)$, $h_j \geq 0$, so that h_j increases pointwise to h as $j \rightarrow \infty$. Then for all $j \geq 1$,

$$\liminf_{n \rightarrow \infty} \int h d\mu_n^{(p)} \geq \liminf_{n \rightarrow \infty} \int h_j d\mu_n^{(p)} = \int h_j d\nu^{(p)}.$$

But $\nu^{(p)} = \times_1^p \nu^{(1)}$ is supported on $[a, b]^p$; therefore $\sup_j \int h_j d\nu^{(p)} = \int h d\nu^{(p)}$, and the claim is proved. \square

CLAIM 5.3.2. *Let $p \in \mathbf{N}$ and suppose f and g are continuous functions on $[0, \infty)^p$ satisfying $g \geq 0$ and $-g \leq f \leq g$, and suppose that $\lim_{n \rightarrow \infty} \int g d\mu_n^{(p)} = \int g d\nu^{(p)}$. Then $\lim_{n \rightarrow \infty} \int f d\mu_n^{(p)} = \int f d\nu^{(p)}$.*

Proof. Applying Claim 5.3.1 to $g - f$ gives $\limsup_{n \rightarrow \infty} \int f d\mu_n^{(p)} \leq \int f d\nu^{(p)}$, while applying Claim 5.3.1 to $g + f$ yields $\liminf_{n \rightarrow \infty} \int f d\mu_n^{(p)} \geq \int f d\nu^{(p)}$; the claim is proved. \square

In order to finish the proof of the lemma, it will suffice to show

$$\lim_{n \rightarrow \infty} \int \lambda_1^{k_1} \lambda_2^{k_2} \cdots \lambda_p^{k_p} d\mu_n^{(p)} = \int \lambda_1^{k_1} \lambda_2^{k_2} \cdots \lambda_p^{k_p} d\nu^{(p)}, \quad (53)$$

for every $p \in \mathbf{N}$ and all integers $k_1, k_2, \dots, k_p \geq 0$. Letting $\ell = k_1 + k_2 + \cdots + k_p$, we have $0 \leq \lambda_1^{k_1} \lambda_2^{k_2} \cdots \lambda_p^{k_p} \leq 1 + \lambda_1^\ell + \lambda_2^\ell + \cdots + \lambda_p^\ell$. Moreover, because $\mu_n^{(1)}$ converges in moments to $\nu^{(1)}$, we have

$$\begin{aligned} \lim_{n \rightarrow \infty} \int (1 + \lambda_1^\ell + \lambda_2^\ell + \cdots + \lambda_p^\ell) d\mu_n^{(p)} &= 1 + p \lim_{n \rightarrow \infty} \int \lambda_1^\ell d\mu_n^{(1)} = 1 + p \int \lambda_1^\ell d\nu^{(1)} \\ &= \int (1 + \lambda_1^\ell + \lambda_2^\ell + \cdots + \lambda_p^\ell) d\nu^{(p)}. \end{aligned}$$

Now (53) follows from Claim 5.3.2, and the lemma is proved. \square

Theorem 5.4. *Let $c \geq 1$ and let $Y(n)$ be an $n \times n$ random matrix whose density with respect to Lebesgue measure on $M_n(\mathbf{C})$ is*

$$K_{c,n}^{(1)} |\det Y|^{2(c-1)n} \exp(-n \operatorname{Tr}(Y^* Y)).$$

Then $Y(n)$ converges in $$ -moments as $n \rightarrow \infty$ to a circular free Poisson element of parameter c .*

Proof. Clearly for every non-random $n \times n$ unitary matrix U , the distribution of $UY(n)$ is equal to the distribution of $Y(n)$. Let σ_n be the symmetrized joint distribution of the eigenvalues of $(Y(n)^* Y(n))^{1/2}$ and let μ_n be the symmetrized joint distribution of the eigenvalues of $Y(n)^* Y(n)$. For $p \in \{1, \dots, n\}$ let $\sigma_n^{(p)}$, respectively $\mu_n^{(p)}$, be the marginal distribution of σ_n , respectively μ_n , corresponding to the first p variables. Given $k_1, \dots, k_p \in \mathbf{N} \cup \{0\}$,

$$\int \lambda_1^{k_1} \lambda_2^{k_2} \cdots \lambda_p^{k_p} d\sigma_n^{(p)}(\lambda_1, \dots, \lambda_p) = \int \lambda_1^{k_1/2} \lambda_2^{k_2/2} \cdots \lambda_p^{k_p/2} d\mu_n^{(p)}(\lambda_1, \dots, \lambda_p).$$

By Theorem 5.1 and Lemma 5.3, it follows that

$$\lim_{n \rightarrow \infty} \int \lambda_1^{k_1} \lambda_2^{k_2} \cdots \lambda_p^{k_p} d\sigma_n^{(p)}(\lambda_1, \dots, \lambda_p) = \prod_{i=1}^p \int \lambda_i^{k_i/2} d\nu_c(\lambda_i),$$

where ν_c is the free Poisson distribution of parameter c . Therefore $\sigma_n^{(p)}$ converges in moments to $\times_1^p \rho$, where ρ has density

$$\frac{d\rho}{dt} = \frac{\sqrt{(d_1^2 - t^2)(t^2 - d_0^2)}}{\pi t} 1_{[d_0, d_1]}(t),$$

with $d_0 = 1 - \sqrt{c}$ and $d_1 = 1 + \sqrt{c}$. Now Theorem 4.6 applies and finishes the proof. \square

Every complex $n \times n$ matrix A is unitarily conjugate to an upper triangular matrix: $A = USU^*$, where U is unitary, the (i, j) th entry of S is zero if $i > j$ and the eigenvalues of A , listed according to generalized multiplicity, may appear in any order down the diagonal of S . If A has n distinct eigenvalues and if their order on the diagonal of S is specified, then the pair (U, S) is unique up to replacement by (UD, D^*SD) , where D is a diagonal unitary. Given a random matrix $X \in \mathcal{M}_n$, we may ask for a corresponding random upper triangular matrix S and random unitary matrix U so that the distribution of USU^* is equal to the distribution of X . Then X and S will have the same $*$ -moments with respect to the functional τ_n . For specificity, we may insist that the distribution of the diagonal entries of S be symmetric and the joint distribution of the pair (U, S) be the same as the joint distribution of (UD, D^*SD) for every non-random diagonal unitary D (i.e. that the joint distribution of (U, S) be invariant under this action of the n -torus \mathbf{T}^n). If the distribution of X is invariant under conjugation by non-random unitaries and if (U, S) is the pair of random matrices as described above, then it is clear that the random unitary U is distributed according to Haar measure on the $n \times n$ unitaries and that U and S are independent. In this case, the relevant question is only the distribution of S . F. Dyson answered this question when $X \in \text{GRM}(n, 1/n)$. We state his result, and then make a slight modification to give, in conjunction with Theorem 5.4, an upper triangular matrix model for a circular free Poisson element.

Theorem 5.5 (Dyson, see [M, A.35]). *Let $T(n) \in \text{UTGRM}(n, 1/n)$ and let $D(n) \in \mathcal{M}_n$ be a diagonal random matrix, whose diagonal entries have joint density*

$$K_n^{(7)} \exp \left(-n \sum_{i=1}^n |z_i|^2 \right) \prod_{1 \leq i < j \leq n} |z_i - z_j|^2 \quad (54)$$

with respect to Lebesgue measure on \mathbf{C}^n , for some constant $K_n^{(7)}$. Let $U(n) \in \text{HURM}(n)$, suppose that $(D(n), T(n), U(n))$ is an independent family of matrix-valued random variables and let

$$X(n) = U(n)(D(n) + T(n))U(n)^*.$$

Then $X(n) \in \text{GRM}(n, 1/n)$. Consequently, $D(n) + T(n)$ converges to a circular element in $*$ -moments as $n \rightarrow \infty$.

COROLLARY 5.6. Let $c \geq 1$, let $T(n) \in \text{UTGRM}(n, 1/n)$ and let $D_c(n) \in \mathcal{M}_n$ be a diagonal random matrix, whose diagonal entries have joint density

$$K_{c,n}^{(8)} \exp \left(-n \sum_{i=1}^n |z_i|^2 \right) \left(\prod_{i=1}^n |z_i| \right)^{2(c-1)n} \prod_{1 \leq i < j \leq n} |z_i - z_j|^2$$

with respect to Lebesgue measure on \mathbf{C}^n , for some constant $K_{c,n}^{(8)}$. Let $U(n) \in \text{HURM}(n)$, suppose that $(D_c(n), T(n), U(n))$ is an independent family of matrix-valued random variables and let

$$Y(n) = U(n)(D_c(n) + T(n))U(n)^*. \quad (55)$$

Then $Y(n)$ has density with respect to Lebesgue measure on $M_n(\mathbf{C})$ equal to

$$K_{c,n}^{(1)} |\det Y|^{2(c-1)n} \exp(-n \text{Tr}(Y^* Y)). \quad (56)$$

Consequently, $D_c(n) + T(n)$ converges, in $*$ -moments as $n \rightarrow \infty$, to a circular free Poisson element of parameter c .

Proof. Let \mathcal{M}_n be the manifold of matrices in $M_n(\mathbf{C})$ having n distinct eigenvalues. Then \mathcal{M}_n has full Lebesgue measure in $M_n(\mathbf{C})$. Let \mathcal{U}_n be the Lie group of $n \times n$ unitary matrices, and let \mathcal{T}_n be the manifold of all upper triangular $n \times n$ complex matrices, no two of whose diagonal elements are the same. Let $\pi : \mathcal{U}_n \times \mathcal{T}_n \rightarrow \mathcal{M}_n$ be given by $\pi(U, S) = USU^*$. Dyson proved his result by evaluating the Jacobian of π (after throwing away the directions in $\ker d\pi$) and thereby finding the measure σ_n on \mathcal{T}_n such that letting μ_n be Haar measure on \mathcal{U}_n , the push-forward measure $\pi_*(\mu_n \times \sigma_n)$ on \mathcal{M}_n has density $K_{1,n}^{(1)} \exp(-n \text{Tr}(Y^* Y))$ with respect to Lebesgue measure on \mathcal{M}_n , i.e. the density of a random matrix $X(n) \in \text{GRM}(n, 1/n)$. This measure σ_n was found to have density

$$K_n^{(9)} \prod_{1 \leq i < j \leq n} |S_{ii} - S_{jj}|^2 \exp(-n \text{Tr}(S^* S)) \quad (57)$$

with respect to Lebesgue measure on \mathcal{T}_n , where for a matrix $S \in \mathcal{T}_n$, S_{ii} is the i th diagonal entry of S ; this density (57) is that of the matrix $D(n) + T(n)$ in Theorem 5.5.

The matrix $D_c(n) + T(n)$ in the corollary has density

$$K_{c,n}^{(10)} \prod_{1 \leq i < j \leq n} |S_{ii} - S_{jj}|^2 \exp(-n \operatorname{Tr}(S^* S)) |\det(S)|^{2(c-1)n}$$

with respect to Lebesgue measure on \mathcal{T}_n ; since $\det(USU^*) = \det(S)$, and building on Dyson's calculation, it follows that the random matrix $Y(n)$ of (55) has density (56) with respect to Lebesgue measure on \mathcal{M}_n , as required.

An application of Theorem 5.4 shows that $Y(n)$, and hence also $D_c(n) + T(n)$, converges, in $*$ -moments as $n \rightarrow \infty$, to a circular free Poisson element. \square

The following lemma shows that the diagonal entries of $D_c(n)$ are in a specific sense asymptotically independent. This will allow their eventual decoupling (see Remark 5.9).

LEMMA 5.7. *For $c \geq 1$ and $n \in \mathbf{N}$ let μ_n be the probability measure on \mathbf{C}^n whose density with respect to Lebesgue measure is*

$$D_n(z_1, \dots, z_n) = K_{c,n}^{(8)} \exp\left(-n \sum_{i=1}^n |z_i|^2\right) \left(\prod_{i=1}^n |z_i|\right)^{2(c-1)n} \prod_{1 \leq i < j \leq n} |z_i - z_j|^2.$$

Given $p \in \{1, 2, \dots, n\}$ let $\mu_n^{(p)}$ be the marginal distribution of μ_n corresponding to the first p variables z_1, \dots, z_p . Then for every $p \in \mathbf{N}$, $\mu_n^{(p)}$ converges in weak topology and in $*$ -moments as $n \rightarrow \infty$ to the product measure $\times_1^p \rho$, where ρ is uniform distribution on the annulus $\{z \in \mathbf{C} \mid \sqrt{c-1} < |z| < \sqrt{c}\}$.*

Proof. This is quite similar to the proof of Lemma 5.3. Let $\alpha = (c-1)n$. Consider first the case $p = 1$. Let $\psi_0, \psi_1, \psi_2, \dots$ be the polynomials obtained via Gram-Schmidt orthonormalization of the sequence $1, z, z^2, \dots$ in $L^2(\mathbf{C}, |z|^{2\alpha} e^{-n|z|^2} d(\operatorname{Re} z) d(\operatorname{Im} z))$. Then

$$\psi_k(z) = \sqrt{\frac{n^{k+\alpha+1}}{\pi \Gamma(k+\alpha+1)}} z^k.$$

Using the Vandermonde determinant we have

$$\prod_{1 \leq i < j \leq n} (z_j - z_i) = K_{c,n}^{(11)} \det \begin{pmatrix} \psi_0(z_1) & \psi_0(z_2) & \cdots & \psi_0(z_n) \\ \psi_1(z_1) & \psi_1(z_2) & \cdots & \psi_1(z_n) \\ \vdots & \vdots & & \vdots \\ \psi_{n-1}(z_1) & \psi_{n-1}(z_2) & \cdots & \psi_{n-1}(z_n) \end{pmatrix}$$

for some constant $K_{c,n}^{(11)}$. Therefore

$$\begin{aligned} D_n(z_1, \dots, z_n) \\ = K_{c,n}^{(12)} \left(\prod_{i=1}^n |z_i| \right)^{2\alpha} \left| \sum_{\pi \in S_n} \text{sign}(\pi) \prod_{i=1}^n \psi_{\pi(i)-1}(z_i) \right|^2 \exp \left(-n \sum_{i=1}^n |z_i|^2 \right). \end{aligned}$$

Writing

$$\prod_{i=1}^n \psi_{\pi(i)-1}(z_i) = \psi_{\pi(1)-1}(z_1) \otimes \psi_{\pi(2)-1}(z_2) \otimes \cdots \otimes \psi_{\pi(n)-1}(z_n)$$

and noting that as π ranges over the permutation group S_n these form an orthonormal family with respect to the measure $\left(\prod_{i=1}^n |z_i| \right)^{2\alpha} \exp(-n \sum_{i=1}^n |z_i|^2)$

on \mathbf{C}^n , we find $K_{c,n}^{(12)} = (n!)^{-1}$. Moreover, the density of $\mu_n^{(1)}$ with respect to Lebesgue measure on \mathbf{C} is

$$\begin{aligned} D_{n,1}(z) &= \int_{\mathbf{C}^{n-1}} D_n(z, z_2, \dots, z_n) d(\text{Re } z_1) d(\text{Im } z_1) \cdots d(\text{Re } z_n) d(\text{Im } z_n) \\ &= \frac{1}{n} \left(\sum_{k=0}^{n-1} |\psi_k(z)|^2 \right) |z|^{2\alpha} e^{-n|z|^2} = \frac{n^\alpha}{\pi} \sum_{k=0}^{n-1} \frac{n^k |z|^{2k+2\alpha}}{\Gamma(k+\alpha+1)} e^{-n|z|^2}. \end{aligned}$$

We shall show that $\mu_n^{(1)}$ converges in $*$ -moments to ρ . Clearly if $a, b \in \mathbf{N} \cup \{0\}$ and if $a \neq b$ then

$$\int_{\mathbf{C}} z^a \bar{z}^b D_{n,1}(z) d(\text{Re } z) d(\text{Im } z) = 0 = \int_{\mathbf{C}} z^a \bar{z}^b d\rho(z).$$

Hence we need only show

$$\lim_{n \rightarrow \infty} \int_{\mathbf{C}} |z|^{2b} d\mu_n^{(1)}(z) = \int_{\mathbf{C}} |z|^{2b} d\rho(z)$$

for all $b \in \mathbf{N} \cup \{0\}$. We have

$$\begin{aligned} \int_{\mathbf{C}} |z|^{2b} d\mu_n^{(1)}(z) &= \frac{n^{(c-1)n}}{\pi} \sum_{k=0}^{n-1} \frac{n^k}{\Gamma(k+(c-1)n+1)} \\ &\quad \cdot \int_{\mathbf{C}} |z|^{2(b+k+(c-1)n)} e^{-n|z|^2} d(\text{Re } z) d(\text{Im } z) \\ &= \sum_{k=0}^{n-1} \frac{n^{k+(c-1)n}}{\Gamma(k+(c-1)n+1)} \int_0^\infty t^{b+k+(c-1)n} e^{-nt} dt. \end{aligned}$$

Writing

$$f_n(t) = \sum_{k=0}^{n-1} \frac{n^{k+(c-1)n}}{\Gamma(k+(c-1)n+1)} t^{b+k+(c-1)n} e^{-nt},$$

we have

$$\begin{aligned} \int_{\mathbf{C}} |z|^{2b} d\mu_n^{(1)}(z) &= \int_0^\infty t^b f_n(t) dt = -\frac{1}{b+1} \int_0^\infty t^{b+1} f'_n(t) dt \\ &= \frac{1}{b+1} \left(\frac{\Gamma(cn+b+1)}{n^{b+1}\Gamma(cn)} - \frac{\Gamma((c-1)n+b+1)}{n^{b+1}\Gamma((c-1)n)} \right) \\ &= \frac{1}{b+1} \left(c \left(c + \frac{1}{n} \right) \cdots \left(c + \frac{b}{n} \right) - (c-1) \left((c-1) + \frac{1}{n} \right) \cdots \left((c-1) + \frac{b}{n} \right) \right) \\ &\xrightarrow{n \rightarrow \infty} \frac{1}{b+1} (c^{b+1} - (c-1)^{b+1}) = \int_{\mathbf{C}} |z|^{2b} d\rho(z). \end{aligned}$$

Hence $\mu_n^{(1)}$ converges in $*$ -moments to ρ as $n \rightarrow \infty$; since ρ is compactly supported it follows that $\mu_n^{(1)}$ converges in the weak* topology to ρ .

The density of $\mu_n^{(2)}$ with respect to Lebesgue measure is

$$\begin{aligned} D_{n,2}(z_1, z_2) &= \int_{\mathbf{C}^{n-2}} D_n(z_1, z_2, \dots, z_n) d(\operatorname{Re} z_3) d(\operatorname{Im} z_3) \cdots d(\operatorname{Re} z_n) d(\operatorname{Im} z_n) \\ &= \frac{1}{n!} \sum_{\pi, \sigma \in S_n} \operatorname{sign}(\pi) \operatorname{sign}(\sigma) \int_{\mathbf{C}^{n-2}} \prod_{i=1}^n \left(\psi_{\pi(i)-1}(z_i) \overline{\psi_{\sigma(i)-1}(z_i)} |z_i|^{2\alpha} e^{-n|z_i|^2} \right) \\ &\quad \cdot d(\operatorname{Re} z_3) d(\operatorname{Im} z_3) \cdots d(\operatorname{Re} z_n) d(\operatorname{Im} z_n) \\ &= \frac{1}{n(n-1)} \sum_{\substack{0 \leq k, \ell \leq n-1 \\ k \neq \ell}} \left(|\psi_k(z_1)|^2 |\psi_\ell(z_2)|^2 - \psi_k(z_1) \psi_\ell(z_2) \overline{\psi_\ell(z_1) \psi_k(z_2)} \right) \\ &\quad \cdot |z_1|^{2\alpha} |z_2|^{2\alpha} e^{-n(|z_1|^2 + |z_2|^2)} \\ &= \frac{n}{n-1} D_{n,1}(z_1) D_{n,1}(z_2) - \frac{1}{n(n-1)} \left| \sum_{k=0}^{n-1} \psi_k(z_1) \overline{\psi_k(z_2)} \right|^2 \\ &\quad \cdot |z_1|^{2\alpha} |z_2|^{2\alpha} e^{-n(|z_1|^2 + |z_2|^2)}. \end{aligned}$$

Hence as a linear functional on $C_0(\mathbf{C}^2)$, the norm of $\mu_n^{(2)} - \mu_n^{(1)} \otimes \mu_n^{(1)}$ is bounded above by $2/(n-1)$. Therefore $\mu_n^{(2)}$ converges to $\rho \times \rho$ in the weak* topology as $n \rightarrow \infty$. Arguing as in the proof of Lemma 5.3 and using Theorem 5.2, we conclude that for every $p \geq 1$, $\mu_n^{(p)}$ converges in weak* topology to $\times_1^p \rho$, which we will denote by $\nu^{(p)}$.

It remains to show that $\mu_n^{(p)}$ converges to $\nu^{(p)}$ in $*$ -moments, namely that

$$\begin{aligned} \lim_{n \rightarrow \infty} \int z_1^{k_1} \overline{z_1}^{\ell_1} \cdots z_p^{k_p} \overline{z_p}^{\ell_p} d\mu_n^{(p)}(z_1, \dots, z_p) \\ = \int z_1^{k_1} \overline{z_1}^{\ell_1} \cdots z_p^{k_p} \overline{z_p}^{\ell_p} d\nu^{(p)}(z_1, \dots, z_p) \end{aligned} \quad (58)$$

for every $k_1, \dots, k_p, \ell_1, \dots, \ell_p \in \mathbf{N} \cup \{0\}$. Exactly as in the proof of Claim 5.3.1, we show that if h is a positive continuous function on \mathbf{C}^p then

$$\liminf_{n \rightarrow \infty} \int h d\mu_n^{(p)} \geq \int h d\nu^{(p)}.$$

Then, considering the real and imaginary parts separately and arguing as in the proof of Claim 5.3.2, we show that if f and g are continuous functions on \mathbf{C}^p , if $g \geq 0$, if $|f| \leq g$ and if $\lim_{n \rightarrow \infty} \int g d\mu_n^{(p)} = \int g d\nu^{(p)}$, then $\lim_{n \rightarrow \infty} \int f d\mu_n^{(p)} = \int f d\nu^{(p)}$. But letting $m = k_1 + \cdots + k_p + \ell_1 + \cdots + \ell_p$, we have

$$|z_1^{k_1} \overline{z_1}^{\ell_1} \cdots z_p^{k_p} \overline{z_p}^{\ell_p}| \leq 1 + |z_1|^{2m} + \cdots + |z_p|^{2m}.$$

Moreover, because $\mu_n^{(1)}$ converges in $*$ -moments to $\nu^{(1)}$, we have

$$\begin{aligned} \lim_{n \rightarrow \infty} \int (1 + |z_1|^{2m} + \cdots + |z_p|^{2m}) d\mu_n^{(p)} \\ = 1 + p \lim_{n \rightarrow \infty} \int |z_1|^{2m} d\mu_n^{(1)} = 1 + p \int |z_1|^{2m} d\nu^{(1)} \\ = \int (1 + |z_1|^{2m} + \cdots + |z_p|^{2m}) d\nu^{(p)}. \end{aligned}$$

Hence, we have (58) and the lemma is proved. \square

LEMMA 5.8. *Let $c \geq 1$ and for every $n \in \mathbf{N}$ let μ_n and μ'_n be the probability measures on \mathbf{C}^n whose densities with respect to Lebesgue measure are, respectively,*

$$D_n(z_1, \dots, z_n) = K_{c,n}^{(8)} \left(\prod_{i=1}^n |z_i| \right)^{2(c-1)n} \left(\prod_{1 \leq i < j \leq n} |z_i - z_j|^2 \right) \exp \left(-n \sum_{i=1}^n |z_i|^2 \right) \quad (59)$$

$$D'_n(z_1, \dots, z_n) = K_{c,n}^{(13)} \left(\prod_{i=1}^n |z_i| \right)^{2(c-1)n} \left(\sum_{\pi \in S_n} \prod_{i=1}^n |z_i|^{2(\pi(i)-1)} \right) \exp \left(-n \sum_{i=1}^n |z_i|^2 \right). \quad (60)$$

For $p \in \{1, \dots, n\}$ let $\mu_n^{(p)}$ and $(\mu'_n)^{(p)}$ denote the marginal distributions of μ_n and, respectively, μ'_n corresponding to the variables z_1, \dots, z_p . Then for

every p , $(\mu'_n)^{(p)}$ is obtained from $\mu_n^{(p)}$ by averaging over the action of the torus \mathbf{T}^p on \mathbf{C}^p given by coordinate-wise multiplication:

$$\mathbf{T}^p \times \mathbf{C}^p \ni ((w_1, \dots, w_p), (z_1, \dots, z_p)) \mapsto (w_1 z_1, \dots, w_p z_p).$$

Consequently, $(\mu'_n)^{(p)}$ converges to the measure $\times_1^p \rho$, in $*$ -moments and in weak* topology as $n \rightarrow \infty$, where ρ is the uniform distribution on the annulus $\{z \in \mathbf{C} \mid \sqrt{c-1} < |z| < \sqrt{c}\}$.

Proof. Using the Vandermonde determinant we find

$$\prod_{1 \leq i < j \leq n} |z_i - z_j|^2 = \sum_{\pi, \sigma \in S_n} \text{sign}(\pi) \text{sign}(\sigma) \prod_{i=1}^n z_i^{\pi(i)-1} \overline{z_i}^{\sigma(i)-1}. \quad (61)$$

Averaging (61) over the action of \mathbf{T}^n gives

$$\sum_{\pi \in S_n} \prod_{i=1}^n |z_i|^{2(\pi(i)-1)}.$$

From this we can easily see that $K_{c,n}^{(8)} = K_{c,n}^{(13)}$ and that μ'_n is obtained from μ_n by averaging. In order to show that $(\mu'_n)^{(p)}$ is obtained from $\mu_n^{(p)}$ by averaging, it suffices to note that for any measure τ on \mathbf{C}^n , the average over the action of \mathbf{T}^p on the marginal distribution, $\tau^{(p)}$, corresponding to the first p variables, is equal to the marginal distribution of the average over the action of \mathbf{T}^n on τ .

Since, by Lemma 5.7, $\mu_n^{(p)}$ converges in $*$ -moments and in weak* topology as $n \rightarrow \infty$ to $\times_1^p \rho$, which is invariant under the action of \mathbf{T}^p , it follows that $(\mu'_n)^{(p)}$ converges in $*$ -moments and in weak* topology to $\times_1^p \rho$. \square

REMARK 5.9. Our main purpose in proving the immediately preceding two lemmas was to be able to conclude that

$$\lim_{n \rightarrow \infty} \left(\int z_1^{k_1} \overline{z_1}^{\ell_1} \dots z_p^{k_p} \overline{z_p}^{\ell_p} d\mu_n^{(p)} - \int z_1^{k_1} \overline{z_1}^{\ell_1} \dots z_p^{k_p} \overline{z_p}^{\ell_p} d(\mu'_n)^{(p)} \right) = 0 \quad (62)$$

for every $p \in \mathbf{N}$ and every $k_1, \dots, k_p, \ell_1, \dots, \ell_p \in \mathbf{N} \cup \{0\}$. It is possible to prove (62) directly using the Vandermonde determinant and combinatorial arguments, though this sort of proof is not as satisfying as the one above involving Lemma 5.7, where the limit measure is found.

Theorem 5.10. Let $c \geq 1$ and $N \in \mathbf{N}$, and let (A, ϕ) be a W^* -noncommutative probability space with random variables $a_1, \dots, a_N \in A$ and $b_{ij} \in A$ ($1 \leq i < j \leq N$), where a_j is a circular free Poisson element of parameter $(c-1)N + j$, where each b_{ij} is a circular element with

$\phi(b_{ij}^* b_{ij}) = 1$, and where the family

$$((\{a_j^*, a_j\})_{1 \leq j \leq N}, (\{b_{ij}^*, b_{ij}\})_{1 \leq i < j \leq N})$$

is free. Consider the W^* -noncommutative probability space $(M_N(A), \phi_N)$, where

$$\phi_N((x_{ij})_{1 \leq i, j \leq N}) = N^{-1} \sum_{j=1}^N \phi(x_{jj}),$$

and consider the random variable

$$x = \frac{1}{\sqrt{N}} \begin{pmatrix} a_1 & b_{12} & b_{13} & \cdots & b_{1,N-1} & b_{1N} \\ 0 & a_2 & b_{23} & \cdots & b_{2,N-1} & b_{2N} \\ 0 & 0 & a_3 & \ddots & \vdots & b_{3N} \\ \vdots & \vdots & \ddots & \ddots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 & a_{N-1} & b_{N-1,N} \\ 0 & 0 & \cdots & 0 & 0 & a_N \end{pmatrix} \in M_N(A).$$

Then x is a circular free Poisson element of parameter c .

Proof. For every $n \in \mathbf{N}$ let $Y(nN)$ be an $nN \times nN$ random matrix whose distribution has density with respect to Lebesgue measure

$$K_{c,nN}^{(1)} |\det(Y)|^{2(c-1)nN} \exp(-nN \operatorname{Tr}(Y^* Y)).$$

Then by Theorem 5.4, $Y(nN)$ converges in $*$ -moments as $n \rightarrow \infty$ to a circular free Poisson element of parameter c . By Corollary 5.6, each $Y(nN)$ has the same $*$ -moments as $S^{(1)}(nN) \stackrel{\text{def}}{=} D^{(1)}(nN) + T(nN)$, where $T(nN) \in \text{UTGRM}(nN, 1/nN)$, $D^{(1)}(nN)$ is a diagonal $nN \times nN$ random matrix, the distribution of whose diagonal entries has density

$$K_{c,nN}^{(8)} \left(\prod_{i=1}^{nN} |z_i| \right)^{2(c-1)nN} \left(\prod_{1 \leq i < j \leq nN} |z_i - z_j|^2 \right) \exp \left(-nN \sum_{i=1}^n |z_i|^2 \right) \quad (63)$$

with respect to Lebesgue measure on \mathbf{C}^n , and where $D^{(1)}(nN)$ and $T(nN)$ are independent. We will use previous results to show that each $S^{(k)}(nN) \stackrel{\text{def}}{=} D^{(k)}(nN) + T(nN)$ ($k \in \{2, 3, 4, 5, 6\}$) also converges in $*$ -moments as $n \rightarrow \infty$ to a circular free Poisson element of parameter c , where $D^{(k)}(nN)$ is a diagonal random matrix such that $D^{(k)}(nN)$ and $T(nN)$ are independent and where the joint distributions of the diagonal entries of $D^{(k)}(nN)$ have,

in various cases, the following densities with respect to Lebesgue measure on \mathbf{C}^n :

$$\begin{aligned}
 k = 2 : \quad & K_{c,nN}^{(13)} \left(\prod_{i=1}^{nN} |z_i| \right)^{2(c-1)nN} \left(\sum_{\pi \in S_{nN}} \prod_{i=1}^{nN} |z_i|^{2(\pi(i)-1)} \right) \exp \left(-nN \sum_{i=1}^{nN} |z_i|^2 \right) \\
 k = 3 : \quad & K_{c,nN}^{(14)} \left(\prod_{i=1}^{nN} |z_i| \right)^{2(c-1)nN} \left(\prod_{i=1}^{nN} |z_i|^{2(i-1)} \right) \exp \left(-nN \sum_{i=1}^{nN} |z_i|^2 \right) \\
 k = 4 : \quad & K_{c,nN}^{(15)} \prod_{j=1}^N \left(\left(\prod_{i=1}^n |z_{(j-1)n+i}| \right)^{2((c-1)N+j-1)n} \left(\prod_{i=1}^n |z_{(j-1)n+i}|^{2(i-1)} \right) \right. \\
 & \quad \left. \cdot \exp \left(-nN \sum_{i=1}^n |z_{(j-1)n+i}|^2 \right) \right) \\
 k = 5 : \quad & K_{c,nN}^{(16)} \prod_{j=1}^N \left(\left(\prod_{i=1}^n |z_{(j-1)n+i}| \right)^{2((c-1)N+j-1)n} \right. \\
 & \quad \left. \cdot \left(\sum_{\pi \in S_n} \prod_{i=1}^n |z_{(j-1)n+i}|^{2(\pi(i)-1)} \right) \exp \left(-nN \sum_{i=1}^n |z_{(j-1)n+i}|^2 \right) \right) \\
 k = 6 : \quad & K_{c,nN}^{(17)} \prod_{j=1}^N \left(\left(\prod_{i=1}^n |z_{(j-1)n+i}| \right)^{2((c-1)N+j-1)n} \right. \\
 & \quad \left. \cdot \left(\prod_{1 \leq i < i' \leq n} |z_{(j-1)n+i} - z_{(j-1)n+i'}|^2 \right) \exp \left(-nN \sum_{i=1}^n |z_{(j-1)n+i}|^2 \right) \right).
 \end{aligned}$$

The proof that $S^{(k)}(nN)$ converges in $*$ -distribution to a circular free Poisson element relies for $k = 2$ on Lemmas 5.7 and 5.8 (see Remark 5.9), and Theorem 3.2; for $k = 3$ we use Theorem 3.6; the density for $k = 4$ is just a rewriting of that for $k = 3$; for $k = 5$ we again use Theorem 3.6; for $k = 6$ we again use Lemmas 5.7 and 5.8, and Theorem 3.2.

We may characterize the above successive transformations as follows: from (63) to $k = 2$ is decoupling; from $k = 2$ to $k = 3$ is desymmetrization; from $k = 3$ to $k = 4$ is regrouping; from $k = 4$ to $k = 5$ is partial resymmetrization; from $k = 5$ to $k = 6$ is partial recoupling.

Taking blocks of consecutive rows and columns to write $D^{(6)}(nN) + T(nN)$ as an $N \times N$ matrix of $n \times n$ random matrices, we have

$$S^{(6)}(nN) = \frac{1}{\sqrt{N}} \begin{pmatrix} A_1^{(6)} & B_{12}^{(6)} & B_{13}^{(6)} & \cdots & B_{1,N-1}^{(6)} & B_{1N}^{(6)} \\ 0 & A_2^{(6)} & B_{23}^{(6)} & \cdots & B_{2,N-1}^{(6)} & B_{2N}^{(6)} \\ 0 & 0 & A_3^{(6)} & \ddots & \vdots & B_{3N}^{(6)} \\ \vdots & \vdots & \ddots & \ddots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 & A_{N-1}^{(6)} & B_{N-1,N}^{(6)} \\ 0 & 0 & \cdots & 0 & 0 & A_N^{(6)} \end{pmatrix},$$

where

$$((A_j^{(6)}(n))_{1 \leq j \leq N}, (B_{ij}^{(6)}(n))_{1 \leq i < j \leq N})$$

is an independent family of matrix-valued random variables, where $B_{ij}^{(6)}(n) \in \text{GRM}(n, 1/n)$ for every $1 \leq i < j \leq N$ and where $A_j^{(6)}(n) = D_j^{(6)}(n) + T_j(n)$ with $T_j(n) \in \text{UTGRM}(n, 1/n)$, with $D_j^{(6)}(n)$ a diagonal random matrix, the joint distribution of whose diagonal entries has density with respect to Lebesgue measure

$$K_{(c-1)N+j,n}^{(8)} \left(\prod_{i=1}^n |z_i| \right)^{2((c-1)N+j-1)n} \left(\prod_{1 \leq i < i' \leq n} |z_i - z_{i'}|^2 \right) \exp \left(-n \sum_{i=1}^n |z_i|^2 \right)$$

and with $D_j^{(6)}(n)$ and $T_j(n)$ independent.

Let $U_j(n) \in \text{HURM}(n)$, ($1 \leq j \leq N$), be such that

$$((A_j^{(6)}(n))_{1 \leq j \leq N}, (B_{ij}^{(6)}(n))_{1 \leq i < j \leq N}, (U_j(n))_{1 \leq j \leq N})$$

is an independent family of matrix-valued random variables. By conjugating the matrix $S^{(6)}(nN)$ with $\text{diag}(U_1(n), U_2(n), \dots, U_N(n))$ and by using Corollary 5.6 and the fact that the class $\text{GRM}(n, 1/n)$ is invariant under left and right multiplication by independent unitaries, it follows that $S^{(6)}(nN)$ has the same $*$ -moments, as

$$S^{(7)}(nN) = \frac{1}{\sqrt{N}} \begin{pmatrix} A_1^{(7)} & B_{12}^{(7)} & B_{13}^{(7)} & \cdots & B_{1,N-1}^{(7)} & B_{1N}^{(7)} \\ 0 & A_2^{(7)} & B_{23}^{(7)} & \cdots & B_{2,N-1}^{(7)} & B_{2N}^{(7)} \\ 0 & 0 & A_3^{(7)} & \ddots & \vdots & B_{3N}^{(7)} \\ \vdots & \vdots & \ddots & \ddots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 & A_{N-1}^{(7)} & B_{N-1,N}^{(7)} \\ 0 & 0 & \cdots & 0 & 0 & A_N^{(7)} \end{pmatrix},$$

where

$$((A_j^{(7)}(n))_{1 \leq j \leq N}, (B_{ij}^{(7)}(n))_{1 \leq i < j \leq N}) \quad (64)$$

is an independent family of matrix-valued random variables, where $B_{ij}^{(7)}(n) \in \text{GRM}(n, 1/n)$ for every $1 \leq i < j \leq N$ and where the distribution of $A_j^{(7)}(n)$ has density

$$K_{(c-1)N+j,n}^{(1)} |\det(A)|^{2((c-1)N+j-1)n} \exp(-n \text{Tr}(A^* A))$$

with respect to Lebesgue measure on $M_n(\mathbf{C})$.

If $(V_j^{(1)})_{1 \leq j \leq N}$, $(V_j^{(2)})_{1 \leq j \leq N}$, $(U_{ij}^{(1)})_{1 \leq i < j \leq N}$, $(U_{ij}^{(2)})_{1 \leq i < j \leq N}$ are non-random $n \times n$ unitary matrices, then

$$\left((V_j^{(1)} A_j^{(7)}(n) V_j^{(2)})_{1 \leq j \leq N}, (U_{ij}^{(1)} B_{ij}^{(7)}(n) U_{ij}^{(2)})_{1 \leq i < j \leq N} \right) \quad (65)$$

continues to be an independent family of matrix-valued random variables, $V_j^{(1)} A_j^{(7)}(n) V_j^{(2)}$ has the same distribution as $A_j^{(7)}$ and $U_{ij}^{(1)} B_{ij}^{(7)}(n) U_{ij}^{(2)}$ has the same distribution as $B_{ij}^{(7)}(n)$. Therefore, the family (65) has the same joint $*$ -moments as the family (64). Taking into account also Theorem 5.1 and Lemma 5.3 (as in the proof of Theorem 5.4), we see that the conditions of Theorem 4.6 are fulfilled, allowing us to conclude that the family (64) is asymptotically $*$ -free as $n \rightarrow \infty$. Moreover (by Theorem 5.4), each $A_j^{(7)}(n)$ converges in $*$ -moments to a circular free Poisson element of parameter $(c-1)N+j$, while $B_{ij}^{(7)}(n)$ converges in $*$ -moments to a circular element. Therefore, the entries of the matrix $S^{(7)}(n)$, as $n \rightarrow \infty$, model the entries of the matrix x in the statement of the theorem. As $S^{(7)}(n)$ converges in $*$ -moments to a circular free Poisson element of parameter c , the theorem is proved. \square

6 Invariant Subspaces of a Circular Free Poisson Element

In this section, we will apply Theorem 5.10 and the general results of §2 to exhibit invariant subspaces for a circular free Poisson element. We will rely on the result of Haagerup and Larsen [HL, Example 5.2] that the spectrum of a circular free Poisson element of parameter c is $\{z \in \mathbf{C} \mid \sqrt{c-1} \leq |z| \leq \sqrt{c}\}$.

Theorem 6.1. *Let (\mathcal{M}, ψ) be a W^* -noncommutative probability space with ψ faithful, let $c \geq 1$ and let $y \in \mathcal{M}$ be a circular free Poisson element of parameter c . Given $r \geq 0$, let $p_r(y) \in \mathcal{M}$ be the projection onto the invariant subspace of y as in Definition 2.4. Then*

$$\psi(p_r(y)) = \begin{cases} 0 & \text{if } r \leq \sqrt{c-1} \\ r^2 - (c-1) & \text{if } \sqrt{c-1} \leq r \leq \sqrt{c} \\ 1 & \text{if } r \geq \sqrt{c}. \end{cases} \quad (66)$$

Proof. We may without loss of generality assume that $\mathcal{M} = \{y\}''$, which implies $\mathcal{M} \cong L(F_2)$ and ψ is a trace. Let $N \in \mathbf{N}$ and let

$$x = \frac{1}{\sqrt{N}} \begin{pmatrix} a_1 & b_{12} & b_{13} & \cdots & b_{1,N-1} & b_{1N} \\ 0 & a_2 & b_{23} & \cdots & b_{2,N-1} & b_{2N} \\ 0 & 0 & a_3 & \ddots & \vdots & b_{3N} \\ \vdots & \vdots & \ddots & \ddots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 & a_{N-1} & b_{N-1,N} \\ 0 & 0 & \cdots & 0 & 0 & a_N \end{pmatrix} \in M_N(A)$$

be the circular free Poisson element of parameter c as in Theorem 5.10, where we take (A, ϕ) to be a W^* -noncommutative probability space. Thus a_j is a circular free Poisson element of parameter $(c-1)N + j$, each b_{ij} is a circular element and the collection of all a_j and b_{ij} is $*$ -free. For $k \in \{1, \dots, N\}$, let

$$e_k = \text{diag}(\underbrace{1, \dots, 1}_{k \text{ times}}, 0, \dots, 0) \in M_n(\mathbf{C}1) \subseteq M_n(A).$$

Another application of Theorem 5.10 shows that in the W^* -noncommutative probability space

$$(e_k M_N(A) e_k, \sqrt{\frac{N}{k}} \phi_N \upharpoonright_{e_k M_N(A) e_k}) \cong (M_k(A), \phi_k),$$

the element $\sqrt{\frac{N}{k}} e_k x e_k$ is a circular free Poisson element of parameter $\frac{N}{k}(c-1) + 1$. Hence by [HL, Example 5.2], $e_k x e_k$ has spectrum

$$\left\{ z \in \mathbf{C} \mid \sqrt{c-1} \leq |z| \leq \sqrt{(c-1) + \frac{k}{N}} \right\}.$$

Similarly, if $k < N$ then denoting by 1_N the identity element of $M_N(A)$, we find that in the W^* -noncommutative probability space,

$$\begin{aligned} ((1_N - e_k) M_N(A) (1_N - e_k), \sqrt{\frac{N}{N-k}} \Phi_N \upharpoonright_{(1_N - e_k) M_N(A) (1_N - e_k)}) \\ \cong (M_{N-k}(A), \phi_{N-k}), \end{aligned}$$

the element $\sqrt{\frac{N}{N-k}} (1_N - e_k) x (1_N - e_k)$ is a circular free Poisson element of parameter $\frac{N}{N-k} c$. Hence $(1_N - e_k) x (1_N - e_k)$ has spectrum

$$\left\{ z \in \mathbf{C} \mid \sqrt{(c-1) + \frac{k}{N}} \leq |z| \leq \sqrt{c} \right\}.$$

Therefore, by Proposition 2.2, if $k \leq N-2$ and if

$$\sqrt{(c-1) + \frac{k}{N}} \leq r < \sqrt{(c-1) + \frac{k+1}{N}}$$

then $e_k \leq p_r(x) \leq e_{k+1}$ and consequently $\frac{k}{N} \leq \psi(p_r(y)) \leq \frac{k+1}{N}$. Letting N grow without bound and choosing k appropriately implies (66). \square

Some further facts concerning these projections $p_r(y)$ are collected below in Theorem 6.3, for the proof of which we will use the following lemma.

LEMMA 6.2. *For every $c \geq 1$ let (A_c, ϕ_c) be a W^* -noncommutative probability space and let $y_c \in A$ be a circular free Poisson element of parameter c . Given $c_0 \geq 1$ and a sequence $(c_n)_1^\infty$ in $[1, \infty)$ converging to c_0 , we have that y_{c_n} converges in $*$ -moments to y_{c_0} as $n \rightarrow \infty$.*

Proof. The positive part h_c of y_c has the same moments as the measure ρ_c on \mathbf{R}_+ whose density with respect to Lebesgue measure is

$$\frac{d\rho_c}{dt} = \frac{\sqrt{(d_1^2 - t^2)(t^2 - d_0^2)}}{\pi t} 1_{[d_0, d_1]}(t),$$

with $d_0 = 1 - \sqrt{c}$ and $d_1 = 1 + \sqrt{c}$. Since y_c has the polar decomposition $y_c = u_c h_c$ where u_c is a Haar unitary and where u_c and h_c are $*$ -free, the $*$ -moments of y_c can be expressed as certain polynomials in the moments of ρ_c . Clearly, the k th moment of ρ_{c_n} converges to the k th moment of ρ_{c_0} as $n \rightarrow \infty$. \square

Theorem 6.3. *Let y be a circular free Poisson element of parameter c in some W^* -noncommutative probability space (\mathcal{M}, ψ) , with ψ faithful. Then*

- (i) $p_s(y)$ converges to $p_r(y)$ in the strong* topology as $s \rightarrow r$.
- (ii) If $\sqrt{c-1} < r < \sqrt{c}$ then in the W^* -noncommutative probability space

$$\left(p_r(y) \mathcal{M} p_r(y), \frac{1}{\psi(p_r(y))} \psi \upharpoonright_{p_r(y) \mathcal{M} p_r(y)} \right), \quad (67)$$

$\psi(p_r(y))^{-1/2} y p_r(y)$ is a circular free Poisson element of parameter $1 + (c-1)/\psi(p_r(y))$. Hence the spectrum of $y p_r(y)$ relative to $p_r(y) \mathcal{M} p_r(y)$ is

$$\sigma(y p_r(y)) = \{z \in \mathbf{C} \mid \sqrt{c-1} \leq |z| \leq \sqrt{c}\}.$$

- (iii) If $\sqrt{c-1} < r < \sqrt{c}$, then in the W^* -noncommutative probability space

$$\left((1 - p_r(y)) \mathcal{M} (1 - p_r(y)), \frac{1}{1 - \psi(p_r(y))} \psi \upharpoonright_{(1 - p_r(y)) \mathcal{M} (1 - p_r(y))} \right),$$

$(1 - \psi(p_r(y)))^{-1/2} (1 - p_r(y)) y$ is a circular free Poisson element of parameter $c/(1 - \psi(p_r(y)))$. Hence the spectrum of $(1 - p_r(y)) y$ relative to $(1 - p_r(y)) \mathcal{M} (1 - p_r(y))$ is

$$\sigma((1 - p_r(y)) y) = \{z \in \mathbf{C} \mid \sqrt{r} \leq |z| \leq \sqrt{c}\}.$$

- (iv) If x is the upper triangular $N \times N$ matrix given in Theorem 5.10, that is a circular free Poisson element of parameter c , then for every $k \in \{0, 1, \dots, N\}$ and letting $r = \sqrt{(c-1) + k/N}$, we have $p_r(x) = \text{diag}(\underbrace{1, \dots, 1}_{k \text{ times}}, 0, \dots, 0)$.

Proof. We know from general principles that $p_{r'}(y) \leq p_r(y)$ if $r' < r$, and from Theorem 6.1 we have that $\lim_{s \rightarrow r} \psi(p_s(y)) = \psi(p_r(y))$; as ψ is faithful we conclude (i).

Let us now prove (iv). Arguing as in the proof of Theorem 6.1, we have $e_k \leq p_r(x)$ whenever $r > \sqrt{(c-1) + k/N}$. We may take the W^* -noncommutative probability space (A, ϕ) so that ϕ is a faithful trace, in which case, since $\inf\{\psi(p_r(x)) \mid r > \sqrt{(c-1) + k/N}\} = k/N$, it follows that

$$e_k = \bigwedge \left\{ p_r(x) \mid r > \sqrt{(c-1) + k/N} \right\}.$$

Thus e_k is the limit in strong* topology of $p_r(x)$ as r tends to $\sqrt{(c-1) + k/N}$ from above. Using (i), it follows that $e_k = p_{\sqrt{(c-1) + k/N}}(x)$.

For (ii), let us show that $\psi(p_r(y))^{-1/2} y p_r(y)$ is circular free Poisson of the desired parameter, first in the case when $\psi(p_r(y)) = k/N$ is rational. We may take (\mathcal{M}, ψ) to be $(M_N(A), \phi_N)$ and y to be equal to the $N \times N$ matrix x as in Theorem 5.10. By (iv), the noncommutative probability space (67) is $(M_k(A), \phi_k)$ and

$$\frac{1}{\sqrt{\psi(p_r(y))}} y p_r(y) = \frac{1}{\sqrt{k}} \begin{pmatrix} a_1 & b_{12} & b_{13} & \cdots & b_{1,k-1} & b_{1k} \\ 0 & a_2 & b_{23} & \cdots & b_{2,k-1} & b_{2k} \\ 0 & 0 & a_3 & \ddots & \vdots & b_{3k} \\ \vdots & \vdots & \ddots & \ddots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 & a_{k-1} & b_{k-1,k} \\ 0 & 0 & \cdots & 0 & 0 & a_k \end{pmatrix},$$

where a_j is circular free Poisson of parameter $(c-1)N + j$. Applying again Theorem 5.10, we obtain that $\psi(p_r(y))^{-1/2} y p_r(y)$ is circular free Poisson of parameter $1 + (c-1)/\psi(p_r(y))$. When r is such that $\psi(p_r(y))$ is irrational, then using (i) we have that $y p_r(y)$ is the strong* limit of $y p_s(y)$ as s tends to r through rational numbers. Hence by Lemma 6.2 and the continuity in r of $\psi(p_r(y))$ implied by Theorem 5.10, it follows that $\psi(p_r(y))^{-1/2} y p_r(y)$ is circular free Poisson of parameter $1 + (c-1)/\psi(p_r(y))$. The statement about the spectrum follows from the result of Haagerup and Larsen [HL] that we've been using repeatedly.

Part (iii) is proved similarly. When $\psi(p_r(y)) = k/N$ is rational then we get

$$\frac{1}{\sqrt{1 - \psi(p_r(y))}}(1 - p_r(y))y = \frac{1}{\sqrt{N - k}} \begin{pmatrix} a_{k+1} & b_{k+1,k+2} & b_{k+1,k+3} & \cdots & b_{k+1,N-1} & b_{k+1,N} \\ 0 & a_{k+2} & b_{k+2,k+3} & \cdots & b_{k+2,N-1} & b_{k+2,N} \\ 0 & 0 & a_{k+3} & \ddots & \vdots & b_{k+3,N} \\ \vdots & \vdots & \ddots & \ddots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 & a_{N-1} & b_{N-1,N} \\ 0 & 0 & \cdots & 0 & 0 & a_N \end{pmatrix},$$

where a_{k+j} is circular free Poisson of parameter $(c-1)N + k + j$. The remaining part of the argument is like for (ii) above. \square

The following proposition shows that $p_r(y)$ is characterized by the spectral conditions in (ii) and (iii) of Theorem 6.3.

PROPOSITION 6.4. *Let Y be a circular free Poisson element of parameter $c \geq 1$ in a W^* -probability space (\mathcal{M}, ψ) , with ψ faithful, and let $\sqrt{c-1} < r < \sqrt{c}$. Suppose $p \in \mathcal{M}$ is a projection such that*

- (i) $yp = pyp$
- (ii) $\sigma_{p\mathcal{M}p}(yp) \subseteq \{z \in \mathbf{C} \mid \sqrt{c-1} \leq |z| \leq r\}$
- (iii) $\sigma_{(1-p)\mathcal{M}(1-p)}((1-p)y) \subseteq \{z \in \mathbf{C} \mid r \leq |z| \leq \sqrt{c}\}$.

Then $p = p_r(y)$.

Proof. Note that (i) implies $(1-p)y = (1-p)y(1-p)$. Let \mathcal{M} be normally and faithfully represented on a Hilbert space \mathcal{H} . For $\xi \in p\mathcal{H}$ we have

$$\limsup_{n \rightarrow \infty} \|y^n \xi\|^{1/n} \leq \limsup_{n \rightarrow \infty} \|(pyp)^n\|^{1/n} \leq r,$$

where the last inequality is because the spectral radius of pyp is $\leq r$. Hence $p \leq p_r(y)$.

In order to prove the reverse inequality, it will suffice to show $p \geq p_s(y)$ for all $0 \leq s < r$, because $s \mapsto p_s(y)$ is strong*-continuous by Theorem 6.3(i). Let $0 \leq s < r$, $\xi \in (1-p)\mathcal{H}$ and let $\eta \in E_s(y)$, i.e.

$$\limsup_{n \rightarrow \infty} \|y^n \eta\|^{1/n} \leq s. \quad (68)$$

Set $\xi_n = ((1-p)y^*(1-p))^{-n}\xi$. Then $\xi_n \in (1-p)\mathcal{H}$ and, because the spectral radius of $((1-p)y^*(1-p))^{-1}$ is $\leq 1/r$, we have $\limsup_{n \rightarrow \infty} \|\xi_n\|^{1/n} \leq 1/r$. Since $(1-p)\mathcal{H}$ is an invariant subspace for y^* , we have $\xi = ((1-p)y^*(1-p))^n \xi_n = (y^*)^n \xi_n$. Therefore $\langle \xi, \eta \rangle = \langle (y^*)^n \xi_n, \eta \rangle = \langle \xi_n, y^n \eta \rangle$, so using (68)

and Schwarz's inequality we have $\limsup_{n \rightarrow \infty} |\langle \xi, \eta \rangle|^{1/n} \leq s/r < 1$, which shows that $\langle \xi, \eta \rangle = 0$. Hence $(1-p)\mathcal{H} \perp \overline{E_s(y)} = p_s(y)\mathcal{H}$ and therefore $p_s(y) \leq p$. \square

REMARK 6.5. *The proof above shows that the subspace*

$$E_r(y) = \left\{ \xi \in \mathcal{H} \mid \limsup_{k \rightarrow \infty} \|y^k \xi\|^{1/k} \leq r \right\}$$

is closed; thus we have $p_r(y)\mathcal{H} = E_r(y)$, without taking the closure.

The next example, however, shows that the sort of spectral decomposition found in Theorem 6.3 and closedness of the subspace $E_r(y)$ do not always hold.

EXAMPLE 6.6. Let $\mathcal{H} = \bigoplus_{k=2}^{\infty} \mathcal{H}_k$, where \mathcal{H}_k is k -dimensional Hilbert space with orthonormal basis $e_1^{(k)}, \dots, e_k^{(k)}$ and let $T = \bigoplus_{k=2}^{\infty} T_k$, where $T_k \in B(\mathcal{H}_k)$ is the nilpotent operator

$$T_k e_j^{(k)} = \begin{cases} 0 & j = 1 \\ e_{j-1}^{(k)} & 2 \leq j \leq k. \end{cases}$$

It is well known that the spectrum of T is the closed unit disk $\overline{\mathbf{D}}$ — see for example Brown [Br, Example 4.10]. However, if $r > 0$ then $E_r(T)$ is dense in \mathcal{H} , so $p_r(T) = 1$ and the spectrum of $T p_r(T)$ is $\overline{\mathbf{D}}$. Moreover, if $0 < r < 1$ then the vector $\sum_{k=2}^{\infty} \frac{1}{k} e_k^{(k)}$ is not an element of $E_r(T)$; this shows that $E_r(T)$ is not closed. Note that $T \in \bigoplus_{k=2}^{\infty} B(\mathcal{H}_k)$, which is a finite von Neumann algebra.

References

- [B] B.V. BRONK, Exponential ensembles for random matrices, J. Math. Phys. 6 (1965), 228–237.
- [Br] L.G. BROWN, Lidskii's Theorem in the type II case, in “Geometric Methods in Operator Algebras (Kyoto, 1983)” (H. Araki, E.G. Effros, eds.), Pitman Res. Notes Math. Ser. 123, Longman Sci. Tech. (1986), 1–35.
- [D] J. DIXMIER, Les algèbres d'opérateurs dans l'espace hilbertien, Gauthier-Villars, 1969.
- [Dy1] K.J. DYKEMA, On certain free product factors via an extended matrix model, J. Funct. Anal. 112 (1993), 31–60.
- [Dy2] K.J. DYKEMA, Free products of hyperfinite von Neumann algebras and free dimension, Duke Math. J. 69 (1993), 97–119.
- [HL] U. HAAGERUP, F. LARSEN, Brown's spectral distribution measure for R -diagonal elements in finite von Neumann algebras, preprint (1999).

- [HT] U. HAAGERUP, S. THORBJØRNSSEN, Random matrices with complex Gaussian entries, preprint (1998).
- [HeS] E. HEWITT, L.J. SAVAGE, Symmetric measures on Cartesian products, Trans. Amer. Math. Soc. 80 (1955), 470–501.
- [M] M.L. MEHTA, Random Matrices, Second edition, Academic Press, 1991.
- [NS] A. NICA, R. SPEICHER, R -diagonal pairs – a common approach to Haar unitaries and circular elements, Fields Inst. Commun. 12 (D.V. Voiculescu, ed.) (1997), 149–188.
- [R1] F. RĂDULESCU, The fundamental group of the von Neumann algebra of a free group with infinitely many generators is $\mathbf{R}_+ \setminus \{0\}$, J. Amer. Math. Soc. 5 (1992), 517–532.
- [R2] F. RĂDULESCU, Random matrices, amalgamated free products and subfactors of the von Neumann algebra of a free group, of noninteger index, Invent. Math. 115 (1994), 347–389.
- [S] E. STØRMER, Symmetric states of infinite tensor products of C^* -algebras, J. Funct. Anal. 3 (1969), 48–68.
- [V1] D. VOICULESCU, Circular and semicircular systems and free product factors, Operator Algebras, Unitary Representations, Algebras, and Invariant Theory, Progress in Mathematics 92, Birkhäuser (1990), 45–60.
- [V2] D. VOICULESCU, Limit laws for random matrices and free products, Invent. Math. 104 (1991), 201–220.
- [V3] D. VOICULESCU, A strengthened asymptotic freeness result for random matrices with applications to free entropy, Internat. Math. Res. Notices (1998), 41–63.
- [VDN] D.V. VOICULESCU, K.J. DYKEMA, A. NICA, Free Random Variables, CRM Monograph Series 1, American Mathematical Society, 1992.

KEN DYKEMA, Department of Mathematics, Texas A&M University, College Station TX 77843-3368, USA

Ken.Dykema@math.tamu.edu

<http://www.math.tamu.edu/~Ken.Dykema/>

UFFE HAAGERUP, Department of Mathematics and Computer Science, University of Southern Denmark – Odense University, Campusvej 55, 5230 Odense M, Denmark

haagerup@imada.sdu.dk

<http://www.imada.sdu.dk/~haagerup/>

Submitted: March 2000

Revision: May 2001