

isibang/ms/2008/7

May 5th, 2008

<http://www.isibang.ac.in/~statmath/eprints>

A Problem of Powers and the Product of Spatial Product Systems

B.V. RAJARAMA BHAT, VOLKMAR LIEBSCHER
and Michael Skeide

Indian Statistical Institute, Bangalore Centre
8th Mile Mysore Road, Bangalore, 560059 India

A Problem of Powers and the Product of Spatial Product Systems*

B.V. Rajarama Bhat

*Statistics and Mathematics Unit, Indian Statistical Institute Bangalore, R. V. College
Post, Bangalore 560059, India, E-mail: bhat@isibang.ac.in, Homepage:
<http://www.isibang.ac.in/Smubang/BHAT/>*

Volkmar Liebscher

*Institut für Mathematik und Informatik, Ernst-Moritz-Arndt-Universität Greifswald,
17487 Greifswald, Germany, E-mail: volkmar.liebscher@uni-greifswald.de,
Homepage:
<http://www.math-inf.uni-greifswald.de/biomathematik/liebscher/>*

Michael Skeide[†]

*Dipartimento S.E.G.e S., Università degli Studi del Molise, Via de Sanctis, 86100
Campobasso, Italy, E-mail: skeide@math.tu-cottbus.de, Homepage:
http://www.math.tu-cottbus.de/INSTITUT/lswas/_skeide.html*

In the 2002 AMS summer conference on “Advances in Quantum Dynamics” in Mount Holyoke Robert Powers proposed a sum operation for spatial E_0 -semigroups. Still during the conference Skeide showed that the Arveson system of that sum is the product of spatial Arveson systems. This product may but need not coincide with the tensor product of Arveson systems. The Powers sum of two spatial E_0 -semigroups is, therefore, up to cocycle conjugacy Skeide’s product of spatial noises.

Keywords: Quantum dynamics, quantum probability, Hilbert modules, product systems, E_0 -semigroups. 2000 AMS-Subject classification: 46L53; 46L55; 46L08; 60J25; 12H20.

*This work is supported by a PPP-project by DAAD and DST and by a Research in Pairs project at MFO.

[†]MS is supported by research funds of the University of Molise and the Italian MIUR (PRIN 2005) and by research funds of the Dipartimento S.E.G.e S. of University of Molise.

2

1. Introduction

Let $\vartheta^i = (\vartheta_t^i)_{t \in \mathbb{R}_+}$ ($i = 1, 2$) be two E_0 -semigroups on $\mathcal{B}(H)$ with associated Arveson systems $\mathfrak{H}^{i \otimes} = (\mathfrak{H}_t^i)_{t \in \mathbb{R}_+}$ (Arveson [Arv89]). Furthermore, let $\Omega^i = (\Omega_t^i)_{t \in \mathbb{R}_+} \subset \mathcal{B}(H)$ be two semigroups of intertwining isometries for ϑ^i (units). Then

$$T_t \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} = \begin{pmatrix} \vartheta_t^1(a_{11}) & \Omega_t^1 a_{12} \Omega_t^{2*} \\ \Omega_t^2 a_{12} \Omega_t^{1*} & \vartheta_t^2(a_{22}) \end{pmatrix} \quad (*)$$

defines a CP-semigroup $T = (T_t)_{t \in \mathbb{R}_+}$ on $\mathcal{B}(H \oplus H)$. In the 2002 AMS summer conference on “Advances in Quantum Dynamics” Robert Powers asked for the Arveson system associated with T (Bhat [Bha96], Arveson [Arv97]). During that conference (see [Ske03a]) Skeide showed that this product system is nothing but the *product of the spatial product systems* introduced in Skeide [Ske06] (published first in 2001). Meanwhile, Powers has formalized the above *sum operation* in [Pow04] and he has proved that the product may, but need not coincide with the tensor product of the involved Arveson systems, a fact suspected already in [Ske03a, Lie03].

In these notes we extend Powers’ construction to the case of spatial E_0 -semigroups ϑ^i on $\mathcal{B}^a(E^i)$ where E^i are Hilbert \mathcal{B} -modules. We obtain the same result as in [Ske03a], namely, the product system of the minimal dilation of the CP-semigroup on $\mathcal{B}^a(E^1 \oplus E^2)$ defined in analogy with (*) is the product of spatial product systems from [Ske06].

Like in [Ske03a], it is crucial to understand the following point (on which we will spend some time in Section 3): In Bhat and Skeide [BS00] to every CP-semigroup on a C^* -algebra \mathcal{B} a product system of \mathcal{B} -algebra has been constructed. However, the C^* -algebra in question here is $\mathcal{B}^a(E^1 \oplus E^2)$, where $\mathcal{B}^a(E)$ denotes the algebra of all adjointable operators on a Hilbert module E . So what has the product system of $\mathcal{B}^a(E^1 \oplus E^2)$ -correspondences to do with the product systems of the E_0 -semigroups ϑ^1 and ϑ^2 , which are product systems of \mathcal{B} -correspondences? The answer to this question, like in [Ske03a], will allow to construct the Arveson system of a CP-semigroup on $\mathcal{B}(H)$ *without* having to find first its minimal dilation. To understand even the Hilbert space case already requires, however, module techniques.

2. Product systems, CP-semigroups, E_0 -semigroups and dilations

Throughout these notes, by \mathcal{B} we denote a unital C^* -algebras. There are no spatial product systems where \mathcal{B} is nonunital. There is no reasonable notion

of unit for product systems of correspondences over nonunital C^* -algebras, where \mathcal{B} could not easily be substituted by a unital ideal of \mathcal{B} .

2.1 Product systems. Product systems of Hilbert modules (*product system* for short) occurred in different contexts in Bhat and Skeide [BS00], Skeide [Ske02], Muhly and Solel [MS02] and other more recent publications. Let \mathcal{B} be a unital C^* -algebra. A product system is a family $E^\odot = (E_t)_{t \in \mathbb{R}_+}$ of *correspondences* E_t over \mathcal{B} (that is, a (right) Hilbert \mathcal{B} -module with a unital representation of \mathcal{B}) with an associative identification

$$E_s \odot E_t = E_{s+t},$$

where $E_0 = \mathcal{B}$ and for $s = 0$ or $t = 0$ we get the canonical identifications. By \odot we denote the (internal) tensor product of correspondences.

If we want to emphasize that we do not put any technical condition, we say *algebraic* product system. There are concise definitions of *continuous* [Ske03b] and *measurable* (separable!) [Hir04] product systems of C^* -correspondences, and *measurable* (separable pre-dual!) [MS07] product systems of W^* -correspondences. Skeide [Ske08] will discuss *strongly continuous* product systems of von Neumann correspondences. We do not consider such constraints in these notes. We just mention for the worried reader that the result from [Ske06,Ske03b] that the product of continuous spatial product systems is continuous.

2.2 Units. A *unit* for a product system E^\odot is family $\xi^\odot = (\xi_t)_{t \in \mathbb{R}_+}$ of elements $\xi_t \in E_t$ such that

$$\xi_s \odot \xi_t = \xi_{s+t}$$

and $\xi_0 = \mathbf{1} \in \mathcal{B} = E_0$. A unit may be *unital* ($\langle \xi_t, \xi_t \rangle = \mathbf{1} \forall t \in \mathbb{R}_+$), *contractive* ($\langle \xi_t, \xi_t \rangle \leq \mathbf{1} \forall t \in \mathbb{R}_+$), or *central* ($b\xi_t = \xi_t b \forall t \in \mathbb{R}_+, b \in \mathcal{B}$).

We do not pose technical conditions on the unit. But, sufficiently continuous units can be used to pose technical conditions on the product system in a nice way; see [Ske03a].

2.3 The product system of a CP-semigroup. Let $T = (T_t)_{t \in \mathbb{R}_+}$ be a (not necessarily unital) CP-semigroup on a unital C^* -algebra \mathcal{B} . According to Bhat and Skeide [BS00] there exists a product system E^\odot with a unit ξ^\odot determined uniquely up to isomorphism (of the pair (E^\odot, ξ^\odot)) by the following properties:

- (1) $\langle \xi_t, b\xi_t \rangle = T_t(b)$.

4

(2) E^\odot is **generated** by ξ^\odot , that is, the smallest subsystem of E^\odot containing ξ^\odot is E^\odot .

In analogy with Paschke's [Pas73] GNS-construction for CP-maps, we call (E^\odot, ξ^\odot) the **GNS-system** of T and we call ξ^\odot the **cyclic unit**. In fact, $\mathcal{E}_t = \overline{\text{span}} \mathcal{B} \xi_t \mathcal{B}$ is the **GNS-module** of T_t with cyclic vector ξ_t . For the comparison of the product system of Powers' CP-semigroup with a product of product systems it is important to note that

$$\begin{aligned} E_t &= \overline{\text{span}} \{ x_{t_n}^n \odot \dots \odot x_{t_1}^1 : n \in \mathbb{N}, t_n + \dots + t_1 = t, x_{t_k}^k \in \mathcal{E}_{t_k} \} \\ &= \overline{\text{span}} \{ b_n \xi_{t_n} \odot \dots \odot b_1 \xi_{t_1} b_0 : n \in \mathbb{N}, t_n + \dots + t_1 = t, b_k \in \mathcal{B} \}. \end{aligned} \quad (2.1)$$

In fact, the product system E_t can be obtained as an inductive limit of the expressions $\mathcal{E}_{t_n} \odot \dots \odot \mathcal{E}_{t_1}$ over refinement of the partitions $t_n + \dots + t_1 = t$ of $[0, t]$.

2.4 The product system of an E_0 -semigroup on $\mathcal{B}^a(E)$. Let E be a Hilbert \mathcal{B} -module with a **unit vector** ξ (that is, $\langle \xi, \xi \rangle = \mathbf{1}$) and let $\vartheta = (\vartheta_t)_{t \in \mathbb{R}_+}$ be an E_0 -**semigroup** (that is, a semigroup of unital endomorphisms) on $\mathcal{B}^a(E)$. Let us denote by xy^* ($x, y \in E$) the **rank-one operator**

$$xy^* : z \longmapsto x \langle y, z \rangle.$$

Then $p_t := \vartheta_t(\xi \xi^*)$ is a projection and the range $E_t := p_t E$ is a Hilbert \mathcal{B} -submodule of E . By defining the (unital!) left action $b x_t = \vartheta_t(\xi b \xi^*) x_t$ we turn E_t into a \mathcal{B} -correspondence. One easily checks that

$$x \odot y_t \longmapsto \vartheta_t(x \xi^*) y_t$$

defines an isometry $u_t : E \odot E_t \rightarrow E$. Clearly, if ϑ_t is **strict** (that is, precisely, if $\overline{\text{span}} \vartheta_t(E E^*) E = E$), then u_t is a unitary. Identifying $E = E \odot E_t$ and using the semigroup property, we find

$$\vartheta_t(a) = a \odot \text{id}_{E_t} \quad (E \odot E_s) \odot E_t = E \odot (E_s \odot E_t). \quad (2.2)$$

The restriction of u_t to $E_s \odot E_t$ is a bilinear unitary onto E_{s+t} and the preceding associativity reads now $(E_r \odot E_s) \odot E_t = E_r \odot (E_s \odot E_t)$. Obviously, $E_0 = \mathcal{B}$ and the identifications $E_t \odot E_0 = E_t = E_0 \odot E_t$ are the canonical ones. Thus, $E^\odot = (E_t)_{t \in \mathbb{R}_+}$ is a product system.

For E_0 -semigroups on $\mathcal{B}(H)$ the preceding construction is due to Bhat [Bha96], the extension to Hilbert modules to Skeide [Ske02]. We would like to mention that Bhat's construction does not give the Arveson system of an E_0 -semigroup, but its *opposite* Arveson system (all orders in tensor

products reversed). By Tsirelson [Tsi00] the two need not be isomorphic. For Hilbert C^* -modules Arveson's construction does not work. For von Neumann modules it works, but gives a product system of von Neumann correspondences over \mathcal{B}' , the commutant of \mathcal{B} ; see [Ske03a,Ske04].

Existence of a unit vector is not a too hard requirement, as long as \mathcal{B} is unital. (If E has no unit vector, then a finite multiple E^n will have one; see [Ske04]. And product systems do not change under taking direct sums.) We would like to mention a further method to construct the product system of an E_0 -semigroup, that works also for nonunital \mathcal{B} . It relies on the representations theory of $\mathcal{B}^a(E)$ in Muhly, Skeide and Solel [MSS06]. See [Ske04] for details.

2.5 Dilation and minimal dilation. Suppose E^\odot is a product system with a unit ξ^\odot . Clearly, $T_t := \langle \xi_t, \bullet \xi_t \rangle$ defines a CP-semigroup $T = (T_t)_{t \in \mathbb{R}_+}$, which is unital, if and only if ξ^\odot is unital. Obviously, E^\odot is the product systems of T , if and only if it is generated by ξ^\odot .

If ξ^\odot is unital, then we may embed E_t as $\xi_s \odot E_t$ into E_{s+t} . This gives rise to an inductive limit E and a factorization $E = E \odot E_t$, fulfilling the associativity condition in (2.2). It follows that $\vartheta_t(a) = a \odot \text{id}_{E_t}$ defines an E_0 -semigroup $\vartheta = (\vartheta_t)_{t \in \mathbb{R}_+}$ on $\mathcal{B}^a(E)$. The embedding $E_t \rightarrow E_{s+t}$ is, in general, only right linear so that, in general, E is only a right Hilbert module.

Under the inductive limit all $\xi_t \in E_t \subset E$ correspond to the same unit vector $\xi \in E$. Moreover, $\xi = \xi \odot \xi_t$, so that the vector expectation $\varphi := \langle \xi, \bullet \xi \rangle$ fulfills $\varphi \circ \vartheta_t(\xi b \xi^*) = T_t(b)$, that is, (E, ϑ, ξ) is a **weak dilation** of T in the sense of [BP94,BS00]. Clearly, the product system of ϑ (constructed with the unit vector ξ) is E^\odot .

Suppose ϑ is a strict E_0 -semigroup on some $\mathcal{B}^a(E)$ and that ξ is a unit vector in E . One may show (see [Ske02]) that $T_t(b) := \varphi \circ \vartheta_t(\xi b \xi^*)$ defines a (necessarily unital) CP-semigroup (which it dilates), if and only if the projections $p_t := \vartheta_t(\xi \xi^*)$ increase. In this case, the product system E^\odot of ϑ has a unit $\xi^\odot = (\xi_t)_{t \in \mathbb{R}_+}$ with $\xi_t := p_t \xi$, which fulfills $T_t = \langle \xi_t, \bullet \xi_t \rangle$. We say the weak dilation (E, ϑ, ξ) of T is **minimal**, if the **flow** $j_t(b) := \vartheta_t(\xi b \xi^*)$ generates E out of ξ . One may show that this is the case, if and only if the product system of ϑ coincides with the product system of T . The minimal (weak) dilation is determined up to suitable unitary equivalence.

2.6 Remark. We would like to emphasize that in order to construct the minimal dilation of a unital CP-semigroup T , we first constructed the prod-

uct system of T and then constructed the dilating E_0 -semigroup ϑ (giving back the product system of T). It is not necessary to pass through minimal dilation to obtain the product system of T , but rather the other way round.

2.7 Spatial product systems. Following [Ske06], we call a product system E^\odot *spatial*, if it has central unital *reference unit* $\omega^\odot = (\omega_t)_{t \in \mathbb{R}_+}$. The choice of the reference unit is part of the spatial structure, so we will write a pair (E^\odot, ω^\odot) . For instance, a *morphism* $w^\odot: E^\odot \rightarrow F^\odot$ between product systems E^\odot and F^\odot is a family $w^\odot = (w_t)_{t \in \mathbb{R}_+}$ of mappings $w_t \in \mathcal{B}^{a,bil}(E_t, F_t)$ (that is, bilinear adjointable mappings from E_t to F_t) fulfilling $w_s \odot w_t = w_{s+t}$ and $w_0 = \text{id}_{\mathcal{B}}$. To be a *spatial* morphism of spatial product systems, w^\odot must send the reference unit of E^\odot to the reference unit of F^\odot .

Our definition matches that of Powers [Pow87] in that an E_0 -semigroup ϑ on $\mathcal{B}^a(E)$ admits a so-called intertwining semigroup of isometries, if and only if the product system of ϑ is spatial. It does not match the usual definition for Arveson systems, where an Arveson system is *spatial*, if it has a unit. The principle result of Barreto, Bhat, Liebscher and Skeide [BBS04] asserts that a product system of von Neumann correspondences is spatial, if it has a (continuous) unit. But, for Hilbert modules this statement fails. In fact, we show in [BLS08a] that, unlike for Arveson systems, a subsystem of a product system of Fock modules need not be spatial (in particular, it need not be Fock).

There are many interesting questions about spatial product systems, open even in the Hilbert space case. Does the spatial structure of the spatial product system depend on the choice of the reference unit? The equivalent question is, whether every spatial product system is *amenable* [Bha01] in the sense that the product system automorphisms act transitively on the set of units. Tsirelson [Tsi04] claims they are not. But, still there is a gap that has not yet been filled. In contrast to this, the question raised by Powers [Pow04], whether the product defined in the next section depends on the reference units, or not, we can answer in the negative sense; see [BLS08b].

2.8 The product of spatial product systems. The basic motivation of [Ske06] was to define an *index* of a product system and to find a *product* of product systems under which the index is *additive*. Both problems could not be solved in full generality, but precisely for the category of spatial product systems.

The mentioned result [BLS08a] is one of the reasons why it is hopeless

to define an index for nonspatial product systems. However, once accepted the necessity to restrict to spatial product systems (anyway, the index of a nonspatial Arveson systems is somewhat an artificial definition), everything works as we know it from Arveson systems, provided we indicate the good product operation.

In the theory of Arveson systems, there is the tensor product (of arbitrary Arveson systems). However, for modules this does not work. (You may write down the tensor product of correspondences, but, in general, it is not possible to define a product system structure.)

The **product** of two spatial product systems $E^{i\odot}$ ($i = 1, 2$) with reference units $\omega^{i\odot}$ is the spatial product system $(E^1 \odot E^2)^{\odot}$ with reference unit ω^{\odot} which is characterized uniquely up to spatial isomorphism by the following properties:

- (1) There are spatial isomorphisms $w^{i\odot}$ from $E^{i\odot}$ onto subsystems of $(E^1 \odot E^2)^{\odot}$.
- (2) $(E^1 \odot E^2)^{\odot}$ is generated by these two subsystems.
- (3) $\langle w_t^1(x_t^1), w_t^2(y_t^2) \rangle = \langle x_t^1, \omega_t^1 \rangle \langle \omega_t^2, y_t^2 \rangle$.

Existence of the product follows by an inductive limit; see [Ske06]. By Condition 1 we may and, usually, will identify the factors as subsystem of the product. Condition 3 means, roughly speaking, that the reference units of the two factors are identified, while components from different factors which are orthogonal to the respective reference unit are orthogonal in the product. Condition 2 means that

$$\begin{aligned} & (E^1 \odot E^2)_t \\ &= \overline{\text{span}}\{x_{t_n}^n \odot \dots \odot x_{t_1}^1 : n \in \mathbb{N}, t_n + \dots + t_1 = t, x_{t_k}^k \in E_{t_k}^i \ (i = 1, 2)\}. \end{aligned}$$

It is important to note (crucial exercise!) that this may be rewritten in the form

$$\begin{aligned} & (E^1 \odot E^2)_t \\ &= \overline{\text{span}}\{x_{t_n}^n \odot \dots \odot x_{t_1}^1 : n \in \mathbb{N}, t_n + \dots + t_1 = t, x_{t_k}^k \in \mathcal{E}_{t_k}^i \ (i = 1, 2)\}, \end{aligned} \tag{2.3}$$

where we put $\mathcal{E}_t := \mathcal{B}\omega_t \oplus (E_t^1 \ominus \mathcal{B}\omega_t^1) \oplus (E_t^2 \ominus \mathcal{B}\omega_t^2)$ (the direct sum of E^1 and E^2 with “identification of the reference vectors” and denoting the new reference vector by ω_t). Written in that way, it is easy to see that the subspaces are actually increasing of the partitions $t_n + \dots + t_1 = t$ of $[0, t]$. This gives an idea how to obtain the product as an inductive limit; see [Ske06].

3. The product system of \mathcal{B} -correspondences of a CP-semigroup on $\mathcal{B}^a(E)$

In Section 2.3 we have said what the product system of CP-semigroup on \mathcal{B} is. It is a product systems of \mathcal{B} -correspondences. On the other hand, if $\mathcal{B}(H)$ -people speak about the product system of a unital CP-semigroup on $\mathcal{B}(H)$, they mean an Arveson system, that is, a product system of Hilbert spaces. Following Bhat [Bha96] and Arveson [Arv97], the Arveson system of a unital CP-semigroup is the Arveson system of its minimal dilating E_0 -semigroup. (To be specific, we mean the product system constructed as in Section 2.4 following [Bha96], not the product system constructed in [Arv89], which is anti-isomorphic to the former.) A precise understanding of the relation between the two product systems, one of $\mathcal{B}(H)$ -modules, the other of Hilbert spaces, will allow to avoid the construction of the minimal dilation. But we will discuss it immediately for CP-semigroups on $\mathcal{B}^a(E)$.

Suppose we have a Hilbert $\mathcal{B}^a(E)$ -module F . Then we may define the Hilbert \mathcal{B} -module $F \odot E$. Every $y \in F$ gives rise to a mapping $y \odot \text{id} \in \mathcal{B}^a(E, F \odot E)$ defined by $(y \odot \text{id}_E)x = y \odot x$ with adjoint $y^* \odot \text{id}_E: y' \odot x \mapsto \langle y, y' \rangle x$. These mappings fulfill $(y \odot \text{id}_E)^*(y' \odot \text{id}_E) = \langle y, y' \rangle$ and $ya \odot \text{id}_E = (y \odot \text{id}_E)a$ for every $a \in \mathcal{B}^a(E)$. Via $a \mapsto y \odot \text{id}_E$ we may identify F as a subset of $\mathcal{B}^a(E, F \odot E)$. This subset is strictly dense but, in general, it need not coincide. In fact, we have always $F \supset \mathcal{K}(E, F \odot E)$ where the *compact operators* between Hilbert \mathcal{B} -modules E_1 and E_2 are defined as $\mathcal{K}(E_1, E_2) := \overline{\text{span}}\{x_1 x_2^* : x_i \in E_i\}$, and $F = \mathcal{K}(E, F \odot E)$ whenever the right multiplication is strict (in the same sense as left multiplication, namely, $\overline{\text{span}} F E E^* = F$).

3.1 Remark. The space $\mathcal{B}^a(E, F \odot E)$ may be thought of as the strict completion of F , and it is possible to define a strict tensor product of $\mathcal{B}^a(E)$ -correspondences. We do not need this here, and refer the interested reader to [Ske04].

Now suppose that F is a $\mathcal{B}^a(E)$ -correspondence with strict left action. If E has a unit vector, then, doing as in Section 2.4, we see that F factors into $E \odot F_E$ (where the F_E is a suitable multiplicity correspondence from \mathcal{B} to $\mathcal{B}^a(E)$) and $a \in \mathcal{B}^a(E)$ acts on $F = E \odot F_E$ as $a \odot \text{id}_{F_E}$. For several reasons we do not follow Section 2.4, but refer to the representation theory of $\mathcal{B}^a(E)$ from [MSS06]. This representation theory tells us that F_E may be chosen as $E^* \odot F$, where E^* is the *dual* \mathcal{B} - $\mathcal{B}^a(E)$ -correspondence of E

with operations $\langle x^*, x'^* \rangle := xx'^*$ and $bx^*a := (a^*xb^*)^*$. Then, clearly,

$$\begin{aligned} F &= \overline{\text{span}} \mathcal{K}(E)F = \mathcal{K}(E) \odot F \\ &= (E \odot E^*) \odot F = E \odot (E^* \odot F) = E \odot F_E \end{aligned}$$

explains both how the isomorphism is to be defined and what the action of a is. Putting this together with the preceding construction, we obtain

$$\mathcal{B}^a(E, E \odot E_F) \supset F \supset \mathcal{K}(E, E \odot E_F) = E \odot E_F \odot E^*,$$

where we defined the \mathcal{B} -correspondence $E_F := E^* \odot F \odot E$.

3.2 Remark. We do not necessarily have equality $F = \mathcal{K}(E, E \odot E_F)$. But if we have (so that F is a *full* Hilbert $\mathcal{K}(E)$ -module), then the operation of *tensor conjugation* with E^* may be viewed as an operation of Morita equivalence for correspondences in the sense of Muhly and Solel [MS00]. In what follows, the generalization to Morita equivalence of product systems [Ske04] is in the background. An elaborate version for the strict tensor product (see Remark 3.1) can be found in [Ske04].

We observe that the assignment (the functor, actually) $F \mapsto E_F := E^* \odot F \odot E$ respects tensor products. Indeed, if F_1 and F_2 are $\mathcal{B}^a(E)$ -correspondences with strict left actions, then

$$\begin{aligned} E_{F_1} \odot E_{F_2} &= (E^* \odot F_{F_1} \odot E) \odot (E^* \odot F_{F_2} \odot E) \\ &= E^* \odot F_{F_1} \odot (E \odot E^* \odot F_{F_2}) \odot E \\ &= E^* \odot F_{F_1} \odot F_{F_2} \odot E = E_{F_1 \odot F_2}. \end{aligned} \quad (3.1)$$

It is, clearly, associative. It respects inclusions and, therefore, inductive limits. If E is full, then $E^* \odot \mathcal{B}^a(E) \odot E = \mathcal{B}$. We summarize:

3.3 Proposition [Ske04]. *Suppose that E is full (for instance, E has a unit vector). Suppose that $F^\odot = (F_t)_{t \in \mathbb{R}_+}$ is a product system of $\mathcal{B}^a(E)$ -correspondences such that the left actions of all F_t are strict.*

Then the family $E^\odot = (E_t)_{t \in \mathbb{R}_+}$ of \mathcal{B} -correspondences $E_t := E^ \odot F_t \odot E$ with product system structure defined by (3.1) is a product system.*

Moreover, if the F_t are inductive limits over families \mathcal{F}_t , then the E_t are inductive limits over the corresponding $\mathcal{E}_t := E^ \odot \mathcal{F}_t \odot E$.*

3.4 Theorem. *Let F^\odot be the GNS-system of a strict unital CP-semigroup T on $\mathcal{B}^a(E)$, and denote by (F, θ, ζ) the minimal dilation of T .*

10

Then E^\odot (from Proposition 3.3) is the product system of the strict E_0 -semigroup ϑ induced on $\mathcal{B}^a(F \odot E) \cong \mathcal{B}^a(F) \odot \text{id}_E = \mathcal{B}^a(F)$ by θ .

The triple $(F \odot E, \vartheta, p = \vartheta_0(\zeta\zeta^*))$ is the unique minimal dilation of T to the operators on a Hilbert \mathcal{B} -module in the sense that

$$p(F \odot E) = \mathbf{1}_{\mathcal{B}^a(E)} \odot E = E$$

and

$$p\vartheta_t(a)p = T_t(a).$$

Proof. We proceed precisely as in the proof of [Ske04, Theorem 5.12]. We know (see Section 2.5) that the product system of the minimal θ is F^\odot . Though, we have a unit vector ζ in F , it is more suggestive to think of the correspondences F_t to be obtained as $F_t = F^* \odot {}_tF$ where ${}_tF$ is F viewed as $\mathcal{B}^a(E)$ -correspondences via θ_t ; see [Ske04, Section 2] for details. In the same way, the product system of ϑ is $(F \odot E)^* \odot {}_t(F \odot E)$. We find

$$\begin{aligned} (F \odot E)^* \odot {}_t(F \odot E) &= (E^* \odot F^*) \odot ({}_tF \odot E) \\ &= E^* \odot (F^* \odot {}_tF) \odot E = E^* \odot F_t \odot E = E_t. \end{aligned}$$

(Note: The first step where ${}_t$ goes from outside the brackets into, is just the definition of ϑ_t .) This shows the first statement.

For the second statement, we observe that $x \mapsto \zeta \odot x$ provides an isometric embedding of E into $F \odot E$ and that p is the projection on the range $\zeta \odot E$ of this embedding. Clearly,

$$\begin{aligned} p\vartheta_t(a)p &= (\zeta\zeta^* \odot \text{id}_E)(\theta_t(a) \odot \text{id}_E)(\zeta\zeta^* \odot \text{id}_E) \\ &= \left(\zeta \langle \zeta, \theta_t(a)\zeta \rangle \zeta^* \right) \odot \text{id}_E = (\zeta T_t(a)\zeta^*) \odot \text{id}_E = T_t(a), \end{aligned}$$

when $\mathcal{B}^a(E)$ is identified with the corner $(\zeta\mathcal{B}^a(E)\zeta^*) \odot \text{id}_E$ in $\mathcal{B}^a(F \odot E)$. ■

3.5 Remark. If T is a normal unital CP-semigroup on $\mathcal{B}(H)$ (normal CP-maps on $\mathcal{B}(H)$ are strict), then E^\odot is nothing but the Arveson system of T (in the sense of Bhat's construction). Note that we did construct E^\odot **without** constructing the minimal dilation first. In the theorem the minimal dilation occurred only, because we wanted to verify that our product system coincides with the one constructed via minimal dilation.

3.6 Remark. We hope that the whole discussion could help to clarify the discrepancy between the terminology and constructions in the case of CP-semigroups on $\mathcal{B}(H)$ and those for CP-semigroups on \mathcal{B} . The semigroups

of this section lie in between, in that they are CP-semigroups on $\mathcal{B}^a(E)$, so not general \mathcal{B} but also not just $\mathcal{B}(H)$. The operation that transforms the product system of $\mathcal{B}^a(E)$ -correspondences into a product system of \mathcal{B} -correspondences is *cum grano salis* an operation of Morita equivalence. (In the von Neumann case and when E is full, it is Morita equivalence.) We obtain \mathcal{B} -correspondences because E is a Hilbert \mathcal{B} -module. For $\mathcal{B}(H)$ we obtain \mathbb{C} -correspondences (or Hilbert spaces), because H is a Hilbert \mathbb{C} -module.

4. Powers' CP-semigroup

We, finally, come to Powers' CP-semigroup and to the generalization to Hilbert modules of the result from [Ske03a] that its product system is the product of the involves spatial product systems.

Let ϑ^i ($i = 1, 2$) be two strict E_0 -semigroup on $\mathcal{B}^a(E^i)$ (E^i two Hilbert \mathcal{B} -modules with unit vectors ω^i) with spatial product systems $E^{i\odot}$ (as in Section 2.4) and unital central reference units $\omega^{i\odot}$. Since ω_t^i commutes with \mathcal{B} , the mapping $b \mapsto \omega_t^i b$ is bilinear. Consequently, $\Omega_t^i := \text{id}_{E^i} \odot \omega_t^i: x^i \mapsto x^i \odot \omega_t^i \in E^i \odot E_t^i = E^i$ defines a semigroup of isometries in $\mathcal{B}^a(E^i)$. (The isometries are *intertwining* in the sense that $\vartheta_t^i(a)\Omega_t^i = (a \odot \text{id}_{E_t^i})(\text{id}_{E^i} \odot \omega_t^i) = (\text{id}_{E^i} \odot \omega_t^i)a = \Omega_t^i a$.) It follows that

$$T_t \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} = \begin{pmatrix} \vartheta_t^1(a_{11}) & \Omega_t^1 a_{12} \Omega_t^{2*} \\ \Omega_t^2 a_{21} \Omega_t^{1*} & \vartheta_t^2(a_{22}) \end{pmatrix}$$

defines a unital semigroup on $\mathcal{B}^a\left(\begin{smallmatrix} E^1 \\ E^2 \end{smallmatrix}\right)$. (We see later on that T_t is completely positive, by giving its GNS-module explicitly.) Using the identifications $E^i = E^i \odot E_t^i$ we find the more convenient form

$$T_t \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} = \begin{pmatrix} a_{11} \odot \text{id}_{E_t^1} & (\text{id}_{E^1} \odot \omega_t^1) a_{12} (\text{id}_{E^2} \odot \omega_t^{2*}) \\ (\text{id}_{E^2} \odot \omega_t^2) a_{21} (\text{id}_{E^1} \odot \omega_t^{1*}) & a_{22} \odot \text{id}_{E_t^2} \end{pmatrix}$$

where T_t maps from $\mathcal{B}^a\left(\begin{smallmatrix} E^1 \\ E^2 \end{smallmatrix}\right)$ to $\mathcal{B}^a\left(\begin{smallmatrix} E^1 \odot E_t^1 \\ E^2 \odot E_t^2 \end{smallmatrix}\right) = \mathcal{B}^a\left(\begin{smallmatrix} E^1 \\ E^2 \end{smallmatrix}\right)$.

Denote by F^\odot the product system of T in the sense of Section 2.4, that is, the F_t are $\mathcal{B}^a\left(\begin{smallmatrix} E^1 \\ E^2 \end{smallmatrix}\right)$ -correspondences. By Proposition 3.3, setting $E_t := \left(\begin{smallmatrix} E^1 \\ E^2 \end{smallmatrix}\right)^* \odot F_t \odot \left(\begin{smallmatrix} E^1 \\ E^2 \end{smallmatrix}\right)$ we define a product system E^\odot of \mathcal{B} -correspondences and

$$\mathcal{K}\left(\left(\begin{smallmatrix} E^1 \\ E^2 \end{smallmatrix}\right), \left(\begin{smallmatrix} E^1 \\ E^2 \end{smallmatrix}\right) \odot E_t\right) \subset F_t \subset \mathcal{B}^a\left(\left(\begin{smallmatrix} E^1 \\ E^2 \end{smallmatrix}\right), \left(\begin{smallmatrix} E^1 \\ E^2 \end{smallmatrix}\right) \odot E_t\right).$$

4.1 Theorem. E^\odot is the product $(E^1 \odot E^2)^\odot$ of the spatial product systems $E^{1\odot}$ and $E^{2\odot}$.

12

Proof. Recall that, by (2.1), F_t is the inductive limit of expressions of the form

$$\mathcal{F}_t := \mathcal{F}_{t_n} \odot \dots \odot \mathcal{F}_{t_1}$$

over the partitions $\mathbf{t} = (t_n, \dots, t_1)$ with $t_n + \dots + t_1 = t$, where \mathcal{F}_t is the GNS-module of T_t with cyclic vector ζ_t .

Put $\mathcal{E}_t = \begin{pmatrix} E^1 \\ E^2 \end{pmatrix}^* \odot \mathcal{F}_t \odot \begin{pmatrix} E^1 \\ E^2 \end{pmatrix}$. Then $\mathcal{F}_t \subset \mathcal{B}^a \left(\begin{pmatrix} E^1 \\ E^2 \end{pmatrix}, \begin{pmatrix} E^1 \\ E^2 \end{pmatrix} \odot \mathcal{E}_t \right)$. We claim that $\mathcal{E}_t = \mathcal{B}\omega_t \oplus (E_t^1 \ominus \mathcal{B}\omega_t^1) \oplus (E_t^2 \ominus \mathcal{B}\omega_t^2)$ and that ζ_t is the operator given by

$$\zeta_t \begin{pmatrix} x^1 \\ x^2 \end{pmatrix} = \begin{pmatrix} z^1 \\ 0 \end{pmatrix} \odot (\langle \omega_t^1, y_t^1 \rangle, p_t^1 y_t^1, 0) + \begin{pmatrix} 0 \\ z^2 \end{pmatrix} \odot (\langle \omega_t^2, y_t^2 \rangle, 0, p_t^2 y_t^2),$$

with $\begin{pmatrix} x^1 \\ x^2 \end{pmatrix} = \begin{pmatrix} z^1 \odot y_t^1 \\ z^2 \odot y_t^2 \end{pmatrix} \in \begin{pmatrix} E^1 \\ E^2 \end{pmatrix} = \begin{pmatrix} E^1 \odot E_t^1 \\ E^2 \odot E_t^2 \end{pmatrix}$ and $p_t^i := \text{id}_{E_t^i} - \omega_t^i \omega_t^{i*}$. To show this, we must check two things. Firstly, we must check whether $\langle \zeta_t, a\zeta_t \rangle = T_t(a)$. This is straightforward and we leave it as an exercise. Secondly, we must check whether elements of the form on the right-hand side of

$$\begin{pmatrix} x^1 \\ x^2 \end{pmatrix}^* \odot \zeta_t \odot \begin{pmatrix} z^1 \odot y_t^1 \\ z^2 \odot y_t^2 \end{pmatrix} \longmapsto \left(\begin{pmatrix} x^1 \\ x^2 \end{pmatrix}^* \odot \text{id}_{\mathcal{E}_t} \right) \zeta_t \begin{pmatrix} z^1 \odot y_t^1 \\ z^2 \odot y_t^2 \end{pmatrix}$$

are total in \mathcal{E}_t . For the right-hand side we find

$$\begin{aligned} & \left\langle \begin{pmatrix} x^1 \\ x^2 \end{pmatrix}, \begin{pmatrix} z^1 \\ 0 \end{pmatrix} \right\rangle \langle \omega_t^1, y_t^1 \rangle, p_t^1 y_t^1, 0 \rangle + \left\langle \begin{pmatrix} x^1 \\ x^2 \end{pmatrix}, \begin{pmatrix} 0 \\ z^2 \end{pmatrix} \right\rangle \langle \omega_t^2, y_t^2 \rangle, 0, p_t^2 y_t^2 \rangle \\ & = \langle x_1, z_1 \rangle \langle \omega_t^1, y_t^1 \rangle + \langle x_2, z_2 \rangle \langle \omega_t^2, y_t^2 \rangle, \langle x_1, z_1 \rangle p_t^1 y_t^1, \langle x_2, z_2 \rangle p_t^2 y_t^2 \rangle. \end{aligned}$$

From this, totality follows.

By Proposition 3.3, we obtain E_t as inductive limit over the expressions

$$\mathcal{E}_t := \mathcal{E}_{t_n} \odot \dots \odot \mathcal{E}_{t_1},$$

which, by the preceding computation, precisely coincides with what is needed, according to (2.3), to obtain $E_t = (E^1 \odot E^2)_t$. ■

References

- Arv89. W. Arveson. *Continuous analogues of Fock space*. Number 409 in Mem. Amer. Math. Soc. American Mathematical Society, 1989.
- Arv97. W. Arveson. Minimal E_0 -semigroups. In P. Fillmore and J. Mingo, editors, *Operator algebras and their applications*, number 13 in Fields Inst. Commun., pages 1–12. American Mathematical Society, 1997.
- BBLS04. S.D. Barreto, B.V.R. Bhat, V. Liebscher, and M. Skeide. Type I product systems of Hilbert modules. *J. Funct. Anal.*, 212:121–181, 2004. (Preprint, Cottbus 2001).

- Bha96. B.V.R. Bhat. An index theory for quantum dynamical semigroups. *Trans. Amer. Math. Soc.*, 348:561–583, 1996.
- Bha01. B.V.R. Bhat. *Cocycles of CCR-flows*. Number 709 in Mem. Amer. Math. Soc. American Mathematical Society, 2001.
- BLS08a. B.V.R. Bhat, V. Liebscher, and M. Skeide. Subsystems of Fock need not be Fock. in preparation, 2008.
- BLS08b. B.V.R. Bhat, V. Liebscher, and M. Skeide. The product of Arveson systems does not depend on the reference units. in preparation, 2008.
- BP94. B.V.R. Bhat and K.R. Parthasarathy. Kolmogorov’s existence theorem for Markov processes in C^* -algebras. *Proc. Indian Acad. Sci. (Math. Sci.)*, 104:253–262, 1994.
- BS00. B.V.R. Bhat and M. Skeide. Tensor product systems of Hilbert modules and dilations of completely positive semigroups. *Infin. Dimens. Anal. Quantum Probab. Relat. Top.*, 3:519–575, 2000. (Rome, Volterra-Preprint 1999/0370).
- Hir04. I. Hirshberg. C^* -Algebras of Hilbert module product systems. *J. Reine Angew. Math.*, 570:131–142, 2004.
- Lie03. V. Liebscher. Random sets and invariants for (type II) continuous tensor product systems of Hilbert spaces. Preprint, arXiv: math.PR/0306365, 2003. To appear in Mem. Amer. Math. Soc.
- MS00. P.S. Muhly and B. Solel. On the Morita equivalence of tensor algebras. *Proc. London Math. Soc.*, 81:113–168, 2000.
- MS02. P.S. Muhly and B. Solel. Quantum Markov processes (correspondences and dilations). *Int. J. Math.*, 51:863–906, 2002. (arXiv: math.OA/0203193).
- MS07. P.S. Muhly and B. Solel. Quantum Markov semigroups (product systems and subordination). *Int. J. Math.*, 18:633–669, 2007. (arXiv: math.OA/0510653).
- MSS06. P.S. Muhly, M. Skeide, and B. Solel. Representations of $\mathcal{B}^a(E)$. *Infin. Dimens. Anal. Quantum Probab. Relat. Top.*, 9:47–66, 2006. (arXiv: math.OA/0410607).
- Pas73. W.L. Paschke. Inner product modules over B^* -algebras. *Trans. Amer. Math. Soc.*, 182:443–468, 1973.
- Pow87. R.T. Powers. A non-spatial continuous semigroup of $*$ -endomorphisms of $\mathcal{B}(\mathfrak{H})$. *Publ. Res. Inst. Math. Sci.*, 23:1053–1069, 1987.
- Pow04. R.T. Powers. Addition of spatial E_0 -semigroups. In *Operator algebras, quantization, and noncommutative geometry*, number 365 in Contemporary Mathematics, pages 281–298. American Mathematical Society, 2004.
- Ske02. M. Skeide. Dilations, product systems and weak dilations. *Math. Notes*, 71:914–923, 2002.
- Ske03a. M. Skeide. Commutants of von Neumann modules, representations of $\mathcal{B}^a(E)$ and other topics related to product systems of Hilbert modules. In G.L. Price, B .M. Baker, P.E.T. Jorgensen, and P.S. Muhly, editors, *Advances in quantum dynamics*, number 335 in Contemporary Mathematics, pages 253–262. American Mathematical Society, 2003.

- (Preprint, Cottbus 2002, arXiv: math.OA/0308231).
- Ske03b. M. Skeide. Dilation theory and continuous tensor product systems of Hilbert modules. In W. Freudenberg, editor, *Quantum Probability and Infinite Dimensional Analysis*, number XV in Quantum Probability and White Noise Analysis, pages 215–242. World Scientific, 2003. Preprint, Cottbus 2001.
- Ske04. M. Skeide. Unit vectors, Morita equivalence and endomorphisms. Preprint, arXiv: math.OA/0412231v5 (Version 5), 2004.
- Ske06. M. Skeide. The index of (white) noises and their product systems. *Infin. Dimens. Anal. Quantum Probab. Relat. Top.*, 9:617–655, 2006. (Rome, Volterra-Preprint 2001/0458, arXiv: math.OA/0601228).
- Ske08. M. Skeide. Dilations of product systems and commutants of von Neumann modules. in preparation, 2008.
- Tsi00. B. Tsirelson. From random sets to continuous tensor products: answers to three questions of W. Arveson. Preprint, arXiv: math.FA/0001070, 2000.
- Tsi04. B. Tsirelson. On automorphisms of type II Arveson systems (probabilistic approach). Preprint, arXiv: math.OA/0411062, 2004.