

write a program to search for solutions, but that won't solve the problem in general, and so it seems that solving this problem for $m > 3$ will require some other approach.

Summary. When is the average of sums of powers of integers itself a sum of the first n integers raised to a power? We provide all solutions when averaging two sums, and provide some conditions regarding when larger averages may have solutions.

References

1. S. Kahan, An Anything-But-Average Average, *Journal of Recreational Math.* **31** (2002–2003) 253–256.
2. G. Mackiw, A Combinatorial Approach to Sums of Integer Powers, *Math. Mag.* **73** (2000) 44–46; available at <http://dx.doi.org/10.2307/2691490>

Uncountably Generated Ideals of Functions

B. Sury (sury@isibang.ac.in), Stat-Math Unit, Indian Statistical Institute, 8th Mile Mysore Road, Bangalore, India

Undergraduates usually think that the study of continuous functions and the study of abstract algebra are divorced from each other. More often than not, they find it very surprising that concepts like rings and ideals could be applied to function spaces as well! Some applications in algebra texts concern the ring $C[0, 1]$ of real-valued continuous functions on $[0, 1]$; however, these texts restrict themselves to a few standard exercises although more could be accomplished with almost the same amount of labor. For instance, the exercises in [1, p. 388], [2, p. 259], and [3, p. 140] ask for a proof that maximal ideals in $C[0, 1]$ are not finitely generated. The fact that these maximal ideals are not *countably* generated does not seem to be as well-known as it should be although the proof is not harder! We will prove this, and then use it to produce some non-prime ideals in $C(0, 1)$ which cannot be countably generated as well. Without further ado, let us begin.

Maximal ideals in $C[0, 1]$ Let $I_c = \{f \in C[0, 1] : f(c) = 0\}$ where $c \in [0, 1]$. Contrary to what we want to prove, assume that I_c is generated by a countable set $\{f_1, f_2, \dots\}$. By re-scaling, we may assume that $|f_n(x)| \leq 1$ for all x and for all n . Consider the function

$$f(x) := \sum_{n=1}^{\infty} \sqrt{\frac{|f_n(x)|}{2^n}}.$$

By uniform convergence, f is continuous. Clearly, $f \in I_c$. By assumption $f = \sum_{i=1}^r g_i f_i$ for suitable g_i in $C[0, 1]$ and natural number r .

Let M be an upper bound for $|g_i|$ for all $i \leq r$ and all x in $[0, 1]$. Then,

$$|f(x)| \leq M \sum_{i=1}^r |f_i(x)|.$$

<http://dx.doi.org/10.4169/college.math.j.42.5.404>
MSC: 13A15, 13E15, 46J20

Now, by continuity, there is a neighborhood U of c such that

$$\sqrt{|f_i(x)|} < \frac{1}{2^i M}$$

for $x \in U$ and $i \leq r$. In addition, since f vanishes only at c , for each $x \in U$, $x \neq c$, $f_i(x) \neq 0$ for some $i \leq r$. Thus, for each $x \in U$, $x \neq c$, we get some i such that

$$|f_i(x)| < \frac{\sqrt{|f_i(x)|}}{2^i M}.$$

Hence,

$$|f(x)| \leq M \sum_{i=1}^r |f_i(x)| < \sum_{i=1}^r \frac{\sqrt{|f_i(x)|}}{2^i} \leq |f(x)|$$

which is a contradiction. This proves that I_c is not countably generated.

Prime ideals in $C[0, 1]$ Recall that a proper ideal P in a commutative ring R is *prime*, if $st \in P$ implies either $s \in P$ or $t \in P$. Equivalently, the quotient ring R/P is an integral domain (a ring in which $xy = 0$ implies either $x = 0$ or $y = 0$). Prime ideals are more general than maximal ideals in that every maximal ideal is prime; for, an ideal M is maximal if and only if, R/M is a field. The basic method (essentially, the only method) for constructing prime ideals begins with a multiplicatively closed subset S of R containing 1. Then, any ideal P which is maximal with respect to the property that P does not intersect S , must be prime. This is so because if $ab \in P$, $a \notin P$, $b \notin P$, then the ideals $P + (a)$ and $P + (b)$ must intersect S , by the maximality property. Thus, if

$$s = p_1 + ar_1 \in S \cap (P + (a)) \text{ and } t = p_2 + br_2 \in S \cap (P + (b)),$$

then $st = p_1(p_2 + br_2) + p_2ar_1 + abr_1r_2 \in S \cap P$, contradicting the construction of P . Note that, according to Zorn's lemma, such ideals always exist.

In $C[0, 1]$, one can take S to be the set of all polynomial functions on $[0, 1]$ whose leading coefficient is 1. Then, all resulting ideals are prime, but not maximal; for, if I_c is a maximal ideal containing P , then the polynomial $x - c$ is in $S \cap I_c$ and cannot, therefore, belong to P . It is not clear if such P are countably generated. In general, the following question is natural: *Are there any finitely generated prime ideals in $C[0, 1]$?*

A non-prime ideal in $C(0, 1)$ There is a nice way to use the previous result to produce an ideal in the ring of continuous functions on the noncompact interval $(0, 1)$, which is neither prime nor countably generated. Let

$$I = \{f \in C(0, 1) : f(1/n) = 0 \text{ for all but finitely many } n\}.$$

Note that I is indeed an ideal but not a prime ideal. For instance, if we consider some $f \in C(0, 1)$ which vanishes at all $1/2n$ and does not vanish at any $1/(2n + 1)$ and a function $g \in C(0, 1)$ which vanishes at all $1/(2n + 1)$ and does not vanish at any $1/2n$, then neither f nor g are in I whereas $fg \in I$.

We will show that I cannot be countably generated. To begin with, fix disjoint closed intervals

$$K_n := \left[\frac{1}{n+1} + \epsilon_n, \frac{1}{n} \right],$$

for all n .

Suppose, contrary to what we want to prove, that I is generated by a countable set $\{f_1, f_2, \dots\}$ in $C(0, 1)$. For each n , we look at the ring $C(K_n)$ and its maximal ideal $I_{1/n}$ consisting of those functions which vanish at $1/n$. For each n , $I_{1/n}$, as we have shown, is not countably generated. Now, the restrictions $f_i|_{K_n}$ which happen to vanish at $1/n$, form a countable subset in $I_{1/n}$. For each n , pick an element $\phi_n \in I_{1/n}$ which is *not* in the ideal generated by the restrictions $\{f_i|_{K_n} : i \geq 1, f_i(1/n) = 0\}$. As ϕ_n are defined on disjoint intervals, they have a continuous extension ϕ to the whole of $(0, 1)$. So, now we have $\phi \in C(0, 1)$ with $\phi|_{K_n} = \phi_n$. Note that this is possible because the limit point 0 is not in the set $(0, 1)$. Since $\phi_n(1/n) = 0$ for every n , $\phi(1/n) = 0$ for each n ; that is, $\phi \in I = (\{f_1, f_2, \dots\})$. Therefore, by assumption, we may write $\phi = \sum_{i=1}^r g_i f_i$, for suitable $g_i \in C(0, 1)$ and some r . As f_1, \dots, f_r vanish at all but finitely many $1/n$, there is a common N (indeed, infinitely many) so that $f_i(1/N) = 0$ for $i = 1, \dots, r$. Therefore $\phi(1/N) = 0$. However, the fact that

$$\phi_N = \phi|_{K_N} = \sum_{i=1}^r g_i|_{K_N} f_i|_{K_N}$$

contradicts our choice of ϕ_N in $I_{1/N}$. Therefore, I cannot be countably generated.

Summary. Maximal ideals in the ring of continuous functions on the closed interval $[0, 1]$ are not finitely generated. This is well-known. What is not as well-known, but perhaps should be, is the fact that these ideals are not countably generated although the proof is not harder! We prove this here and use the result to produce some non-prime ideals in the ring of continuous functions on the open interval $(0, 1)$ which also cannot be countably generated.

References

1. M. Artin, *Algebra*, 3rd Indian edition, Prentice-Hall of India, New Delhi, 1994.
2. D. S. Dummit & R. M. Foote, *Abstract Algebra*, 2nd ed., John Wiley & Sons, Inc., New York (Asian Edn.), Singapore, 2003.
3. I. N. Herstein, *Topics in Algebra*, 2nd ed., John Wiley & Sons, Inc., New York, 1975.